

# Heart disease and single-vitamin supplementation<sup>1-4</sup>

Maret G Traber

## ABSTRACT

Heart disease is the number one cause of death in the United States and has long been recognized to be multifactorial. A growing body of evidence suggests that not only free radical-mediated reactions but also inflammatory responses play major roles in atherogenesis. Vitamin E has both antioxidant and antiinflammatory properties and is the most widely studied vitamin in clinical trials and thus will be the primary example used in this review. Clinical trials of vitamin E efficacy, in hindsight, have been overly optimistic in their expectation that a vitamin could reverse poor dietary habits and a sedentary lifestyle as well as provide benefit beyond that of pharmaceutical agents in treating heart disease. However, it is also apparent that most Americans do not consume dietary amounts adequate to meet established vitamin E requirements. In response to oxidative stressors, vitamin E can decrease biomarkers of lipid peroxidation, is itself killed, and requires optimal vitamin C status to function most effectively. Thus, adequate vitamin E intakes are clearly needed, but what is adequate for what function has yet to be defined. It is noteworthy that in most trials, biomarkers were not used nor were oxidative stress and lipid peroxidation markers used or plasma vitamin E concentrations measured. *Am J Clin Nutr* 2007;85(suppl):293S-9S.

**KEY WORDS** Vitamin E, folic acid, biomarkers, oxidative stress, carboxyethyl hydroxy chromans

## INTRODUCTION

Heart disease is the number one cause of death in the United States and according to the American Heart Association is also the leading cause of death of women (1). For this presentation, heart diseases include coronary artery disease, hypertensive heart disease, congestive heart failure, peripheral vascular disease, and atherosclerosis, including cerebral artery disease and strokes (2). This review evaluates the use of single-nutrient supplements in the prevention, treatment, or amelioration of heart disease. Vitamin E is the most widely studied with the most clinical trial evidence and thus will be the primary example used.

## HEALTHY DIET

Heart disease has long been recognized to be multifactorial; its onset takes decades in most individuals, and diet is well recognized as an important risk factor. Hu and Willett (3) estimated that 74% of coronary events among nonsmokers might have been prevented by eating a healthy diet (nonhydrogenated unsaturated fats, whole grains, abundant fruit and vegetables, and adequate n-3 fatty acids), maintaining a healthy body weight, exercising

regularly for  $\geq 30$  min/d, and consuming a moderate amount of alcohol ( $>5$  g/d). Nonetheless, such a diet is apparently extremely difficult to achieve for most Americans. The Women's Health Initiative Randomized Controlled Dietary Modification Trial (4) aggressively attempted to change women's diets (reduce total fat intake to 20% of energy, increase intakes of vegetables and fruit to 5 servings/d, and increase grains to  $\geq 6$  servings/d), yet the women in the intervention arm only increased fruit and vegetable intake to 1.1 servings/d and had some success in decreasing fat intake. The diet had no significant effect on coronary heart disease but trends toward greater reductions were observed with lower intakes of saturated fat or *trans* fat or higher intakes of vegetables and fruit (4). Given the poor record of individuals making successful changes to their dietary intakes, the possibility of using individual nutrient supplements to stave off the disorder has been a long-standing quest.

## CHOLESTEROL AS A BIOMARKER OF HEART DISEASE

Reduction of plasma cholesterol by diet to reduce heart disease risk was unsuccessfully attempted for decades. With the advent of drugs that inhibited cholesterol synthesis (HMGCoA reductase inhibitors, or statins), it was possible to demonstrate that serum cholesterol reduction decreased mortality from coronary heart disease (5), suggesting the use of over-the-counter statins to lower serum cholesterol (6). The data from the statin studies have shown that serum cholesterol is an appropriate biomarker for heart disease risk in that decreasing serum cholesterol also decreases heart disease mortality (5). For example, the Food and Drug Administration has permitted a health claim for phyosterols in lowering heart disease risk given that they lower serum cholesterol (7).

Cholesterol, as an example of a biomarker, gives some guidance as to what might be useful in linking a dietary component, a biomarker, and a disease. Thus, cholesterol as a component of the atherosclerotic lesion is not measurable but cholesterol in the serum or specific lipoproteins can readily be measured. The drug

<sup>1</sup> From the Linus Pauling Institute, Oregon State University, Corvallis, OR.

<sup>2</sup> Presented at the conference "Multivitamin/Mineral Supplements and Chronic Disease Prevention," held at the National Institutes of Health, Bethesda, MD, May 15-17, 2006.

<sup>3</sup> Supported by the National Institutes of Health (NIH DK 59576 and ES11536).

<sup>4</sup> Address reprint requests and correspondence to MG Traber, Linus Pauling Institute, Oregon State University, 571 Weniger Hall, Corvallis, OR 97331. E-mail: maret.traber@oregonstate.edu.

studies showed that lowering serum cholesterol resulted in decreased mortality from heart disease. Since then it has only been necessary to show that a dietary component reduces serum cholesterol. Unfortunately, no such biomarkers have been described for antioxidants or many of the vitamins tested.

### ANTIOXIDANTS AND HEART DISEASE

The question as to why elevated plasma cholesterol, specifically LDL cholesterol, causes atherosclerosis led Steinberg et al (7) to posit that modification of LDLs made these plasma cholesterol carriers more susceptible to uptake by scavenger cells (eg, macrophages). The proposed modification of the protein resulted from lipid peroxidation of the particle. This hypothesis became known as the oxidative modification theory and led to the corollary that antioxidants would be beneficial in stopping lipoprotein oxidation.

#### Vitamin E: lipid peroxidation and other functions

Vitamin E functions in vivo as a chain-breaking antioxidant that prevents the propagation of free radical damage in biological membranes (8). Specifically, it is a potent peroxy radical scavenger and especially protects polyunsaturated fatty acids (PUFAs) in membranes and in plasma lipoproteins (9). When lipid hydroperoxides are oxidized to peroxy radicals, these react 1000 times faster with vitamin E than with PUFAs (10). Specifically, the phenolic hydroxyl group of tocopherol reacts with the peroxy radical to form the corresponding lipid hydroperoxide and the tocopheroxyl radical. In this way, vitamin E acts as a chain-breaking antioxidant, preventing further lipid peroxidation.

A growing body of evidence suggests that not only free radical-mediated reactions but also inflammatory responses play a major role in atherogenesis (11–13). Additionally, scavenger cells (eg, monocyte-macrophages) are critically involved in atherogenesis and secrete biologically active mediators: proinflammatory, proatherogenic cytokines, such as interleukin 1 $\beta$  (IL-1 $\beta$ ) and tumor necrosis factor  $\alpha$ , as well as chemokines, such as monocyte chemoattractant protein 1. Both IL-1 $\beta$  and tumor necrosis factor  $\alpha$  stimulate expression of the adhesion molecules vascular adhesion molecule 1 (VCAM-1), intercellular cell adhesion molecule 1 (ICAM-1), and E-selectin (14).

Importantly,  $\alpha$ -tocopherol appears to have potent effects on cellular functions that may modulate heart attack risk (15).  $\alpha$ -Tocopherol can modulate the inflammatory response by inhibiting 5-lipoxygenase, which decreases monocyte IL-1 $\beta$  release (16).  $\alpha$ -Tocopherol also decreases monocyte-endothelial cell adhesion in vitro, which correlates with decreased endothelial cell E-selectin expression (17) and decreased adhesion molecules (ICAM-1 and VCAM-1) induced by oxidized LDL (18, 19). The decreased ICAM-1 expression is mediated by decreased expression of CD11b and VLA-4, possibly by  $\alpha$ -tocopherol inhibiting the activation of nuclear factor  $\kappa$ B (20).  $\alpha$ -Tocopherol inhibits some of these cellular functions through mechanisms mediated by protein kinase C, as shown in model systems such as smooth muscle cell proliferation (21) and platelet aggregation and adhesion (22, 23).  $\alpha$ -Tocopherol supplementation also decreases monocyte superoxide production via inhibition of protein kinase C (20, 24).

Vascular homeostasis is another key function regulated by  $\alpha$ -tocopherol. Normal vascular function requires responsiveness

to nitric oxide.  $\alpha$ -Tocopherol mediates nitric oxide production in endothelial cells (25). In an ex vivo study of rabbit aorta, delivery of  $\alpha$ -tocopherol mediated by plasma phospholipid transfer protein resulted in the maintenance of relaxation in response to acetylcholine (26). In human aortic endothelial cells in culture, vitamin E resulted in a net increase in the production of vasodilator prostanoids (27).

Unfortunately, many of the studies described were carried out in tissue culture or are a result of ex vivo treatments with  $\alpha$ -tocopherol. Very few measurements were made in supplemented humans, so the health benefits of these observations made in tissue culture studies are lacking.

#### Vitamin E supplement interventions to ameliorate heart disease

The first of the clinical trials to test the efficacy of vitamin E in heart attack prevention was in an English population and showed that vitamin E prevented second heart attacks (28), but subsequent larger trials largely did not show vitamin E benefit (29, 30). Some examples of the better-known trials are listed in **Table 1**. Nearly 200 trials using vitamin E supplements were carried out in the 10 y after the first trial, and a recent review and meta-analysis claims that vitamin E has neither benefit nor harm (2). It is noteworthy that in most trials, biomarkers were not used nor were oxidative stress and lipid peroxidation markers or plasma vitamin E concentrations measured.

The literature abounds with basic science showing that vitamin E is an antioxidant and inhibits smooth muscle cell proliferation, platelet adhesion and aggregation, and monocyte endothelial adhesion; it has benefit in animal atherosclerosis models and LDL oxidation, platelet effects, and antiinflammatory effects in humans (as reviewed in reference 37). Thus, it is difficult to argue that the mechanistic aspects of vitamin E's actions are unknown and need further study. Nonetheless, the actual processes that generate the necessity of vitamin E for the human body are unknown, leading to confusion as to why humans require vitamin E.

#### Vitamin E absorption, transport, metabolism, and excretion

Dietary components with vitamin E antioxidant activity include  $\alpha$ -,  $\beta$ -, and  $\gamma$ -tocopherols and -tocotrienols (38). All these molecules have a chromanol ring with a various number of methyl groups and have either a phytyl tail (tocopherols) or an unsaturated tail (tocotrienols). The naturally occurring form of  $\alpha$ -tocopherol is *RRR*- $\alpha$ -tocopherol; chiral carbons are in the *R*-conformation at positions 2, 4', and 8'. Chemical synthesis of  $\alpha$ -tocopherol results in an equal mixture of 8 different stereoisomers (*RRR*, *RSR*, *RRS*, *RSS*, *SRR*, *SSR*, *SRS*, and *SSS*) called *all-rac*- $\alpha$ -tocopherol. The 2 position is critical for in vivo  $\alpha$ -tocopherol activity; only *2R*- $\alpha$ -tocopherol forms meet human vitamin E requirements (38).

The liver preferentially secretes  $\alpha$ -tocopherol into plasma, under control of the  $\alpha$ -tocopherol transfer protein ( $\alpha$ -TTP) as shown in vitamin E-deficient humans with genetic  $\alpha$ -TTP defects (39, 40) and in  $\alpha$ -TTP knockout mice (*Ttpa*<sup>-/-</sup>) (41, 42).  $\alpha$ -Tocopherol is excreted into bile via MDR2 (ABCB4) (43), an ATP-binding cassette transporter that facilitates biliary phospholipid excretion.

Unlike other fat-soluble vitamins, vitamin E is not accumulated in the liver (or any other tissue) much beyond a 2- to 3-fold

**TABLE 1**Examples of intervention trials with vitamin E<sup>1</sup>

Study	Subjects	Dose	Duration	Relative risk (95% CI)	Endpoint	Reference
	<i>n</i>		<i>y</i>			
CHAOS	2002	400 or 800 IU <i>RRR</i> - $\alpha$ -tocopherol	1.4	0.23 (0.11, 0.47)	Nonfatal myocardial infarction	(28)
GISSI	11 324	300 mg <i>dl</i> - $\alpha$ -tocopherol	3.5	0.98 (0.87, 1.10)	Cardiovascular death, nonfatal myocardial infarction, nonfatal stroke	(30)
HOPE	9541	400 mg <i>RRR</i> - $\alpha$ -tocopherol	4.5	1.05 (0.95, 1.16)	Myocardial infarction, stroke, or cardiovascular death	(29)
SPACE	196	800 IU <i>RRR</i> - $\alpha$ -tocopherol	2	0.46 (0.27, 0.78)	Fatal and nonfatal myocardial infarction, fatal and nonfatal ischemic stroke, unstable angina, peripheral vascular disease, excluding sudden death	(31)
PPP	4495	300 mg <i>dl</i> - $\alpha$ -tocopherol	3.6	1.07 (0.74, 1.56)	Cardiovascular death, nonfatal myocardial infarction, nonfatal stroke	(32)
MICRO-HOPE	3654	400 IU <i>RRR</i> - $\alpha$ -tocopheryl acetate	4.5	1.03 (0.88, 1.21)	Myocardial infarction, stroke, or cardiovascular death	(33)
VEAPS	353	400 IU <i>dl</i> - $\alpha$ -tocopherol	3	<i>P</i> = 0.81 for cardiovascular events (14 placebo and 11 vitamin E)	Fatal, nonfatal, or non-Q-wave myocardial infarction, coronary artery bypass graft surgery, percutaneous transluminal coronary angioplasty, unstable angina, transient ischemic attack, cerebrovascular accident	(34)
HOPE-TOO	3994	400 IU <i>RRR</i> - $\alpha$ -tocopheryl acetate	7	1.04 (0.96, 1.14)	Myocardial infarction, stroke, or cardiovascular death	(35)
Women's Health Study	39 876	600 IU <i>RRR</i> - $\alpha$ -tocopherol	10.1	0.93 (0.82, 1.05)	Nonfatal myocardial infarction, nonfatal stroke, or cardiovascular death	(36)

<sup>1</sup> CHAOS, Cambridge Heart Antioxidant Study; GISSI, Gruppo Italiano per lo Studio della Sopravvivenza nell' Infarto Miocardico; HOPE, Heart Outcomes Prevention Evaluation; SPACE, secondary prevention with antioxidants of cardiovascular disease in endstage renal disease; PPP, Primary Prevention Project; VEAPS, Vitamin E Atherosclerosis Prevention Study.

increase above concentrations seen in unsupplemented subjects (44), which suggests that excretion and metabolism are important in preventing hypervitaminosis E. The vitamin E metabolites  $\alpha$ -CEHC (2,5,7,8-trimethyl-2-[2'-carboxyethyl]-6-hydroxychroman) and  $\gamma$ -CEHC (2,7,8-trimethyl-2-[2'-carboxyethyl]-6-hydroxychroman) are derived from  $\alpha$ - and  $\gamma$ -tocopherols, respectively (45). Vitamin Es are metabolized similarly to xenobiotics in that they are metabolized by cytochrome P450s (CYPs), conjugated, and excreted in urine (45) or bile (46). After  $\omega$ -hydroxylation by CYPs,  $\beta$ -oxidation takes place, and CEHCs can then be sulfated or glucuronidated (47–49).

Vitamin E metabolism seems to be a key event in preventing the accumulation of forms of vitamin E other than  $\alpha$ -tocopherol. After administration of an equimolar dose of differently deuterium-labeled  $\alpha$ - and  $\gamma$ -tocopherols to humans, the disappearance rates for  $\gamma$ -tocopherol and  $\alpha$ -CEHC were virtually identical and much faster than for  $\alpha$ -tocopherol (50). These data emphasize that the body preferentially retains  $\alpha$ -tocopherol and that  $\gamma$ -tocopherol is rapidly metabolized and removed from the circulation.

Why CEHC production from  $\gamma$ -tocopherol is so much greater than that from  $\alpha$ -tocopherol is unknown, but studies in mice suggest that hepatic vitamin E regulation depends on

$\alpha$ -tocopherol concentrations. Hepatic Cyp3a protein is correlated with hepatic  $\alpha$ -tocopherol concentrations, even in mice fed diets high in  $\gamma$ -tocopherol (51), and hepatic Cyp3A mRNA is also up-regulated by dietary  $\alpha$ -tocopherol but not by  $\gamma$ -tocotrienol (52). CYP3A4 inhibition in rats by sesame lignans (sesamin and sesaminol) elevates plasma  $\gamma$ -tocopherol concentrations and decreases  $\gamma$ -CEHC concentrations; similar effects were seen in rats fed ketoconazole, a known inhibitor of CYP3A (53). These data suggest that  $\alpha$ -tocopherol stimulates the metabolism and excretion of non- $\alpha$ -tocopherol forms by increasing CYP3A levels and that inhibition of CYP3A decreases the metabolism of  $\gamma$ -tocopherol, as discussed (51). Moreover, these data emphasize that vitamin E concentrations are limited to prevent excess accumulation.

CYP3A is regulated by a xenobiotic nuclear receptor, the pregnane X receptor (PXR) (54). PXR regulates a constellation of genes involved in xenobiotic detoxification (54–56). PXR is a broad-specificity nuclear receptor that binds lipophilic xenobiotics, including various vitamin E forms (57, 58). On ligand binding, PXR, as a heterodimer with the retinoid X receptor, interacts with the promoter region of genes and induces some cytochrome P450 oxidation systems (phase I), conjugation systems (phase II), and transporters (phase III) that effectively clear

xenobiotics from the liver (59).  $\alpha$ -Tocopherol, as a PXR ligand, could alter these PXR-mediated pathways; however, definitive proof of whether  $\alpha$ -tocopherol regulates its own hepatic concentrations through these pathways or by some other mechanisms is currently lacking and is an area of intense investigation.

Reports of adverse effects of vitamin E supplements in humans are so rare that for the Food and Nutrition Board to set the upper tolerance level (UL), data from studies in rats had to be used. The UL for  $\alpha$ -tocopherol is 1000 mg/d (38). However, findings from 2 clinical trials suggested that adverse vitamin E effects do exist. One study was a 3-y, double-blind, placebo-controlled trial of antioxidants (vitamins E and C,  $\beta$ -carotene, and selenium) in 160 subjects taking the drug combination simvastatin and niacin (60, 61). The drugs raised HDL cholesterol less than expected in subjects taking antioxidants (60) and there was an increase in adverse clinical endpoints (61). The other study was the Women's Angiographic Vitamin and Estrogen Trial, a randomized, double-blind trial of 423 postmenopausal women with  $\geq 1$  coronary stenosis at baseline coronary angiography. In postmenopausal women taking hormone replacement therapy, all-cause mortality was higher in women assigned to antioxidant vitamins compared with placebos (hazard ratio: 2.8; 95% CI: 1.1, 7.2;  $P = 0.047$ ) (62). Both simvastatin (63) and estrogen (64) are metabolized by CYP3A4, lending support to the hypothesis that  $\alpha$ -tocopherol in pharmacologic doses stimulates drug metabolism, potentially decreasing drug concentrations in the 2 described trials. Unfortunately, the extent to which pharmacologic or even dietary  $\alpha$ -tocopherol intakes modulate these pathways in vivo has not been determined. Moreover, these data emphasize the necessity for appropriate biomarkers to evaluate vitamin E efficacy and excess.

### Vitamin E as an antioxidant in humans

Vitamin E is depleted in humans during oxidative stress. In cigarette smokers, as a result of their increased oxidative stress (assessed by plasma  $F_2$ -isoprostane concentrations), plasma vitamin E is more rapidly depleted than in nonsmokers (65). Remarkably, smokers with the lowest plasma vitamin C concentrations had the fastest vitamin E disappearance, presumably because vitamin C regenerates vitamin E (10). To test this hypothesis, smokers and nonsmokers were supplemented for 2 wk with placebo or vitamin C (1000 mg/d), then their plasma vitamin E disappearance rates were measured (66). Using a crossover design, the same subjects were tested again on the opposite supplement. Marginal vitamin C status in smokers was associated with increased rates of vitamin E disappearance from plasma [as previously observed (65)], and these rates were normalized by prior vitamin C supplementation (66). Importantly, both  $\alpha$ - and  $\gamma$ -tocopherols were similarly affected by vitamin C status, which suggests that oxidation of the tocopherols is the mechanism for the faster vitamin E disappearance (66).

The studies of the smokers also made it apparent that additional biomarkers of oxidative and nitrosative damage would be useful. One biomarker of nitration reactions is 5-nitro- $\gamma$ -tocopherol (67). It is higher in smokers than in nonsmokers (68), coronary heart disease patients (69), and Alzheimer patients (70). Other biomarkers currently under investigation include ascorbyl-4-hydroxy nonenal (71, 72). In addition to  $F_2$ -isoprostanes (73),  $\beta$ -oxidation metabolites of  $F_2$ -isoprostanes may be a useful marker of whole-body oxidative stress when

measured in 24-h urine collections (74).  $F_2$ -isoprostanes are free radical-mediated breakdown products of arachidonic acid; longer-chain PUFAs that lead to other breakdown products, such as the neuroprostanes (73) or  $F_3$ -isoprostanes (75), may also be useful biomarkers of oxidative stress.

In contrast with smokers and others experiencing chronic oxidative stress, persons who exercise experience bursts of increased reactive oxygen species. For example, when oxidative stress was evaluated in endurance runners during a 50-km race compared with a rest day, both  $F_2$ -isoprostane concentrations and vitamin E disappearance rates were elevated during the race (76). Daily supplementation with vitamin E (400 IU) and vitamin C (1000 mg) for 6 wk before the race prevented the increase in  $F_2$ -isoprostanes but not in markers of inflammation (77). Taken together, these data strongly support the concept that vitamin E is required for its antioxidant properties, specifically as a lipid-soluble antioxidant preventing the propagation of lipid peroxidation, and that inadequate levels would lead to greater oxidative stress. Supplementation with vitamin E would then be predicted to be beneficial for the prevention of oxidative stress and related disorders.

The longest trial to test whether vitamin E supplements would prevent heart disease was the Women's Health Study (36). In that study, 40 000 women aged  $\geq 45$  y were randomly assigned to receive vitamin E (600 IU) or placebo and aspirin or placebo every other day. The study lasted 10 y. A significant 24% reduction in cardiovascular death was largely attributed to fewer sudden deaths in the vitamin E group (38 compared with 51 in the placebo group). Women aged  $\geq 65$  y (10% of study participants) assigned to vitamin E had a significant 26% reduction in major cardiovascular events, a 34% reduction in myocardial infarction, and a 49% reduction in cardiovascular deaths and no reduction in stroke rate. There was no significant effect of vitamin E on total mortality. The only significant adverse effect was an increase in the risk of epistaxis (nosebleeds) (36). It is noteworthy that biomarkers were not used to evaluate vitamin E efficacy in this trial. The authors of the study reported that vitamin E provided no overall benefit and do not support recommending vitamin E supplementation for cardiovascular disease prevention among healthy women (36).

### Dietary intakes of vitamin E

The current Recommended Dietary Allowance (RDA) for vitamin E is 15 mg  $\alpha$ -tocopherol/d for women and men (38). The Estimated Average Requirement is 12 mg  $\alpha$ -tocopherol and is not met by 96% of women and 90% of men (84). Vitamin E is fat soluble so it is found in limited amounts in fruit and vegetables;  $\alpha$ -tocopherol is found in seeds, nuts, and some vegetable oils (eg, almond, sunflower seed, and olive oils) (78, 79). On the basis of vitamin E kinetic studies done with deuterium-labeled vitamin E, the RDA values appear correct (80), and cardiovascular disease risk is lower with higher intakes (81). Thus, we are left with increased risk of heart disease with low vitamin E intakes, a population that consumes insufficient amounts of vitamin E to meet its needs, and conflicting data concerning the use of supplements (82, 83). Clearly, additional studies are needed to evaluate the required amount of vitamin E to provide optimal protection against oxidative stress.

## OTHER DIETARY COMPONENTS

Fish oil is one of the possible foods that could be used as a dietary supplement to decrease the risk of heart disease. A recent meta-analysis suggested a role for fish oil (eicosapentaenoic acid, docosahexaenoic acid) or fish in secondary prevention because significant reductions in total mortality, coronary heart disease death, and sudden death were reported (84). However, as noted for the vitamin E studies, no biomarkers or intermediate clinical endpoints exist that could be used for evaluating other nutrients. Additionally, because fish oils are highly polyunsaturated, they require adequate protection from oxidation, reinforcing the concept that adequate vitamin E amounts be consumed.

Hyperhomocysteinemia is associated with increased risk of heart disease. It is unclear as to whether supplementation with folic acid will decrease heart disease risk in addition to decreasing homocysteine concentrations. The B-Vitamin Treatment Trialists' Collaboration just reviewed the design and statistical power of 12 randomized trials assessing the effects of lowering homocysteine with B-vitamin supplements on risk of cardiovascular disease (85). They concluded that the individual trials may not have involved a sufficient number of vascular events or have lasted long enough to have a good chance on their own to detect reliably plausible effects of homocysteine lowering on cardiovascular disease risk, but the combined analysis of these trials should have adequate power to determine whether lowering homocysteine reduces the risk of cardiovascular events within a few years. Findings from both the HOPE2 trial (86) and NORVIT (87) showed that the combination of vitamin B-6, vitamin B-12, and folic acid decreased plasma homocysteine but had no demonstrable benefit for heart disease-related outcomes.

## CONCLUSION

Several studies reported that vitamin E is associated with decreased chronic disease risk. The Women's Health Study, a 10-y prevention trial in normal, healthy women, found that 600 IU vitamin E decreased overall cardiovascular mortality by 24% and in women over 65 y decreased it by 49% (36). Antioxidant treatment with vitamins E and C slowed atherosclerotic progression in intimal thickness of coronary and carotid arteries in patients with hypercholesterolemia (88) and heart-transplant patients (89). The Cache County Study reported that antioxidant use (vitamin E > 400 IU and vitamin C > 500 mg) was associated with reduced Alzheimer disease prevalence and incidence in the elderly (90). Regular vitamin E supplement use for  $\geq 10$  y was associated with a lower risk of dying from amyotrophic lateral sclerosis (Lou Gehrig disease), a neurodegenerative disease (91). It is therefore not surprising that vitamin E supplements are taken daily by >35 million persons in the United States (92).

Heart disease is such a complex disorder that it is also not surprising that no single remedy has been found to ameliorate it. Dietary and lifestyle changes must be dramatic to achieve heart-healthy goals for most adult Americans. Research on how to achieve and maintain these changes is urgently needed. Biomarkers to assess progress toward these goals are also needed. Although it appears overly optimistic to assume that 1 or 2 nutrients might have benefit, if such nutrients were discovered, it would be important to have a clear idea of adverse effects and the intakes at which these adverse effects become problematic. 

The author had no financial or personal interest in any company or organization sponsoring the research, including advisory board affiliations.

## REFERENCES

1. Thom T, Haase N, Rosamond W, et al. Heart disease and stroke statistics—2006 update. A report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation* 2006; 113:e85–151.
2. Shekelle PG, Morton SC, Jungvig LK, et al. Effect of supplemental vitamin E for the prevention and treatment of cardiovascular disease. *J Gen Intern Med* 2004;19:380–9.
3. Hu FB, Willett WC. Optimal diets for prevention of coronary heart disease. *JAMA* 2002;288:2569–78.
4. Howard BV, Van Horn L, Hsia J, et al. Low-fat dietary pattern and risk of cardiovascular disease: the Women's Health Initiative Randomized Controlled Dietary Modification Trial. *JAMA* 2006;295:655–66.
5. Hebert PR, Gaziano JM, Chan KS, Hennekens CH. Cholesterol lowering with statin drugs, risk of stroke, and total mortality. An overview of randomized trials. *JAMA* 1997;278:313–21.
6. Brass EP, Allen SE, Melin JM. Potential impact on cardiovascular public health of over-the-counter statin availability. *Am J Cardiol* 2006;97: 851–6.
7. Steinberg D, Parthasarathy S, Carew TE, Khoo JC, Witztum JL. Beyond cholesterol. Modifications of low-density lipoprotein that increase its atherogenicity. *N Engl J Med* 1989;320:915–24.
8. Burton GW, Joyce A, Ingold KU. First proof that vitamin E is major lipid-soluble, chain-breaking antioxidant in human blood plasma. *Lancet* 1982;8293:327.
9. Dieber-Rotheneder M, Puhl H, Waeg G, Striegl G, Esterbauer H. Effect of oral supplementation with D-alpha-tocopherol on the vitamin E content of human low density lipoproteins and resistance to oxidation. *J Lipid Res* 1991;32:1325–32.
10. Buettner GR. The pecking order of free radicals and antioxidants: lipid peroxidation, alpha-tocopherol, and ascorbate. *Arch Biochem Biophys* 1993;300:535–43.
11. Diaz MN, Frei B, Vita JA, Keaney JFJ. Antioxidants and atherosclerotic heart disease. *N Engl J Med* 1997;337:408–16.
12. Ross R. Atherosclerosis—an inflammatory disease. *N Engl J Med* 1999; 340:115–26.
13. Chait A, Han CY, Oram JF, Heinecke JW. Thematic review series: the immune system and atherogenesis. Lipoprotein-associated inflammatory proteins: markers or mediators of cardiovascular disease? *J Lipid Res* 2005;46:389–403.
14. Carlos TM, Harlan JM. Leukocyte-endothelial adhesion molecules. *Blood* 1994;84:2068–101.
15. Singh U, Devaraj S, Jialal I. Vitamin E, oxidative stress, and inflammation. *Annu Rev Nutr* 2005;25:151–74.
16. Devaraj S, Jialal I.  $\alpha$ -Tocopherol decreases interleukin-1 $\beta$  release from activated human monocytes by inhibition of 5-lipoxygenase. *Arterioscler Thromb Vasc Biol* 1999;19:1125–33.
17. Faruqi R, de la Motte C, DiCorleto PE. Alpha-tocopherol inhibits agonist-induced monocytic cell adhesion to cultured human endothelial cells. *J Clin Invest* 1994;94:592–600.
18. Cominacini L, Garbin U, Pasini AF, et al. Antioxidants inhibit the expression of intercellular cell adhesion molecule-1 and vascular adhesion molecule-1 induced by oxidized LDL on human umbilical vein endothelial cells. *Free Radic Biol Med* 1997;22:117–27.
19. Martin A, Foxall T, Blumberg JB, Meydani M. Vitamin E inhibits low-density lipoprotein-induced adhesion of monocytes to human aortic endothelial cells in vitro. *Arterioscler Thromb Vasc Biol* 1997;17:429–36.
20. Islam KN, Devaraj S, Jialal I.  $\alpha$ -Tocopherol enrichment of monocytes decreases agonist-induced adhesion to human endothelial cells. *Circulation* 1998;98:2255–61.
21. Azzi A, Boscoboinik D, Marilley D, Ozer NK, Stauble B, Tasinato A. Vitamin E: a sensor and an information transducer of the cell oxidation state. *Am J Clin Nutr* 1995;62(suppl):1337S–46S.
22. Freedman JE, Farhat JH, Loscalzo J, Keaney JFJ.  $\alpha$ -Tocopherol inhibits aggregation of human platelets by a protein kinase C-dependent mechanism. *Circulation* 1996;94:2434–40.
23. Steiner M. Influence of vitamin E on platelet function in humans. *J Am Coll Nutr* 1991;10:466–73.
24. Cachia O, Benna JE, Pedruzzi E, Descomps B, Gougerot-Pocidalo MA,

- Leger CL.  $\alpha$ -Tocopherol inhibits the respiratory burst in human monocytes. Attenuation of p47(phox) membrane translocation and phosphorylation. *J Biol Chem* 1998;273:32801–5.
25. Keaney JF Jr, Simon DI, Freedman JE. Vitamin E and vascular homeostasis: implications for atherosclerosis. *FASEB J* 1999;13:965–75.
  26. Desrumaux C, Deckert V, Athias A, et al. Plasma phospholipid transfer protein prevents vascular endothelium dysfunction by delivering  $\alpha$ -tocopherol to endothelial cells. *FASEB J* 1999;13:883–92.
  27. Wu D, Liu L, Meydani M, Meydani SN. Vitamin E increases production of vasodilator prostanoids in human aortic endothelial cells through opposing effects on cyclooxygenase-2 and phospholipase A2. *J Nutr* 2005;135:1847–53.
  28. Stephens NG, Parsons A, Schofield PM, Kelly F, Cheeseman K, Mitchinson MJ. Randomised controlled trial of vitamin E in patients with coronary disease: Cambridge Heart Antioxidant Study (CHAOS). *Lancet* 1996;347:781–6.
  29. Yusuf S, Dagenais G, Pogue J, Bosch J, Sleight P. Vitamin E supplementation and cardiovascular events in high-risk patients. The Heart Outcomes Prevention Evaluation Study Investigators. *N Engl J Med* 2000;342:154–60.
  30. Gruppo Italiano per lo Studio della Streptochinasi nell'Infarto Miocardico. Dietary supplementation with n-3 polyunsaturated fatty acids and vitamin E after myocardial infarction: results of the GISSI-Prevenzione trial. *Lancet* 1999;354:447–55.
  31. Boaz M, Smetana S, Weinstein T, et al. Secondary prevention with antioxidants of cardiovascular disease in endstage renal disease (SPACE): randomised placebo-controlled trial. *Lancet* 2000;356:1213–8.
  32. de Gaetano G. Low-dose aspirin and vitamin E in people at cardiovascular risk: a randomised trial in general practice. Collaborative Group of the Primary Prevention Project. *Lancet* 2001;357:89–95.
  33. Lonn E, Yusuf S, Hoogwerf B, et al. Effects of vitamin E on cardiovascular and microvascular outcomes in high-risk patients with diabetes: results of the HOPE study and MICRO-HOPE substudy. *Diabetes Care* 2002;25:1919–27.
  34. Hodis HN, Mack WJ, LaBree L, et al. Alpha-tocopherol supplementation in healthy individuals reduces low-density lipoprotein oxidation but not atherosclerosis: the Vitamin E Atherosclerosis Prevention Study (VEAPS). *Circulation* 2002;106:1453–9.
  35. Lonn E, Bosch J, Yusuf S, et al. Effects of long-term vitamin E supplementation on cardiovascular events and cancer: a randomized controlled trial. *JAMA* 2005;293:1338–47.
  36. Lee IM, Cook NR, Gaziano JM, et al. Vitamin E in the primary prevention of cardiovascular disease and cancer: the Women's Health Study: a randomized controlled trial. *JAMA* 2005;294:56–65.
  37. Jialal I, Devaraj S. Scientific evidence to support a vitamin E and heart disease health claim: research needs. *J Nutr* 2005;135:348–53.
  38. Institute of Medicine. Dietary reference intakes for vitamin C, vitamin E, selenium, and carotenoids. Washington, DC: National Academy Press, 2000.
  39. Traber MG, Sokol RJ, Burton GW, et al. Impaired ability of patients with familial isolated vitamin E deficiency to incorporate  $\alpha$ -tocopherol into lipoproteins secreted by the liver. *J Clin Invest* 1990;85:397–407.
  40. Traber MG, Sokol RJ, Kohlschütter A, et al. Impaired discrimination between stereoisomers of  $\alpha$ -tocopherol in patients with familial isolated vitamin E deficiency. *J Lipid Res* 1993;34:201–10.
  41. Terasawa Y, Ladha Z, Leonard SW, et al. Increased atherosclerosis in hyperlipidemic mice deficient in  $\alpha$ -tocopherol transfer protein and vitamin E. *Proc Natl Acad Sci U S A* 2000;97:13830–4.
  42. Leonard SW, Terasawa Y, Farese RV Jr, Traber MG. Incorporation of deuterated *RRR*- or *all-rac*- $\alpha$ -tocopherol into plasma and tissues of  $\alpha$ -tocopherol transfer protein-null mice. *Am J Clin Nutr* 2002;75:555–60.
  43. Mustacich DJ, Shields J, Horton RA, Brown MK, Reed DJ. Biliary secretion of  $\alpha$ -tocopherol and the role of the mdr2 P-glycoprotein in rats and mice. *Arch Biochem Biophys* 1998;350:183–92.
  44. Burton GW, Traber MG, Acuff RV, et al. Human plasma and tissue  $\alpha$ -tocopherol concentrations in response to supplementation with deuterated natural and synthetic vitamin E. *Am J Clin Nutr* 1998;67:669–84.
  45. Brigelius-Flohé R, Traber MG. Vitamin E: function and metabolism. *FASEB J* 1999;13:1145–55.
  46. Kiyose C, Saito H, Kaneko K, et al. Alpha-tocopherol affects the urinary and biliary excretion of 2,7,8-trimethyl-2-(2'-carboxyethyl)-6-hydroxychroman, gamma-tocopherol metabolite, in rats. *Lipids* 2001;36:467–72.
  47. Swanson JE, Ben RN, Burton GW, Parker RS. Urinary excretion of 2,7,8-trimethyl-2-(beta-carboxyethyl)-6-hydroxychroman is a major route of elimination of gamma-tocopherol in humans. *J Lipid Res* 1999;40:665–71.
  48. Stahl W, Graf P, Brigelius-Flohe R, Wechter W, Sies H. Quantification of the  $\alpha$ - and gamma-tocopherol metabolites 2,5,7,8-tetramethyl-2-(2'-carboxyethyl)-6-hydroxychroman and 2,7,8-trimethyl-2-(2'-carboxyethyl)-6-hydroxychroman in human serum. *Anal Biochem* 1999;275:254–9.
  49. Pope SA, Burtin GE, Clayton PT, Madge DJ, Muller DP. Synthesis and analysis of conjugates of the major vitamin E metabolite,  $\alpha$ -CEHC. *Free Radic Biol Med* 2002;33:807–17.
  50. Leonard SW, Paterson E, Atkinson JK, Ramakrishnan R, Cross CE, Traber MG. Studies in humans using deuterium-labeled  $\alpha$ - and  $\gamma$ -tocopherol demonstrate faster plasma  $\gamma$ -tocopherol disappearance and greater  $\gamma$ -metabolite production. *Free Radic Biol Med* 2005;38:857–66.
  51. Traber MG, Siddens LK, Leonard SW, et al.  $\alpha$ -Tocopherol modulates Cyp3a expression, increases  $\gamma$ -CEHC production and limits tissue  $\gamma$ -tocopherol accumulation in mice fed high  $\gamma$ -tocopherol diets. *Free Radic Biol Med* 2005;38:773–85.
  52. Kluth D, Landes N, Pfluger P, et al. Modulation of Cyp3a11 mRNA expression by  $\alpha$ -tocopherol but not gamma-tocotrienol in mice. *Free Radic Biol Med* 2005;38:507–14.
  53. Ikeda S, Tohyama T, Yamashita K. Dietary sesame seed and its lignans inhibit 2,7,8-trimethyl-2-(2'-carboxyethyl)-6-hydroxychroman excretion into urine of rats fed gamma-tocopherol. *J Nutr* 2002;132:961–6.
  54. Lehmann JM, McKee DD, Watson MA, Willson TM, Moore JT, Klierer SA. The human orphan nuclear receptor PXR is activated by compounds that regulate CYP3A4 gene expression and cause drug interactions. *J Clin Invest* 1998;102:1016–23.
  55. Maglich JM, Stoltz CM, Goodwin B, Hawkins-Brown D, Moore JT, Klierer SA. Nuclear pregnane X receptor and constitutive androstane receptor regulate overlapping but distinct sets of genes involved in xenobiotic detoxification. *Mol Pharmacol* 2002;62:638–46.
  56. Rushmore TH, Kong AN. Pharmacogenomics, regulation and signaling pathways of phase I and II drug metabolizing enzymes. *Curr Drug Metab* 2002;3:481–90.
  57. Landes N, Pfluger P, Kluth D, et al. Vitamin E activates gene expression via the pregnane X receptor. *Biochem Pharmacol* 2003;65:269–73.
  58. Zhou C, Tabb MM, Sadatrafiei A, Grun F, Blumberg B. Tocotrienols activate the steroid and xenobiotic receptor, SXR, and selectively regulate expression of its target genes. *Drug Metab Dispos* 2004;32:1075–82.
  59. Dussault I, Yoo HD, Lin M, et al. Identification of an endogenous ligand that activates pregnane X receptor-mediated sterol clearance. *Proc Natl Acad Sci U S A* 2003;100:833–8.
  60. Cheung MC, Zhao XQ, Chait A, Albers JJ, Brown BG. Antioxidant supplements block the response of HDL to simvastatin-niacin therapy in patients with coronary artery disease and low HDL. *Arterioscler Thromb Vasc Biol* 2001;21:1320–6.
  61. Brown BG, Zhao XQ, Chait A, et al. Simvastatin and niacin, antioxidant vitamins, or the combination for the prevention of coronary disease. *N Engl J Med* 2001;345:1583–92.
  62. Waters DD, Alderman EL, Hsia J, et al. Effects of hormone replacement therapy and antioxidant vitamin supplements on coronary atherosclerosis in postmenopausal women: a randomized controlled trial. *JAMA* 2002;288:2432–40.
  63. Williams D, Feely J. Pharmacokinetic-pharmacodynamic drug interactions with HMG-CoA reductase inhibitors. *Clin Pharmacokinet* 2002;41:343–70.
  64. Lee AJ, Kosh JW, Conney AH, Zhu BT. Characterization of the NADPH-dependent metabolism of 17 $\beta$ -estradiol to multiple metabolites by human liver microsomes and selectively expressed human cytochrome P450 3A4 and 3A5. *J Pharmacol Exp Ther* 2001;298:420–32.
  65. Bruno RS, Ramakrishnan R, Montine TJ, Bray TM, Traber MG.  $\alpha$ -Tocopherol disappearance is faster in cigarette smokers and is inversely related to their ascorbic acid status. *Am J Clin Nutr* 2005;81:95–103.
  66. Bruno RS, Leonard SW, Atkinson JK, et al. Faster vitamin E disappearance in smokers is normalized by vitamin C supplementation. *Free Radic Biol Med* 2006;40:689–97.

67. Hensley K, Benaksas EJ, Bolli R, et al. New perspectives on vitamin E: gamma-tocopherol and carboxyethylhydroxychroman metabolites in biology and medicine. *Free Radic Biol Med* 2004;36:1–15.
68. Leonard SW, Bruno RS, Paterson E, et al. 5-Nitro- $\gamma$ -tocopherol increases in human plasma exposed to cigarette smoke in-vitro and in-vivo. *Free Radic Biol Med* 2003;38:813–9.
69. Morton LW, Ward NC, Croft KD, Puddey IB. Evidence for the nitration of gamma-tocopherol in vivo: 5-nitro-gamma-tocopherol is elevated in the plasma of subjects with coronary heart disease. *Biochem J* 2002;364:625–8.
70. Williamson KS, Gabbita SP, Mou S, et al. The nitration product 5-nitro-gamma-tocopherol is increased in the Alzheimer brain. *Nitric Oxide* 2002;6:221–7.
71. Sowell J, Frei B, Stevens JF. Vitamin C conjugates of genotoxic lipid peroxidation products: structural characterization and detection in human plasma. *Proc Natl Acad Sci U S A* 2004;101:17964–9.
72. Sowell J, Conway HM, Bruno RS, Traber MG, Frei B, Stevens JF. Ascorbylated 4-hydroxy-2-nonenal as a potential biomarker of oxidative stress response. *J Chromatogr B Analyt Technol Biomed Life Sci* 2005;827:139–45.
73. Roberts LJ 2nd, Fessel JP. The biochemistry of the isoprostane, neuroprostane, and isofuran pathways of lipid peroxidation. *Chem Phys Lipids* 2004;128:173–86.
74. Davies SS, Zackert W, Luo Y, Cunningham CC, Frisard M, Roberts LJ 2nd. Quantification of dinor, dihydro metabolites of F(2)-isoprostanes in urine by liquid chromatography/tandem mass spectrometry. *Anal Biochem* 2006;348:185–91.
75. Gao L, Yin H, Milne GL, Porter NA, Morrow JD. Formation of F-ring isoprostane-like compounds(F3-isoprostanes) in vivo from eicosapentaenoic acid. *J Biol Chem* 2006;281:14092–9.
76. Mastaloudis A, Leonard SW, Traber MG. Oxidative stress in athletes during extreme endurance exercise. *Free Radic Biol Med* 2001;91:1–22.
77. Mastaloudis A, Morrow JD, Hopkins DW, Deveraj S, Traber MG. Antioxidant supplementation prevents exercise-induced lipid peroxidation, but not inflammation, in ultramarathon runners. *Free Radic Biol Med* 2004;36:1329–41.
78. Maras JE, Bermudez OI, Qiao N, Bakun PJ, Boody-Alter EL, Tucker KL. Intake of alpha-tocopherol is limited among US adults. *J Am Diet Assoc* 2004;104:567–75.
79. Moshfegh A, Goldman J, Cleveland L. What we eat in America, NHANES 2001-2001: usual nutrient intakes from food compared to dietary reference intakes. Washington, DC: US Department of Agriculture, Agricultural Research Service, 2005.
80. Bruno RS, Leonard SW, Park S-I, Zhao Y, Traber MG. Human vitamin E requirements assessed with the use of apples fortified with deuterium-labeled  $\alpha$ -tocopheryl acetate. *Am J Clin Nutr* 2006;83:299–304.
81. Knekt P, Ritz J, Pereira MA, et al. Antioxidant vitamins and coronary heart disease risk: a pooled analysis of 9 cohorts. *Am J Clin Nutr* 2004;80:1508–20.
82. Muntwyler J, Hennekens CH, Manson J, Buring JE, Gaziano JM. Vitamin supplement use in a low-risk population of US male physicians and subsequent cardiovascular mortality. *Arch Intern Med* 2002;162:1472–6.
83. Holmquist C, Larsson S, Wolk A, de Faire U. Multivitamin supplements are inversely associated with risk of myocardial infarction in men and women—Stockholm Heart Epidemiology Program (SHEEP). *J Nutr* 2003;133:2650–4.
84. Harper CR, Jacobson TA. Usefulness of omega-3 fatty acids and the prevention of coronary heart disease. *Am J Cardiol* 2005;96:1521–9.
85. B-Vitamin Treatment Trialists' Collaboration. Homocysteine-lowering trials for prevention of cardiovascular events: a review of the design and power of the large randomized trials. *Am Heart J* 2006;151:282–7.
86. Lonn E, Yusuf S, Arnold MJ, et al. Homocysteine lowering with folic acid and B vitamins in vascular disease. *N Engl J Med* 2006;354:1567–77.
87. Bønaa KH, Njølstad I, Ueland PM, et al. Homocysteine lowering and cardiovascular events after acute myocardial infarction. *N Engl J Med* 2006;354:1578–88.
88. Salonen RM, Nyyssonen K, Kaikkonen J, et al. Six-year effect of combined vitamin C and E supplementation on atherosclerotic progression: the Antioxidant Supplementation in Atherosclerosis Prevention (ASAP) Study. *Circulation* 2003;107:947–53.
89. Fang JC, Kinlay S, Beltrame J, et al. Effect of vitamins C and E on progression of transplant-associated arteriosclerosis: a randomised trial. *Lancet* 2002;359:1108–13.
90. Zandi PP, Anthony JC, Khachaturian AS, et al. Reduced risk of Alzheimer disease in users of antioxidant vitamin supplements: the Cache County Study. *Arch Neurol* 2004;61:82–8.
91. Ascherio A, Weiskopf MG, O'Reilly EJ, et al. Vitamin E intake and risk of amyotrophic lateral sclerosis. *Ann Neurol* 2005;57:104–10.
92. Ford ES, Ajani UA, Mokdad AH. Brief communication: the prevalence of high intake of vitamin E from the use of supplements among U.S. adults. *Ann Intern Med* 2005;143:116–20.