

Culture Expansion in Low-Glucose Conditions Preserves Chondrocyte Differentiation and Enhances Their Subsequent Capacity to Form Cartilage Tissue in Three-Dimensional Culture

Hannah K. Heywood,¹ Giovanna Nalesso,² David A. Lee,^{1,*} and Francesco Dell'Accio^{2,*}

Abstract

Culture conditions that preserve a stable chondrocyte phenotype are desirable in cell-based cartilage repair to maximize efficacy and clinical outcome. This study investigates whether low-glucose conditions will preserve the chondrocyte phenotype during culture expansion. Articular chondrocytes were culture-expanded in media supplemented with either low (1 mM) or high (10 mM) glucose. The metabolic phenotype, reactive oxygen species generation, and mRNA expression of markers of differentiation or catabolism were assessed by reverse-transcription quantitative polymerase chain reaction after four population doublings (PDs) and subsequent tissue formation capacity determined using pellet cultures. Continuous monolayer culture was used to determine the population doubling limit. After expansion in monolayer for four PDs, chondrocytes expanded in low-glucose conditions exhibited higher expression of the differentiation markers SOX9 and COL2A1 and reduced expression of the catabolic metalloproteinase matrix metalloproteinase 13. When chondrocytes expanded in low glucose were cultured in micropellets, they consistently generated more cartilaginous extracellular matrix than those expanded in high glucose, as evaluated by wet weight, sulfated glycosaminoglycan content, and hydroxyproline assay for collagen content. The same pattern was observed whether high or low glucose was used during the pellet culture. During expansion, chondrocytes in high-glucose generated 50% more reactive oxygen species than low-glucose conditions, despite a lower dependence on oxidative phosphorylation for energy. Furthermore low-glucose cells exhibited >30% increased population doubling limit. These data suggests that low-glucose expansion conditions better preserve the expression of differentiation markers by chondrocytes and enhance their subsequent capacity to form cartilage *in vitro*. Therefore, low glucose levels should be considered for the expansion of chondrocytes intended for tissue engineering applications.

Key words: chondrocyte; Crabtree effect; differentiation markers; glucose concentration; monolayer expansion potential; regenerative potential; tissue engineering

Introduction

IT IS ESTIMATED THAT ANNUALLY in the United Kingdom, ~10,000 patients suffer cartilage damage warranting repair.¹ The *ex vivo* expansion of chondrocyte populations via proliferation in monolayer is a fundamental part of cell based cartilage repair strategies, including autologous chondrocyte implantation (ACI), a clinically established technique to repair chronic symptomatic isolated chondral

defects.² To obtain sufficient cell numbers, chondrocytes recovered by biopsy are typically cultured for three to four population doublings before re-implantation at high density to the damaged area.^{2,3} The success of these procedures is dependent on the capacity of the implanted cells to form cartilage once implanted *in vivo*.^{4,5}

Chondrocytes lose their expression of differentiation markers upon serial passaging, a phenomenon described as “dedifferentiation.”^{4,6,7} Importantly, dedifferentiation is

¹Institute of Bioengineering, School of Engineering and Materials Science, Queen Mary University of London, London, United Kingdom.

²Centre for Experimental Medicine and Rheumatology, William Harvey Research Institute, Barts and The London, Queen Mary's School of Medicine and Dentistry, London, United Kingdom.

*These authors share senior authorship.

associated with lost capacity of chondrocytes to regenerate cartilage when implanted *in vivo* and with a worse clinical outcome of ACI both in terms of structural repair and symptoms.^{4,6-9} Specific gene expression profiles, including a robust expression of the extracellular matrix (ECM) constituent collagen type 2, can predict the subsequent regenerative capacity of expanded chondrocytes.^{4,10,11} It is well established that type II collagen and aggrecan are direct transcriptional targets of SOX9.^{12,13} Thus optimal cell processing techniques for ACI should aim to sustain the expression of these genes within expanded cells, without unduly compromising proliferation.

Typical monolayer expansion involves culture for several weeks under atmospheric oxygen conditions in a high glucose media (10–25 mM glucose). By contrast, the native environment is scarce in oxygen and glucose.¹⁴⁻¹⁷ The environment experienced by chondrocytes is critical to their behavior, and it is well known that reduced oxygen conditions, more comparable to the native niche, can support the chondrocyte phenotype and promotes redifferentiation of monolayer expanded chondrocytes.¹⁸⁻²²

Estimates predict that glucose concentrations may reach as low as 1 mM within articular cartilage *in vivo*¹⁶ and chondrocytes appear well adapted to effectively obtain glucose from this extracellular environment.^{16,23,24} Interestingly, oxygen consumption of differentiated articular chondrocytes is determined by glucose availability and is suppressed above a threshold of about 5 mM,²⁵⁻²⁸ indicating that these cells are also metabolically sensitive to an altered glucose environment. Therefore, this study investigates the effect of glucose concentration on the preservation of chondrocyte differentiation markers throughout expansion and their potential to produce a cartilaginous ECM. Furthermore, we test whether any difference induced by expansion in low-glucose conditions is a stable feature or is transient.

Methods

Chondrocyte isolation and culture

Full-depth cartilage was dissected from bovine metacarpal-phalangeal joints of 18- to 24-month-old animals under sterile conditions. Chondrocytes were isolated from the ECM by sequential incubation at 37°C in Dulbecco's modified Eagle's medium (DMEM) supplemented with 16 mM HEPES buffer, 1.6 mM L-glutamine, 120 mg/L ascorbic acid, and 16% (v/v) fetal calf serum (FCS) (DMEM+16% FCS) with 5.7 mg/mL pronase for 1 h followed by incubation in DMEM+16% FCS with 100 U/mL collagenase for 12 h. All reagents were from Sigma-Aldrich (Poole, United Kingdom).

Freshly isolated chondrocytes isolated from at least four animals were seeded into culture flasks at a density of 2×10^4 cells/cm² and cultured under standard conditions of 20% (v/v) O₂ and 5% CO₂ with glucose-free DMEM+10% (v/v) FBS, supplemented with 2 mM L-glutamine, 18 mM HEPES buffer, 88 U/mL penicillin, and 88 mg/mL streptomycin (all from Sigma-Aldrich). Cells obtained from the different animals were not pooled. The DMEM of half the flasks from each donor was supplemented with 10 mM glucose (high-glucose medium) and the remaining half with 1 mM glucose (low-glucose medium). Galactose (9 mM) was added to the low glucose media, ensuring osmotic balance.

Cell proliferation was monitored by manually counting the cells *in situ* daily from representative fields of view from each flask, captured with a microscope-mounted camera. Once four population doublings were achieved, the cells were detached with trypsin-EDTA (Sigma-Aldrich) and resuspended in fresh culture media. A portion of the cell suspension was used immediately for metabolic analysis, the determination of reactive oxygen species (ROS) generation, or tissue-forming potential (pellet culture). Gene expression following expansion for four population doublings was determined by quantitative polymerase chain reaction (qPCR), described below. Additional monolayer cultures were maintained continuously in either low- or high-glucose media for over 200 days to examine long-term proliferation characteristics. For the long-term culture the flasks were passaged as they reached 90% confluence using 1:8 split ratio.

Metabolic analysis

The cellular rates of glycolysis and oxidative phosphorylation were calculated from the rate of lactate release and oligomycin-sensitive oxygen consumption, respectively. A fluorescence-based oxygen biosensor with a 96-well plate format (BD Biosciences, Oxford, United Kingdom) was used to monitor the dissolved oxygen concentration in cell suspensions over time, using a method adapted from the suppliers and described in detail previously.²⁸ The oxygen consumption rate of each sample was calculated from the gradient of the oxygen–time curve and normalized to cell number. Additional cell samples were treated with 2 mg/mL oligomycin (Sigma-Aldrich) to determine the F₁F₀ coupled oxygen consumption rate, representative of oxidative phosphorylation.

The lactate released by the cells in suspension was determined from media samples collected at 30-min intervals, as described in a previous study²⁹ and summarized here. β nicotinamide adenine dinucleotide+ (β NAD) was reconstituted with 0.2 mL of glycine buffer, 0.4 mL of water, and 0.01 mL of lactate dehydrogenase suspension (1000 U/mL, Sigma L3916) per milligram of β NAD (Sigma-Aldrich). Two hundred eighty microliters of the reconstituted β NAD+ reagent was added to 20 μ L of each sample and standards in duplicate with incubation at 37°C for 30 min. The concentration of lactate in the samples was calculated by comparison of the absorbance at 340 nm to lactate standards ranging in concentration from 0 to 5 mM.

ROS generation

Cells were suspended at 1×10^6 cells/mL in phosphate-buffered saline containing 10 μ M dihydroethidium (Invitrogen) and supplemented with either low or high glucose according to the expansion conditions. Cellular fluorescence (320-nm excitation, 590-nm emission) was monitored at 5-min intervals for 30 min using a BMG fluostar optima fluorimeter, during which the cells were incubated at 37°C and mixed by orbital shaking. The rate of fluorescence increase was recorded as a measure of ROS generation and xanthine/xanthine oxidase (100 μ M/10 mU/mL) used as a positive control for superoxide generation.

Gene expression (RNA extraction, reverse transcription, and qPCR)

Gene expression was performed on cells cultured in monolayer to four population doublings (PDs). Total RNA of

monolayer cells was extracted using TRIzol reagent (Invitrogen) according to the manufacturer's instructions, and RNA yield was assessed using a NanoDrop spectrophotometer. Five hundred nanograms of total RNA from each sample was reverse transcribed to cDNA using ThermoScript Reverse transcription kit (Invitrogen) with oligo dT primer according to the manufacturer's instructions. Quantitative PCR was performed using a MX3000P thermal cycler (Stratagene, La Jolla, CA) with 1 μ L of cDNA (or water control) in a 10- μ L final reaction volume using hot-start DNA polymerase (Qiagen, Crawley, United Kingdom) in the presence of SYBR green (Sigma-Aldrich), ROX dye (Invitrogen), and 0.25 μ M of specific primers with melt curve analysis of the final product. Details of the primers are available in Table 1. Gene expression was assessed using a standard curve prepared by serial dilution of cDNA of freshly isolated chondrocytes with normalization to the geometric mean of β -actin and β 2 microglobulin (B2M) controls. Collagen, type X, alpha 1 (COL10A1) and matrix metalloproteinase 13 (MMP13) are expected to have low expression in freshly isolated cells, and therefore the standards were prepared by dilution series of a preliminary PCR amplification.

Pellet cultures

One million cells were centrifuged at 380 g in a 15-mL tube to form a cell pellet and cultured in 4 mL of either low- or high-glucose media (described previously) with loose caps under standard CO₂ incubator conditions for 6 weeks, with medium replacement on alternate days. After this time the wet weight of the cell pellets was recorded, and the pellets were either frozen prior to biochemical analysis or fixed in 4% formaldehyde overnight for histological analysis of ECM constituents. Samples for biochemical analysis were digested with papain as described previously,³⁰ prior to the dimethylmethylene blue assay for quantification of sulfated glycosaminoglycans³¹ and hydroxyproline assay, for estimation of collagen content.³² Fixed samples were embedded in paraffin wax and sectioned to 2 μ m for staining with Alcian blue (pH 2.5) for sulfated glycosaminoglycans or Sirius red for collagen content using standard protocols.

Statistical tests

Data was collected from at least four donor animals, which were maintained separately during culture. Statistical comparisons were made by two-factor analysis of variance (ANOVA) and subsequently two-tail *t*-tests, with Bonferroni correction for multiple comparisons where appropriate.

Results

Expansion in low glucose resulted in a marginally longer population doubling time


Chondrocytes lose their differentiated traits during culture expansion *in vitro*.^{4,6,8} Therefore, the effect of glucose concentration on the rate and extent of chondrocyte expansion in monolayer was determined. Population expansion progressed marginally, but consistently, more slowly in low-glucose media compared with high-glucose cultures (Fig. 1A, B). Low-glucose cultures exhibited an extended lag phase of 5.1 days compared with 4.3 days in high glucose (Fig. 1C). The low-glucose cultures also exhibited an

TABLE 1. PRIMER SEQUENCES USED FOR QUANTITATIVE GENE EXPRESSION ANALYSIS

| Gene | Accession no. | Sense | Antisense | Product size (bp) | Annealing temp (°C) |
|----------------|----------------|-----------------------------------|-----------------------------------|-------------------|---------------------|
| β -actin | NM_173979.3 | AGC AGT CGG TTG GAT CGA GCA | GGG AAG GCA AAG GAC TTC CTG TAA C | 137 | 60 |
| B2M | NM_173893.3 | GGG TGC TAC ATG TCC ATG TTT GAC C | TGC AGA AGA CAC CCA GAT GTT GAT G | 118 | 60 |
| SOX9 | AF_278703.1 | ACT CTG GGC AAG CTC TGG AGA CT | GGC GCG GCT GGT ACT TGT AGT CC | 121 | 60 |
| COL2A1 | NM_001001135.2 | GCA AGA GGC CCC TGC AGG TG | AGC ACA AAG CAC AAG CCA GTA TGT | 155 | 60 |
| Aggrecan | NM_173981.2 | GAT GCT TCT ATC CCA GCC TCC GC | CGG TCC GGG AAG TGG CCG TAA | 125 | 60 |
| MMP13 | NM_174389.2 | AGC ACC CTT CCC ATG ACC TTA TCT | TCC TCA AAG TGA AGG GCT CGGC | 154 | 60 |
| COL10A1 | NM_174634.1 | AAA GGT CTA AGT GGC CCC TTT TGT C | GAG GTT CAT GAC AAA AGC ACC TTG C | 138 | 60 |
| PTGS2 | NM_174445.2 | ACT CAC CAG CCC ACA GAG AAC GT | TCC CTT TGC CTA ACA GGC ATA CCT | 225 | 55 |

B2M, β 2 microglobulin; COL2A1, collagen, type II; MMP13, matrix metalloproteinase 13; COL10A1, collagen, type X, alpha 1; PTGS2, prostaglandin-endoperoxide synthase 2.

FIG. 1. Population expansion kinetics and morphology. Cell morphology at different time points in either 1 mM glucose or 10 mM glucose media (A). Population duplication dynamics: Cells expanded in 1 mM glucose initially proliferated more slowly (B), with an extended lag phase and increased population doubling time compared to 10 mM glucose cultures (C), such that four population doublings were obtained after 13 and 11 days, respectively (D). Data represents the mean \pm SD of four donors, and *p*-values indicate significance level of two-tailed paired *t*-tests. PD, population doubling.



increased population doubling (PD) time of 2 days in comparison with 1.6 days in high glucose, such that four PDs (considered a standard expansion for cell products used in ACI procedures) were achieved by day 13 and day 11, respectively (Fig. 1D). Interestingly, however, replicative potential was consistently greater with long-term expansion in the low-glucose condition, while chondrocytes cultured in high glucose stopped dividing after approximately 30 PDs (Supplementary Fig. S1), a potential indication of cumulative cell stress.^{33,34}

Glucose conditions determine chondrocyte oxygen metabolism and ROS throughout culture expansion

The metabolic phenotype of differentiated chondrocytes is known to be responsive to glucose concentration, with oxidative phosphorylation being suppressed by high-glucose conditions (Crabtree effect).^{25–28} To confirm that low-glucose conditions effectively modified the metabolism of chondrocytes in the present study, rates of glycolysis, aerobic metabolism, and ROS levels were investigated following four PDs, which was considered a relevant time point because it is a typical expansion required for ACI. Both oxidative phosphorylation and glycolysis (Fig. 2A, B) increased following expansion compared to freshly isolated cells, in keeping with

previous reports.²⁹ Chondrocytes exhibited significantly greater oxidative phosphorylation in 1 mM glucose compared with 10 mM glucose, both at day 0 and after four PDs (Fig. 2). We calculate, using an ATP to O₂ ratio of 5 (as reported by Brand *et al.*³⁵) and an ATP to lactate ratio of 1, that the cells expanded in low glucose derived 57% of their ATP from aerobic metabolism, compared with 23% in high glucose (Fig. 2C). Increased ROS generation was observed following monolayer culture expansion compared to day 0 cells, and statistically significantly greater ROS levels were observed following culture in high-glucose than in low-glucose conditions (Fig. 2D), despite a lower dependence on aerobic metabolism for ATP.

The stability of the acquired metabolic phenotype was then investigated. If, after four PDs in low glucose, chondrocytes were switched to high glucose conditions and *vice versa*, the metabolic differences observed between the expansion conditions were mostly reversible (Fig. 3A–D). In particular, chondrocytes expanded in 10 mM glucose and resuspended in 1 mM glucose prior to analysis derived the same proportion of ATP from aerobic metabolism (57%) as continuous culture in 1 mM glucose (Fig. 3C). Again, this switch towards oxidative metabolism on transfer to 1 mM glucose was accompanied with decreasing ROS generation (Fig. 3D).




FIG. 2. Chondrocyte metabolic phenotype is sensitive to glucose level. Oxidative phosphorylation (**A**) and glycolytic activity (**B**) are shown for freshly isolated cells (day 0) and cells expanded in monolayer for four population doublings. Combined, these data indicate metabolic sensitivity to glucose concentration at both day 0 and after expansion, such that cells in 1 mM glucose derived a greater proportion of ATP from aerobic metabolism (**C**). Reactive oxygen species, determined by the rate of increase in dihydroethidium fluorescence, increased with monolayer expansion (**D**), but was significantly lower following expansion for four population doublings in 1 mM glucose compared to 10 mM glucose. Data represents the mean and SD from four donors ($n=4$), and p -values represent two-tailed t -tests with Bonferroni correction for multiple comparisons. ROS, reactive oxygen species; DHE, dihydroethidium.

High-glucose conditions resulted in increased chondrocyte dedifferentiation and irreversibly decreased capacity to form cartilage tissue in three-dimensional culture in vitro

To investigate whether high- or low-glucose conditions during culture expansion affected chondrocyte phenotype, gene expression of key chondrocyte markers was compared at the end of a 4-PD expansion period. As expected,^{4,6} the mRNA expression of the differentiation markers SOX9, collagen, type II, alpha 1 (COL2A1), and aggrecan all decreased with culture expansion. However, after four PDs (the typical expansion length for ACI), COL2A1 and SOX9 were more expressed in chondrocytes that had been expanded in low-glucose conditions (Fig. 4A). In addition, the expression of MMP13, a proteolytic enzyme involved in ECM catabolism, was significantly lower following expansion in low glucose compared to the high-glucose expansion conditions (Fig. 4B). The expression of COL10A1 (a marker of chondrocyte hypertrophic differentiation) and prostaglandin-endoperoxide synthase 2 (PTGS2), involved in the synthesis of prostaglandin E2) were not significantly affected by expansion glucose level.

To investigate whether this difference in the chondrocyte gene expression profile also resulted in increased cartilage formation *in vitro*, chondrocytes expanded in either high- or low-glucose condition for 4 PDs were replated in three-dimensional (3D) pellets, a condition known to promote the formation of cartilage organoids,^{8,36} all of them in high-glucose medium. Consistent with the higher expression of differentiation markers at the end of the expansion phase, cartilage tissue formation in pellet culture was more efficient when the chondrocytes had been expanded in low glucose, exhibiting 48% greater wet weight and significantly greater GAG and collagen per pellet (Figs. 5A–C, 6A) compared to cells expanded in high glucose. Histochemical staining supported the presence of cartilaginous ECM within the pellets, with Alcian blue staining for sulfated GAGs, and Sirius red for fibrillar collagens (Fig. 6). Differences in the GAG and hydroxyproline content of the pellets were not significant (two-way ANOVA) when expressed as percent wet weight, indicating a greater amount of qualitatively similar matrix in the two groups.

In ACI, *in vitro* expanded chondrocytes are implanted into the joint defect where they face different conditions during engraftment and cartilage formation. Therefore, the preservation




FIG. 3. The effect of glucose level on chondrocyte metabolic phenotype is largely reversible. Oxidative phosphorylation (**A**) and glycolytic activity (**B**), proportion of ATP from aerobic metabolism (**C**) and reactive oxygen species (ROS) generation (**D**) are shown for cells expanded in monolayer for four population doublings in either 1 mM or 10 mM glucose. The data compare cell metabolism in the respective expansion media and in cells exposed to short-term (< 1 h) switchover of glucose levels to assess reversibility of metabolic traits developed during expansion. Data represents the mean and SD from four donors ($n = 4$), and p -values represent two-tailed t -tests.


of the differentiation and capacity to form cartilage achieved by expansion in low-glucose conditions must be retained in different conditions to maintain clinical relevance. To test the reversibility of the phenotype and of the cartilage formation capacity, chondrocytes were first expanded in either high or low glucose, but then cultured as pellets in low glucose. Also in this case, chondrocytes expanded in high glucose were associated with poorer cartilage formation in pellet culture under these conditions (Fig. 5A–C). Thus, high-glucose conditions during the expansion phase irreversibly reduced the potential for cartilage formation in pellet culture. However, once in pellet culture, high-glucose conditions favored cartilage formation (greater wet weight, total GAG and collagen, ANOVA $p < 0.01$), in keeping with previous reports.³⁷ To summarize the data so far, during the expansion phase, low-glucose conditions better and irreversibly preserved the expression of chondrocyte differentiation markers and increased subsequent cartilage formation in 3D pellet culture; however, once in pellet culture, high-glucose conditions favored the formation of ECM.

Discussion

Culture expansion is a necessary step in ACI, but it is associated with a loss of the differentiated chondrocyte phenotype known as dedifferentiation,^{6,7} a reduced capacity to

form stable cartilage *in vivo*^{4,12,13,38} and ultimately with a poorer clinical outcome of ACI.^{10,11} In this study we have demonstrated that expanded chondrocytes are metabolically sensitive to glucose concentration, undergoing a transient suppression of oxidative phosphorylation and increased ROS production in high glucose. This was followed by a reduced expression of chondrocyte differentiation markers at the end of the expansion phase compared with low-glucose conditions and an irreversible loss of cartilage formation capacity in 3D pellet culture.

Chondrocytes expanded in low glucose media took 2 days longer to reach 4 PDs (Fig. 1D). Despite this additional time *in vitro* required to reach four PDs, the low-glucose samples exhibited superior chondrogenic potential in comparison to high-glucose conditions. Therefore, the additional culture time seems an acceptable cost considering that enhanced cell quality would reduce treatment failure rates and thus requirement for re-interventions.^{4,10,11} Indeed, in a trial in which retention of chondrocyte differentiation markers was used as a quality control, 11.8% of the patients (6 patients out of 57) could not receive the treatment because the chondrocytes had insufficient retention of phenotypic traits.^{9,39} These patients had already received surgery in order to obtain a cartilage biopsy and start chondrocyte expansion, such that an additional 2 days required to achieve the same



level of cell expansion while preserving chondrocyte quality seemed worthwhile in terms of patient benefit and resources.

It is interesting to note that although the chondrocytes in high-glucose conditions initially proliferated faster, they subsequently stopped dividing at around 30 PDs, whereas chondrocytes in low-glucose conditions still displayed a nearly linear growth curve at 80 PDs (Supplementary Fig. S1). Although this is not relevant to chondrocyte expansion for tissue engineering applications since chondrocytes are expanded for much shorter time, such a reduced population

doubling limit is often a feature of cumulative cell stress,^{33,34,40} which may have biological consequences during the long remodeling that the repair tissue undergoes *in vivo* prior to maturation.⁴¹ For example, oxidative stress may reduce the anabolic response of human articular chondrocytes to insulin-like growth factor-1⁴² or induce a hypertrophic phenotype.^{43,44} Furthermore, the difference in longevity *in vitro* may be taken into account for the expansion of mesenchymal stem cells for which low-glucose conditions may warrant higher expansion capacity.





FIG. 5. The glucose level during the culture expansion phase irreversibly affects subsequent tissue formation in pellet culture. Chondrocytes expanded in 1 mM glucose produced micropellets with greater wet weight (A) and containing more GAG (B) and collagen (C) compared with chondrocytes expanded in 10 mM glucose. Data represents the mean \pm SEM of pellets from eight chondrocyte donors ($n=8$), and *p*-values represent two-tailed paired *t*-tests.

FIG. 6. Gross morphology and histological analysis of cartilage tissue formation in cell pellet cultures. Pellets produced from chondrocytes derived from four donors are illustrated in (A). The left-hand image illustrated pellets formed from chondrocytes that had been monolayer-expanded using 1 mM glucose. The right-hand image shows micromasses that had been monolayer-expanded in 10 mM glucose prior to pellet formation. During the three-dimensional pellet phase, all cell pellets illustrated were cultured in 10 mM glucose. Histological staining using Alcian blue (pH 2.5) and Sirius red indicate the presence of sulfated GAG and collagen in the pellets (B).



In this study, monolayer expansion in 1 mM glucose had favorable effects on phenotypic expression profiles (increased SOX9, COL2A1) and reduced expression of catabolic MMP13 compared to typical 10 mM glucose media. An association between levels of these differentiation markers in monolayer and subsequent cartilage formation in 3D has been examined previously.^{4,11,45} However, existing literature regarding the optimal glucose environment for retention of such phenotypic expression profiles *in vitro* is scarce and predominantly focuses on the effects of short-term exposure to exceptionally high glucose levels, such as found in hyperglycemia. For example, Rosa et al.⁴⁶ examined the effect of media containing 30 mM glucose on human chondrocyte phenotype in comparison to 10 mM glucose. By contrast, this study examines physiological glucose levels compared to typical culture media used in ACI. To put the values used in our study in context, the glucose concentration in synovial fluid is 1.4 mM lower than arterial plasma, which is regulated between 4 and 6 mM in healthy humans^{15,47,48} and is estimated to reach levels as low as ~1 mM deep within the cartilage.^{15,16}

Expansion in low glucose represents an effective, safe, and inexpensive option to increase the quality of repair in ACI. Our findings may be further exploited in alternative tissue engineering practices where the cells are seeded into a scaffold and cultured for a further period *in vitro* (where glucose can be controlled and monitored). Here, our findings indicate an optimal protocol would be a combination of low-glucose conditions during expansion followed by high glucose during the 3D phase.

Two caveats are important in terms of relevance to chondrocyte transplantation. In this study we used bovine chondrocytes, which may experience lower glucose levels *in vivo*⁴⁹ and altered uptake kinetics^{16,50} compared with human chondrocytes such that glucose concentrations cannot be directly extrapolated to human chondrocytes. The 3D pellet culture system we used in this study is not directly vali-

dated to predict efficacy in ACI in humans. Therefore, our results warrant further substantiation in appropriate validated *in vivo* models such as the cartilage formation assay in nude mice described by Dell'Accio et al.^{4,9,39}

In our study, the preservation of the cartilage phenotype and increased cartilage formation associated with expansion in low-glucose conditions was also associated with relatively increased oxidative phosphorylation and lower ROS production measured during the expansion phase (Fig. 2). However, while the effects of glucose concentration on oxygen metabolism and ROS generation were mostly transient (they were not retained following media switch-over, Fig. 3), the loss of cartilage-forming capacity on transfer to pellet culture was persistent (Fig. 4). Thus glucose conditions, specifically in the expansion phase, may induce modifications that condition subsequent cartilage formation in pellet culture. This is consistent with epigenetic modification during proliferation.^{51–53}

Acknowledgments

We thank Em Fitzgerald (Islington, London, United Kingdom) for photography of chondrocyte pellet specimens. This study was funded by the Medical Research Council/Engineering and Physical Sciences Research Council (EPSRC) discipline bridging initiative grant PPA026, EPSRC Platform Grant EP/E046975/1; Human Frontier Science Program Grant RGP0025/2009-C and Arthritis Research U.K. grants 19654 and 19344.

Author Disclosure Statement

No competing financial interests exist.

References

1. National Institute for Health and Clinical Excellence. The use of autologous chondrocyte implantation for the

- treatment of cartilage defects in knee joints. *Technology Appraisal 89*. National Institute for Health and Clinical Excellence: London, 2005.
2. Brittberg M. Autologous chondrocyte transplantation. *Clin Orthop Relat Res*. 1999;366(Suppl):S147–S155.
 3. Jones DG, Peterson L. Autologous chondrocyte implantation. *Instr Course Lect*. 2007;56:429–445.
 4. Dell'Accio F, De Bari C, Luyten FP. Molecular markers predictive of the capacity of expanded human articular chondrocytes to form stable cartilage in vivo. *Arthritis Rheum*. 2001;44:1608–1619.
 5. Dell'Accio F, Vanlauwe J, Bellemans J, et al. Expanded phenotypically stable chondrocytes persist in the repair tissue and contribute to cartilage matrix formation and structural integration in a goat model of autologous chondrocyte implantation. *J Orthop Res*. 2003;21:123–131.
 6. Benya PD, Shaffer JD. Dedifferentiated chondrocytes reexpress the differentiated collagen phenotype when cultured in agarose gels. *Cell*. 1982;30:215–224.
 7. Benya PD, Padilla SR, Nimni ME. Independent regulation of collagen types by chondrocytes during the loss of differentiated function in culture. *Cell*. 1978;15:1313–1321.
 8. Giovannini S, Diaz-Romero J, Aigner T, et al. Population doublings and percentage of S100-positive cells as predictors of in vitro chondrogenicity of expanded human articular chondrocytes. *J Cell Physiol*. 2010;222:411–420.
 9. Vanlauwe J, Saris DB, Victor J, et al. Five-year outcome of characterized chondrocyte implantation versus microfracture for symptomatic cartilage defects of the knee: early treatment matters. *Am J Sports Med*. 2011;39:2566–2574.
 10. Saris DB, Vanlauwe J, Victor J, et al. Treatment of symptomatic cartilage defects of the knee: characterized chondrocyte implantation results in better clinical outcome at 36 months in a randomized trial compared to microfracture. *Am J Sports Med*. 2009;37 Suppl 1: 10S-19S.
 11. Niemeyer P, Pestka JM, Salzmann GM, et al. Influence of cell quality on clinical outcome after autologous chondrocyte implantation. *Am J Sports Med*. 2012;40:556–561.
 12. Han Y, Lefebvre V. L-Sox5 and Sox6 drive expression of the aggrecan gene in cartilage by securing binding of Sox9 to a far-upstream enhancer. *Mol Cell Biol*. 2008;28:4999–5013.
 13. Lefebvre V, Huang W, Harley VR, et al. SOX9 is a potent activator of the chondrocyte-specific enhancer of the pro alpha1(II) collagen gene. *Mol Cell Biol*. 1997;17:2336–2346.
 14. Maroudas A, Bullough P, Swanson SA, et al. The permeability of articular cartilage. *J. Bone Joint Surg Br*. 1968;50:166–177.
 15. Levick JR. Microvascular architecture and exchange in synovial joints. *Microcirculation*. 1995;2:217–233.
 16. Windhaber RA, Wilkins RJ, Meredith D. Functional characterisation of glucose transport in bovine articular chondrocytes. *Pflugers Arch*. 2003;446:572–577.
 17. Zhou S, Cui Z, Urban JPG. Factors influencing the oxygen concentration gradient from the synovial surface of articular cartilage to the cartilage-bone interface: a modeling study. *Arthritis Rheum*. 2004;50:3915–3924.
 18. Grimshaw MJ, Mason RM. Bovine articular chondrocyte function in vitro depends upon oxygen tension. *Osteoarthritis Cartilage*. 2000;8:386–392.
 19. Lafont JE, Talma S, Hopfgarten C, et al. Hypoxia promotes the differentiated human articular chondrocyte phenotype through SOX9-dependent and -independent pathways. *J Biol Chem*. 2008;283:4778–4786.
 20. Murphy CL, Sambanis A. Effect of oxygen tension and alginate encapsulation on restoration of the differentiated phenotype of passaged chondrocytes. *Tissue Eng*. 2001;7:791–803.
 21. Duval E, Leclercq S, Elissalde JM, et al. Hypoxia-inducible factor 1alpha inhibits the fibroblast-like markers type I and type III collagen during hypoxia-induced chondrocyte redifferentiation: hypoxia not only induces type II collagen and aggrecan, but it also inhibits type I and type III collagen in the hypoxia-inducible factor 1alpha-dependent redifferentiation of chondrocytes. *Arthritis Rheum*. 2009;60:3038–3048.
 22. Grimshaw MJ, Mason RM. Modulation of bovine articular chondrocyte gene expression in vitro by oxygen tension. *Osteoarthritis Cartilage*. 2001;9:357–364.
 23. Lee RB, Wilkins RJ, Razaq S, et al. The effect of mechanical stress on cartilage energy metabolism. *Biorheology*. 2002;39:133–143.
 24. Mobasheri A, Vannucci SJ, Bondy CA, et al. Glucose transport and metabolism in chondrocytes: a key to understanding chondrogenesis, skeletal development and cartilage degradation in osteoarthritis. *Histol Histopathol*. 2002;17:1239–1267.
 25. Bywaters EGL. The metabolism of joint tissues. *J Pathol Bacteriol*. 1937;44:247–268.
 26. Otte P. Basic cell metabolism of articular cartilage. Manometric studies. *Z Rheumatol*. 1991;50:304–312.
 27. Heywood HK, Bader DL, Lee DA. Rate of oxygen consumption by isolated articular chondrocytes is sensitive to medium glucose concentration. *J Cell Physiol*. 2006;206:402–410.
 28. Heywood HK, Knight MM, Lee DA. Both superficial and deep zone articular chondrocyte subpopulations exhibit the Crabtree effect but have different basal oxygen consumption rates. *J Cell Physiol*. 2010;223:630–639.
 29. Heywood HK, Lee DA. Monolayer expansion induces an oxidative metabolism and ROS in chondrocytes. *Biochem Biophys Res Commun*. 2008;373:224–229.
 30. Heywood HK, Sembi PK, Lee DA, et al. Cellular utilization determines viability and matrix distribution profiles in chondrocyte-seeded alginate constructs. *Tissue Eng*. 2004;10:1467–1479.
 31. Farndale RW, Buttle DJ, Barrett AJ. Improved quantitation and discrimination of sulphated glycosaminoglycans by use of dimethylmethylene blue. *Biochim Biophys Acta*. 1986;883:173–177.
 32. Kafienah W, Sims TJ. Biochemical methods for the analysis of tissue-engineered cartilage. *Methods Mol Biol*. 2004;238:217–230.
 33. Shawi M, Autexier C. Telomerase, senescence and ageing. *Mech Ageing Dev*. 2008;129:3–10.
 34. Martin JA, Klingelutz AJ, Moussavi-Harami F, et al. Effects of oxidative damage and telomerase activity on human articular cartilage chondrocyte senescence. *J Gerontol A Biol Sci Med Sci*. 2004;59:324–337.
 35. Brand MD, Harper ME, Taylor HC. Control of the effective P/O ratio of oxidative phosphorylation in liver mitochondria and hepatocytes. *Biochem J*. 1993;291(Pt 3):739–748.
 36. Barbero A, Grogan S, Schafer D, et al. Age related changes in human articular chondrocyte yield, proliferation and post-expansion chondrogenic capacity. *Osteoarthritis Cartilage*. 2004;12:476–484.
 37. Heywood HK, Bader DL, Lee DA. Glucose concentration and medium volume influence cell viability and glycosaminoglycan synthesis in chondrocyte-seeded alginate constructs. *Tissue Eng*. 2006;12:3487–3496.

38. Dell'Accio F, De Bari C, Luyten FP. Microenvironment and phenotypic stability specify tissue formation by human articular cartilage-derived cells in vivo. *Exp Cell Res.* 2003; 287:16–27.
39. Saris DB, Vanlauwe J, Victor J, et al. Characterized chondrocyte implantation results in better structural repair when treating symptomatic cartilage defects of the knee in a randomized controlled trial versus microfracture. *Am J Sports Med.* 2008;36:235–246.
40. Moussavi-Harami F, Duwayri Y, Martin JA, et al. Oxygen effects on senescence in chondrocytes and mesenchymal stem cells: consequences for tissue engineering. *Iowa Orthop J.* 2004;24:15–20.
41. Roberts S, Hollander AP, Caterson B, et al. Matrix turnover in human cartilage repair tissue in autologous chondrocyte implantation. *Arthritis Rheum.* 2001;44: 2586–2598.
42. Yin W, Park JI, Loeser RF. Oxidative stress inhibits insulin-like growth factor-I induction of chondrocyte proteoglycan synthesis through differential regulation of phosphatidylinositol 3-Kinase-Akt and MEK-ERK MAPK signaling pathways. *J Biol Chem.* 2009;284:31972–31981.
43. Kishimoto H, Akagi M, Zushi S, et al. Induction of hypertrophic chondrocyte-like phenotypes by oxidized LDL in cultured bovine articular chondrocytes through increase in oxidative stress. *Osteoarthritis Cartilage.* 2010;18:1284–1290.
44. Morita K, Miyamoto T, Fujita N, et al. Reactive oxygen species induce chondrocyte hypertrophy in endochondral ossification. *J Exp Med.* 2007;204:1613–1623.
45. Tew SR, Li Y, Pothacharoen P, et al. Retroviral transduction with SOX9 enhances re-expression of the chondrocyte phenotype in passaged osteoarthritic human articular chondrocytes. *Osteoarthritis Cartilage.* 2005;13:80–89.
46. Rosa SC, Rufino AT, Judas FM, et al. Role of glucose as a modulator of anabolic and catabolic gene expression in normal and osteoarthritic human chondrocytes. *J Cell Biochem.* 2011;112:2813–2824.
47. Ropes MW, Bennett GA, Bauer W. The origin and nature of normal synovial fluid. *J Clin Invest.* 1939;18:351–372.
48. Cryer PE, Davis SN, Shamoon H. Hypoglycemia in diabetes. *Diabetes Care.* 2003;26:1902–1912.
49. Kennedy WL AA, Bechdel SI, Shigley JF. Studies on the composition of bovine blood as influenced by gestation, lactation, and age. *J Dairy Sci.* 1939;22:251–260.
50. Rosa SC, Goncalves J, Judas F, et al. Impaired glucose transporter-1 degradation and increased glucose transport and oxidative stress in response to high glucose in chondrocytes from osteoarthritic versus normal human cartilage. *Arthritis Res Ther.* 2009;11:R80.
51. Sassone-Corsi P. Physiology. When metabolism and epigenetics converge. *Science.* 2013;339:148–150.
52. Dvir-Ginzberg M, Gagarina V, Lee EJ, et al. Regulation of cartilage-specific gene expression in human chondrocytes by SirT1 and nicotinamide phosphoribosyltransferase. *J Biol Chem.* 2008;283:36300–36310.
53. Gabay O, Sanchez C, Dvir-Ginzberg M, et al. Sirtuin 1 enzymatic activity is required for cartilage homeostasis in vivo in a mouse model. *Arthritis Rheum.* 2013;65:159–166.

Address correspondence to:

Hannah K. Heywood, PhD
Institute of Bioengineering
School of Engineering and Materials Science
Queen Mary University of London
Mile End Road
London, E1 4NS
United Kingdom

E-mail: h.k.heywood@qmul.ac.uk

Abbreviations Used

3D = three-dimensional
 ACI = autologous chondrocyte implantation
 ANOVA = analysis of variance
 B2M = β 2 microglobulin
 β NAD = β nicotinamide adenine dinucleotide
 COL10A1 = collagen, type X, alpha 1
 COL2A1 = collagen, type II, alpha 1
 DMEM = Dulbecco's modified Eagle's medium
 ECM = extracellular matrix
 FCS = fetal calf serum
 MMP13 = matrix metalloproteinase 13
 PD = population doubling
 PTGS2 = prostaglandin-endoperoxide synthase 2
 qPCR = quantitative polymerase chain reaction
 ROS = reactive oxygen species