



OPEN ACCESS

EDITED BY
Yongqing Li,
University of Michigan, United States

REVIEWED BY
Xin Li,
Shenzhen Bay Laboratory, China
Guizhen Zhao,
University of Michigan, United States

*CORRESPONDENCE
Wei Huang
huangwei1212520@163.com

Yong Xu
xywyll@aliyun.com

†These authors have contributed
equally to this work and share
the first authorship

SPECIALTY SECTION
This article was submitted to
Comparative Immunology,
a section of the journal
Frontiers in Immunology

RECEIVED 29 June 2022
ACCEPTED 22 August 2022
PUBLISHED 12 September 2022

CITATION
Zhou T, Cheng X, He Y, Xie Y,
Xu F, Xu Y and Huang W (2022)
Function and mechanism of
histone β -hydroxybutyrylation
in health and disease.
Front. Immunol. 13:981285.
doi: 10.3389/fimmu.2022.981285

COPYRIGHT
© 2022 Zhou, Cheng, He, Xie, Xu, Xu
and Huang. This is an open-access
article distributed under the terms of
the [Creative Commons Attribution
License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution
or reproduction in other forums is
permitted, provided the original
author(s) and the copyright owner(s)
are credited and that the original
publication in this journal is cited, in
accordance with accepted academic
practice. No use, distribution or
reproduction is permitted which
does not comply with these terms.

Function and mechanism of histone β -hydroxybutyrylation in health and disease

Tingting Zhou^{1,2,3,4†}, Xi Cheng^{1,2,3,4†}, Yanqiu He^{1,2,3,4},
Yumei Xie^{1,2,3,4}, Fangyuan Xu⁵, Yong Xu^{1,2,3,4*} and Wei Huang^{1,2,3,4*}

¹Department of Endocrinology, Affiliated Hospital of Southwest Medical University, Luzhou, China,

²Metabolism, Metabolic Vascular Diseases Key Laboratory of Sichuan Province, Luzhou, China,

³Sichuan Clinical Research Center for Nephropathy, Luzhou, China, ⁴Cardiovascular and Metabolic Diseases Key Laboratory of Luzhou, Luzhou, China, ⁵Department of Rehabilitation, Affiliated Hospital of Southwest Medical University, Luzhou, China

Histone post-translational modifications (HPTMs) are essential epigenetic mechanisms that affect chromatin-associated nuclear processes without altering the DNA sequence. With the application of mass spectrometry-based proteomics, novel histone lysine acylation, such as propionylation, butyrylation, crotonylation, malonylation, succinylation, glutarylation, and lactoylation have been successively discovered. The emerging diversity of the lysine acylation landscape prompted us to investigate the function and mechanism of these novel HPTMs in health and disease. Recently, it has been reported that β -hydroxybutyrate (BHB), the main component of the ketone body, has various protective roles beyond alternative fuel provision during starvation. Histone lysine β -hydroxybutyrylation (Kbhb) is a novel HPTMs identified by mass spectrometry, which regulates gene transcription in response to carbohydrate restriction or elevated BHB levels *in vivo* and *in vitro*. Recent studies have shown that histone Kbhb is strongly associated with the pathogenesis of metabolic cardiovascular diseases, kidney diseases, tumors, neuropsychiatric disorders, and metabolic diseases suggesting it has different functions from histone acetylation and methylation. This review focuses on the writers, erasers, sites, and underlying functions of histone Kbhb, providing a glimpse into their complex regulation mechanism.

KEYWORDS

β -hydroxybutyrylation, histone post-translational modifications, epigenetics, gene regulation, immune

1 Introduction

In eukaryotes, the core component of nucleosome is histone octamer, which is formed by two copies each of histone H2A, H2B, H3, and H4, whereas histone H1 binds to the linker DNA between nucleosomes (1). Histones amino acid residues, particularly in the tail of H3 and H4, such as lysine, arginine, serine, tyrosine, and threonine, are easily catalyzed by specific modification enzymes or enzyme complexes to form different kinds of histone post-translational modifications (HPTMs) that include methylation (Kme) (2), acetylation (Kac) (3) and phosphorylation (4). Those play an important role by regulating chromatin structure and function in determining and maintaining a specific gene expression pattern without changing the DNA sequence, ultimately affecting various biological events in eukaryotic cells. For example, histone acetylation is associated with cell cycle regulation (5), chromatin structure (6), DNA damage and repair (7), histone methylation is involved in gene repression or activation (8), and histone phosphorylation is a classical modification associated with cell division (4) and apoptosis (9).

Most frequently, HPTMs are analyzed using specific antibodies, however, despite being highly sensitive, these methods depend on the generation of specific antisera and do not allow a thorough analysis of combinations of HPTMs on the same polypeptide. Mass spectrometry (MS) is ideal for the analysis of HPTMs and is becoming the gold standard since it allows the quantification of modifications and combinations (10). Following the development of high-throughput and high-sensitivity MS, until now, more than ten types of novel HPTMs, such as lysine propionylation (Kpr) (11), lysine butylation (Kbu) (11), lysine crotonylation (Kcr) (12), lysine malonylation (Kma) (13), lysine succinylation (Ksucc) (14), lysine glutarylation (Kglu) (15), lysine 2-hydroxyisobutyrylation (Khib) (16), lysine β -hydroxybutyrylation (Kbhb) (17), lysine lactoylation (Kla) (18), lysine isonicotinylation (Kinic) (19), lysine benzoylation (Kbz) (20) and hundreds of modification sites, have been discovered in succession. Mounting evidence suggests that these novel histone marks influence gene regulation and are functionally distinct from the commonly studied HPTMs, thus adding another level of complexity to chromatin biology (21). Aberrant patterns of these novel HPTMs have been implicated in various diseases including cancer, neuropsychiatric disorders and cardiovascular disease (18, 22–24).

Ketone bodies, including acetone, acetoacetate, and β -hydroxybutyrate (BHB), are usually produced through fatty acid metabolism and utilized by extrahepatic tissues, such as heart and brain, especially during states of diminished carbohydrate availability. The failing or diabetic heart relies on ketone bodies as a source of energy (25, 26), and the use of ketogenic diets (KD) has therapeutic effects on cardiovascular homeostasis, metabolic diseases, and tumors (27). Recently,

ketosis and BHB have been linked to covalent modifications, and serve as dynamic regulators of chromatin architecture and gene transcription, including acetylation and methylation, suggesting that BHB has functions beyond serving as an alternative fuel provision during carbohydrate restriction (28, 29).

In a recent landmark study, histone Kbhb marks were discovered by MS proteomics (17). BHB-induced histone Kbhb is a novel epigenetic regulatory mark that links metabolism to gene expression and provides a new avenue for studying chromatin regulation and the various functions of BHB in physiological and pathophysiological processes (30–32). Till now, histone Kbhb has been detected in yeast, flies, mice, and human cells, and a total of 46 histone Kbhb sites have been identified *in vivo* and *in vitro*. Here in this review, the discovery and recent advances of histone Kbhb are described, in addition, the role and mechanism of Kbhb in health and disease are also been discussed.

2 The discovery and recent advances of histone Kbhb

2.1 The discovery of histone Kbhb

Epigenetics and HPTMs regulated by ketogenic process have long been reported in the literature (33). Evidence strongly supports that many protective effects of KD are mediated by inhibition of Class I histone deacetylases (HDACs), most likely by BHB, thereby activating expression of protective genes (34). However, in 2016, Yingming Zhao and co-workers identified that BHB works as a substrate for the histone Kbhb, which then activates gene expression, independent of acetylation. In this study, histone Kbhb marks are dramatically induced in response to elevated BHB levels in cultured cells, and in livers of mice subjected to prolonged fasting or streptozotocin (STZ)-induced diabetic ketoacidosis (17). Further confirmed by cellular experiments, the levels of histone Kbhb were elevated by BHB dose-dependent manner. The BHB itself enables Kbhb of histone lysine in the promoter regions of some starvation-related genes, helping the body to rapidly adjust and adapt to the metabolic changes (17).

As shown in Figure 1, till now, a total of 46 histone Kbhb sites have been detected in four core histones (H3, H4, H2A, and H2B) and linker histone H1 *in vivo* and *in vitro* (17, 35). 44 non-redundant histone Kbhb sites are identified in human and mouse cells (17). A total of 44 histone Kbhb sites were identified, of which 38 were from BHB-treated human HEK293 cells and 26 from livers of fasted or STZ-induced diabetic mice, indicating that Kbhb is a widespread histone mark (17). Interestingly, the histone Kbhb mark was significantly elevated in mouse liver during starvation,




FIGURE 1

Histone Kbhb sites on histones. The basic subunit of chromatin is the nucleosome, which comprises an octamer of two copies of each of the core histone proteins (H2A, H2B, H3 and H4) wrapped by 146 bp of chromosomal DNA. Till now, histone Kbhb have been detected in four core histones (H3, H4, H2A and H2B) and linker histone H1 in yeast, flies, mice, and human cells, and a total of 46 histone Kbhb sites have been identified *in vivo* and *in vitro*. These histone Kbhb modification sites cross-talk influence on other types of HPTMs dynamically in health and disease.

especially in the H3K9bhb mark localized in the gene promoter (17). Subsequently, pathways associated with the disease were extracted from the database for separate analysis and found to be involved in amino acid catabolism, circadian rhythms, redox balance, PPAR signaling pathways, and oxidative phosphorylation, as marked by elevated H3K9bhb.


Evidence from previous MS data suggests that histone Kbhb has been described not only in histone proteins but also in non-histone proteins, particularly in transcription factors and as a key enzyme in metabolism (30, 36). p53 Kbhb attenuates p53 acetylation levels, as well as the transcriptional activity of p53 at canonical p53 target genes by which ketone bodies have oncogenic roles (37). Moreover, Kbhb inhibited the activity of S-adenosyl-L-homocysteine hydrolase (AHCY) and altered the metabolite levels (30). However, this review focuses on the histone Kbhb, non-histone Kbhb are not the scope of this review.

2.2 The process of histone Kbhb

The process of ketone body metabolism and histone Kbhb has been shown in Figure 2. Fasting, exercise, calorie restriction, ketogenic diets, and any other state that induces endogenous BHB production will only produce R-type BHB(R-BHB) (38). However, S-type BHB (S-BHB) itself is not a normal product of human metabolism (39). BHB in driving histone Kbhb is provided by fatty acid β -oxidation during ketogenesis (38). Previous studies believe that BHB are transported in the blood

to tissues where they are converted to acetyl-CoA, a substrate in the first step of the TCA cycle. However, the isotopic metabolic labelling shows that BHB also binds to free CoA and then form β -hydroxybutyryl-CoA (BHB-CoA), a high energy donor for histone Kbhb (40). It is worth noting that there are two potential sources of this chiral PTM, R- β -hydroxybutyrate (R-BHB) and S- β -hydroxybutyl CoA (S-BHB-CoA) (41). However, S-BHB-CoA is a transient intermediate in the last round of β -oxidation of fatty acids (42). According to the current research, we speculate that the acyl-CoA synthase short-chain family member 2 (ACSS2) is the key enzyme regulating this process because many acyl-CoAs are derived from their respective short-chain fatty acids (SCFAs) *via* ACSS2, for example, ACSS2 is thought to be the enzyme that generates crotonyl-CoA from crotonate and butyryl-CoA from butyrate (43, 44). Exhaustion of ACSS2 resulted in reducing levels of histone Kcr, demonstrating that SCFAs such as crotonate may be an endogenous source of non-acylated CoAs (45). Whether ACSS2 is responsible for the conversion of BHB to BHB-CoA has not been confirmed. Thus, the identification of ACSS2 is valuable to further elucidate the chromatin signaling pathways in which they are involved.

Chromatin modifications sensitize the genome to cellular metabolism changes, thereby establishing different functional adaptive states. The epigenome acts as a metabolic sensor, allowing rapid transcriptional adaptation to regulate cellular metabolism (46). BHB produced in fasting induces histone Kbhb, which in turn transcriptionally regulates the starvation



response (17), and acetyl-CoA produced by fatty acid β -oxidation appears to be important for controlling the expression program of genes involved in lipid metabolism (47). Thus, the epigenome lies between cell metabolism and the regulation of cell fate. Notably, the enrichment and removal of these epigenetic marks are also dependent on the production of metabolites, which affect the activity of chromatin modifiers, and the enzymes in histone H3K9hb pathways (40). Histone H3K9hb are regulated by R-BHB in a concentration-dependent manner (17). Meanwhile, 3-Hydroxy-3-Methylglutaryl-CoA Synthase 2 (HMGCS2) and β -hydroxybutyrate dehydrogenase 1 (BDH1), both of which act as key enzymes in ketone body metabolism, also affect the modification of histone H3K9hb. Loss of intestinal HMGCS2 impairs the aggregation of H3K9hb and affects H3K9hb-related metabolic gene programs (48). However, inhibition of BDH1 resulted in the aggregation of BHB and the increase of H3K9hb, which promoted the proliferation of hepatocellular carcinoma stem cells (49). Moreover, metabolic stress induced BHB is a natural endogenous HDAC inhibitor, which induces H3K9ac and H3K4ac (50, 51). These studies suggest that there is an interaction between cell metabolome and epigenome, which translate metabolic sensors, such as R-BHB, HMGCS2, or BDH1 into histone H3K9hb and regulate metabolic reprogramming (52).

2.2.1 Writers and readers

Histone H3K9hb is not only influenced by the key enzymes of ketone body metabolism and metabolic substrates themselves,

but also regulated by “readers”, “writers” and “erasers”. The term “writers” refers to enzymes that promote histone acylation modifications (53). Studies are currently focused on acetylation modifications and three families of histone acetyltransferases (HATs) are predominant: the p300/CREB-binding protein (p300/CBP), myst (Moz, Ybf2, SAS2, and Tip60), and GNAT (GCN5-associated N-acetyltransferase) families (54). The enzyme p300 has been identified as an acyltransferase for acetylation (55), propionylation (11) and crotonylation (56). Further experiments have shown that the Ac-CoA binding pocket of p300 can preferentially accommodate uncharged short-chain acyl-CoAs (57). According to recent studies, the acyltransferase p300 acts as a histone H3K9hb “writer” to catalyze the addition of BHB to lysine, and the level of histone H3K9hb at the H3K9, H3K18, H3K27, H4K8 sites decreases in response to p300 knockdown and addition of its some of these histone H3K9hb sites were more sensitive to p300 knockdown than the corresponding Kac sites (35). Whether there is competition between BHB-CoA and acetyl-CoA with p300 or other mechanisms leading to greater sensitivity of histone H3K9hb to p300 remains to be investigated. p300 is a co-writer of multiple acylation modifications. We speculate whether there are writers that can specifically install histone H3K9hb and cause downstream alterations in different cellular events.

The “reader” proteins may specifically recognize certain HPTMs, which in turn regulate downstream gene transcription and influence cellular events. For example, three

classical histone Kac readers were found to recognize croton groups bound by histone lysine residues. These include bromodomains, YEATS (Yaf9, ENL, AF9, Taf14, and Sas5), and double plant homeodomain finger (DPF) domains (58). In order to further explore the biological effects of histone Kbh, the identification of candidate chromatin-associated proteins that are able to “readers” histone Kbh of lysine residues is put on the agenda. However, till now, none of the unique reader proteins have been discovered for Kbh. Given the large structural differences between histone Kbh and other types of HPTMs, there may be reader proteins that specifically recognise histone Kbh and cause downstream changes in different biological processes. It is probable that multiple mechanisms act synergistically in the regulation of the histone Kbh acylation pathway.

2.2.2 Erasers

Enzymes that removed acylation modifications are known as “erasers”. HDACs are classified into four classes: class I, II, and IV, HDACs are zinc-dependent, while class III HDACs (also known as sirtuins) are NAD-dependent. So far, the “erasers” of histone Kbh are mainly the zinc-dependent HDAC1 and HDAC2 and the NAD-dependent SIRT3 (35), which mediate the removal of Kbh through enzymatic activity. However, the activity of SIRT3 against Kbh is dependent on the site of modification, which differs from its broad class specificity for Kcr modification (59). SIRT3 is distinguished from zinc-dependent HDACs by the fact that SIRT3 shows class-selective activity for histone de- β -hydroxybutyrylation, favoring H3 K4, K9, K18, K23, K27, and H4K16, but not H4 K5, K8, K12 (60). Moreover, lysine R- β -hydroxybutyrylation and lysine S- β -hydroxybutyrylation showed differences in their preference for the deacetylases HDAC3 and SIRT3 (41). S-BHB is stabilized by hydrogen bonding and hydrophobic contacts, but R-BHB has imperfect hydrogen bonding and weak SIRT3 binding (59). In addition, Zhao and co-workers found that HDAC3 prefers lysine R- β -hydroxybutyrylation modification compared to lysine S- β -hydroxybutyrylation *in vitro* (61). The preference of different chiral Kbh for deacetylases implies that we should consider the importance of the chiral structure of Kbh. HDAC1 to HDAC3, SIRT1 and SIRT2 were recently found to have significant de- β -hydroxybutyrylation activity against core histones *in vitro* by high-performance liquid chromatography (HPLC) analysis (35). However, it is interesting that only HDAC1 and HDAC2 are histone Kbh deacetylases in cells. In summary, HDAC1-3, and SIRT1-3 are the enzymes that have been identified for de- β -hydroxybutyrylation. However, most of the different PTMs do not occur on the same lysine residues, suggesting that a large proportion of the PTMs we studied have unique

functions. Whether there are histone Kbh specific “erasers” needs to be further explored.

3 The biological effects of histone Kbh

3.1 Regulation of gene transcription

The involvement of histone Kbh in the transcriptional regulation of genes affects the life activities of organisms by reprogramming the epigenetic landscape on histone and modulating gene expression (See Table 1 for details). Transcription of genes is promoted by histone Kbh of most histones (17). To date, although a total of 46 histone sites have been identified, studies on histone Kbh have focused mainly on H3K9bh. For example, the mechanism of action of BHB antidepressants is related to the activation of brain derived neurotrophic factor (BDNF) by H3K9bh (62). Histone Kbh upregulation of vascular endothelial growth factor (VEGF) expression antagonizes the damage of aortic endothelial cells by diabetes (63). Similarly, BHB epigenetically modifies H3K9 of Foxo1, Ppargc1a (which encodes PGC-1 α), and adiponectin with Kbh, upregulating the expression of these genes (31, 65). Notably, the identified histone Kbh sites include those lysine residues, such as H4K12, H3K4, H3K79, and H3K56, whose acetylation and methylation are essential for chromatin structure and function (23). Therefore, whether Kbh on these histone lysine residues also affects the function of chromatin needs further investigation. In fact, chromatin is not an inert structure, and post-translational modifications of histones alter the chromatin state in response to external changes (23). In a recent study, epigenetic landscape analysis using ChromHMM and k-Means clustering revealed that fasting-induced H3K9bh enrichment was associated with active/permissive chromatin status of H3K4me3 and H3K27ac co-enriched (48).

Chromatin accessibility is a chromatin property that measures the ability of chromatin-binding factors to bind to DNA, and is one of the important indicators to evaluate the structural homeostasis of chromatin (66). Previous studies have found that chromatin accessibility affects gene transcription (67). Recently, histone H3K9bh kept chromatin in an activated state (48), suggesting a close association between histone Kbh and chromatin accessibility. However, to date, no investigations have focused on the relationship between histone Kbh and chromatin accessibility. “Assay for Transposase Accessible Chromatin sequencing” (ATAC-seq) is an efficient and easy to implement protocol to measure chromatin accessibility that has been widely used in multiple applications studying gene regulation (68). Further studies need to explore how histone Kbh regulates chromatin accessibility by ATAC-seq.

TABLE 1 Kbbh *in vivo* and *in vitro* model of human diseases.

Study Type	Model Species	Mode and Dose	Sites	Target Gene/Enzyme	Result	Ref
In vivo	Spatial restraint stress; Dexamethas (DEX)one-induced mouse(20mg/kg)	BHB (300mg/kg/day)by i.p; ketogenic diet	H3K9	BDNF	Antidepressant effect↑, BDNF↑,	(62)
	3-Hit strategy induced mouse and sirt3 ^{-/-} mouse	ketone ester (1 mg/g/day); EMPA (10μg/g/day)	K395 of CS	CS	BDH1↓, NLRP3 inflammasome↓, BNP mRNA↓, Citrate synthase activity↑	(36)
	STZ-induced male SD rats	BHB (160, 200, and 240 mg/kg/day) by i.p	H3K9	VEGF	NO↑, VEGF↑, H3K9bhb↑	(63)
	STZ-induced male SD rats	BHB (160, 200, and 240 mg/kg/day) by i.p	H3K9	MMP-2	Serum creatinine↓, 24 h-urine protein↓, Col IV↓, MMP-2↑, H3K9bhb↑	(64)
	Not described	Fast (24,48hr); ketogenic diet (KD); 10% (w/w)3-butanedioldiet; STZ	K188, K389, K405 of AHCY	AHCY	PanKbbh↑ (liver, kidney), AHCY activity↓, AHCY-Kbbh↑	(30)
	KKAy mice (obese diabetes)	Dapagliflozin drinks freely(0.02mg/ml)	H3K9	Adiponectin	Adiponectin↑, plasma insulin↓, plasma triglyceride↓, MCP-1↓, PAI-1↓, IL-6↓, TNF-α↓	(65)
	DEX(20um)-induced Cortical neurons	BHB (10 mM) overnight	H3K9	BDNF	BDNF ↑, H3K9bhb↑	(62)
In vitro	Isoproterenol and NLRP3 -enriched macrophage induced H9C2 cells	BHB (5 mM)	K395 of CS	CS	COL1α1↓, COL1α2↓, COL3α1↓, CS activity↑, CS-Kbbh↑, Fatty acid uptake↓	(36)
	IL-15-induced T _{mem} cells	BHB (2, 5, 10 mM)	H3K9	Foxo1; Ppargc1a	Tcf7↑, Lef1↑ and Bcl6↓, Pck1↑, Foxo1↑, Ppargc1a↑, PGC-1α↑	(31)
	Not described	BDH1 K.O (knockout)	H3K9	Not described	H3K9bhb↑, JMJD6↑, GREB3↑, GTPBP4↑, NPM1↑, and TIMM23↑	(49)
	Not described	BHB (0.5, 2, 5, 10 mM)	K188 K389 K405 of AHCY	AHCY	PanKbbh↑, AHCY activity↓, AHCY-Kbbh↑	(30)
	3T3-L1Adipocytes	BHB (10mM), Dapagliflozin (10uM)	H3K9	Adiponectin	Adiponectin↑, MCP-1↓, PAI-1↓, IL-6↓	(65)

↑, upregulation; ↓, downregulation. H3K9bhb, histone H3 lysine 9 β-hydroxybutyrylation; BDNF, brain derived neurotrophic factor; EMPA, empagliflozin; CS, citrate synthase; BDH1, β-hydroxybutyrate dehydrogenase 1; NLRP3, nod-like receptor protein 3; BNP, brain natriuretic peptide; STZ, streptozotocin; VEGF, vascular endothelial growth factor; NO, nitric oxide; MMP-2, matrix metalloproteinase-2; Col IV, IV collagen; KD, ketogenic diet; AHCY, S-adenosyl-L-homocysteine hydrolase; MCP-1, monocyte chemoattractant protein 1; PAI-1, plasminogen activator inhibitor-1; IL-6, interleukin-6; TNF-α, tumor necrosis factor-α; DEX, dexamethas; COL1α1, collagen type I α 1 chain; COL1α2, collagen type I α 2 chain; Col3α1, collagen type 3α1 chain; CS-Kbbh, Kbbh of citrate synthase; IL-15, interleukin-15; Foxo1, forkhead homeobox type protein O1; Ppargc1α, peroxisome proliferative activated receptor-γ co-activator 1α; Tcf7, transcription factor 7; Lef1, lymphoid enhancer-binding factor 1; Bcl6, B-cell CLL/lymphoma 6; Pck1, phosphoenolpyruvate carboxykinase 1; JMJD6, jumonji domain containing 6; GTPBP4, GTP binding protein 4; TIMM23, translocase of inner mitochondrial membrane 23 homolog; NPM1, nucleophosmin 1

3.2 Metabolic reprogramming

The metabolic reprogramming of cells enables them to obtain the necessary energy from an energy-deficient environment to maintain continued cell growth and proliferation. More recently, Terranova and co-workers discovered that H3K9bhb is enriched at the proximal promoter of a subset of key genes associated with lipolytic and ketogenic metabolic pathways in the small intestine cells (SICs) crypt during fasting (48). However, intestinal knockdown of HMGCS2 resulted in significant loss of H3K9bhb-related genes, suggesting that local production of BHB is responsible for chromatin reprogramming within the SI crypt. A recent study has revealed that metabolic reprogramming occurs in damaged proximal tubular epithelial cells (PTECs), characterized by a shift from lipolysis to keratolysis (69). In this study, SGLT2 inhibition-associated renal protection was mediated by the elevation of ketone bodies, which suppressed the mechanistic

target of rapamycin complex1 (mTORC1) signaling (as assessed by phosphorylation of the S6 protein) hyperactivation, thereby alleviating DKD (69).

In addition, elevating the levels of histone Kbbh *in vivo* or intracellularly through the KD, starvation, dietary restriction, or direct administration of BHB regulates the activity of key enzymes in metabolism, such as AHCY (30) and citrate synthase (CS) (36), mediating disease physiopathological processes. This implies that, in the future, a KD could be used to increase BHB levels *in vivo* and regulate Kbbh of key enzymes of metabolism and important genes to slow down disease progression. The biological function of histone Kbbh should not be underestimated, as its modification itself up-regulates gene expression and indirectly regulate the level of other modifications, suggesting that the level of histone Kbbh modification can be enhanced in the future through KD or BHB interventions to regulate the transcription of disease-related genes.




FIGURE 3
 Histone Kbbh in cardiovascular diseases. Increasing studies have demonstrated the involvement of BHB and their cognate acylation reactions in cardiovascular diseases. BHB suppresses high salt induced NLRP3 inflammasome activation and then attenuates atherosclerosis and hypertension by GPR109A signaling. BHB activates citrate synthase(CS)via Kbbh modification of K395 and alleviates HFpEF. SIRT3, acts as a histone de-β-hydroxybutyrylation enzyme, inhibits P53 activity through de-Kbbh modification of K120, K170, K319 and reverses BHB-induced vascular senescence. BHB increased histone H3K9bhb, which potentially promote expression of vascular endothelial growth factor (VEGF) and antagonize aortic endothelial injury.

4 Histone Kbbh in disease and therapeutics

Previous studies have shown that KD, BHB treatment, dietary restriction, and fasting enhance histone Kbbh and be involved in the progression of multiple systemic diseases. Next, we will elaborate on the role of histone Kbbh in cardiovascular

diseases, neuropsychiatric disorders, kidney diseases, tumors, and metabolic diseases.

4.1 Cardiovascular diseases

Cardiovascular disease is a major cause of morbidity and mortality in the world. Increasing studies have demonstrated the




FIGURE 4
 Histone Kbbh in neuropsychiatric disorders. Protective effects of BHB on the brain have been extensively researched. BHB has been found to modulate the levels of antioxidant genes, including MnSOD, Foxo3a, and glutathione, and protect neurons from contusion injury by epigenetic mechanism. BHB-reduced H2AK119ub is dependent on activation of L-type calcium channels and elevated intracellular calcium ion concentrations, which in turn activates the Ca²⁺/CaMKII/p-CREB signalling pathway and promotes the occupation of p-CREB and CBP in the brain derived neurotrophic factor(BDNF)promoter, thereby enhancing BDNF expression. However, BHB-increased H3K4me3 is dependent on cAMP/PKA activation. The latest research has shown that the intraperitoneal injection of BHB induced histone H3K9bhb, which promotes the expression of BDNF and alleviate major depressive disorder (MDD).

downregulation is rescued by antidepressants (62). Treatment of spatial restraint stress and dexamethasone-induced MDD mice with BHB showed that BHB alleviated depression and attenuated the down-regulation of BHB and H3K9bhb. The protein level of H3K9bhb but not H3k9ac was increased in a concentration-dependent way with BHB, and BHB reversed the down-regulation of BDNF *in vivo* and *in vitro*. Therefore, H3K9bhb played a crucial role in depression and was induced by BHB and then possibly activated the BDNF gene (82).


In addition, BHB enhanced BDNF expression in hippocampal neurons *in vivo* and *in vitro* by upregulating the transcriptional activation mark H3K4me3 and decreasing the transcriptional repression mark H2AK119ub. BHB-induced changes in H2AK119ub are dependent on the activation of L-type voltage-sensitive Ca²⁺ channels (L-VSCCs) and the elevation of intracellular, which consequently activates the Ca²⁺+/CaMKII/p-CREB signaling pathway and promotes the occupation of p-CREB and CBP in the BDNF promoter, thereby enhancing BDNF expression. However, the increased H3K4me3 by BHB is dependent on the activation of cAMP/PKA (83).


The molecular mechanism of histone Kbh of the BDNF promoter is one of the many potential mechanisms of histone Kbh to protect the brain. Whether the molecular mechanism of dietary restriction to alleviate neurological disease in the KD described above is related to the modification of histone Kbh needs to be further investigated.

4.3 Kidney diseases

While few reports have explored the relationship between histone Kbh and kidney diseases, as summarized in Figure 5,

evidence shows that it has a potentially beneficial effect. Increased serum BHB levels are associated with a higher probability of death in hemodialysis patients (84), but BHB inhibits oxidative stress and may be beneficial in chronic kidney disease (CKD). KD improved type 1 (Akita) and type 2 (db/db mice) diabetic kidney disease (DKD) by increasing the expression of anti-oxidative stress genes, such as Duox1 and SOD1 (85). This effect is thought to be mediated by BHB, which protects neuronal cells from glucose-induced oxidative stress, but this has not been tested in renal cells (85). Treatment of STZ-induced diabetic rats with BHB confirmed that BHB reduced serum creatinine, 24 h-urine protein, and attenuated glomerular morphological changes of the diabetic rats, with IV collagen (COL IV) content, decreased in a concentration-dependent manner. Then, BHB treatment was found to up-regulate renal matrix metalloproteinase-2 (MMP-2) generation of the diabetic rats significantly, while not affecting the increased TGF- β /Smad3 contents. Furthermore, medium and high concentrations of BHB could antagonize diabetic nephropathy injury, potentially by causing H3K9bhb to promote the generation of MMP-2 in diabetic rats (64). Moreover, it was shown that BHB inhibition of HDACs may induce oxidative stress resistance through the expression of FOXO3a and MT2 in mouse kidneys, thus improving the metabolic profile and redox status (50). In time-restricted feeding (TRF)-treated polycystic kidney disease (PKD) rats, mTORC1 activity was found to be inhibited and phosphorylation of its downstream target, phosphorylated S6^{S235/236}, was significantly reduced in cystic wall epithelial cells. Therefore, BHB may slow down the progression of polycystic kidney disease (PKD) in mouse models by inhibiting the mTOR signaling pathway (86).





However, further studies are needed to investigate the overall effect of histone Kbbh on CKD and the underlying mechanisms to find potential therapeutic means for CKD.

4.4 Tumors

Carcinoma cells require large amounts of energy to support their enhanced proliferation. In contrast to non-tumor cells, tumor cells produce energy mainly through glycolysis, which is considered to be an adaptive response (87). As shown in Figure 6, studies suggested that BHB or BHB-induced PTMs may play an important role in the pathogenesis and treatment of tumors. The KD decreased the ratio of IGF to IGF-binding protein 3 (IGFBP3) in mouse serum (88, 89), which is associated with metabolic syndrome and cancer. The link between Kbbh of histones and hepatocellular carcinoma was also identified, and MTA2 interacts with HDAC2/CHD4 and represses BDH1 *via* R-loop transcription, leading to the accumulation of BHB and an increase in H3K9bhb, resulting in a cascade effect that promotes HCC formation and progression. H3K9bhb level was negatively correlated with BDH1. Five target genes were elevated, namely, JMJD6, GREB3, GTPBP4, NPM1, and TIMM23, which led to a poor prognosis in HCC (49). Reducing AHCY activity leads to

reduced invasiveness in breast cancer and glioblastoma cell lines (90). Furthermore, non-histone Kbbh also affects the progression of tumors. Mild inactivation of AHCY activity leads to the late onset of typical disease symptoms and the pathway for the development of hepatocellular carcinoma due to AHCY dysfunction (91). While AHCY is a potential target for tumor therapy, BHB inhibits AHCY through Kbbh modification of the rate-limiting methionine cyclase AHCY K188, K389, K405, site-inhibit AHCY activity (30). The important role of P53, an important tumor suppressor whose activity is fine-tuned by post-translational modifications of P53 in tumors has been demonstrated by numerous experiments in mice in which loss of function of p53 predisposed cells to permanent damage and tumor transformation (92–94). It was recently found that BHB-mediated P53 Kbbh inhibits the level of p53 acetylation and reduces the transcriptional activity of p53 on typical p53 target genes (including p21 and PUMA), thereby reducing the effect of p53 on apoptosis and cell growth (37). The body's energy metabolism and microenvironment are also important conditions for tumor formation and development, enlightening us about the role of histone post-translational modifications, especially histone Kbbh, lactylation, crotonylation, and other HPTMs closely related to metabolism, in tumor metabolism.

4.5 Metabolic diseases

The increasing prevalence of metabolic diseases such as diabetes, hyperlipidemia, obesity, and non-alcoholic fatty liver is a serious burden on human health and socioeconomics. As shown in [Figure 7](#), the studies suggest that BHB or BHB-induced PTMs may play an important role in the pathogenesis and treatment of metabolic diseases and in metabolic reprogramming. Previous studies show that the treatment with SGLT2 inhibitors has been reported to be associated with high levels of serum BHB (95). In obese diabetic mice, dapagliflozin treatment resulted in elevated BHB at plasma and adipose tissue levels which mediated the promotion of apolipoprotein gene expression in adipocytes by H3K9bhb (65). Apolipoprotein has anti-inflammatory and anti-atherogenic effects (96), and BHB may be protective against obese diabetes through the H3K9bhb of the adiponectin gene. Admittedly, diabetes is a classic metabolic disease with complications that involve injury to several organs and systems, such as the kidneys and cardiovascular system (97, 98). BHB inhibits HDAC3 and leads to acetylation of H3K14 in the Claudin-5 promoter, thereby promoting Claudin-5 production and antagonizing diabetes-related cardiac microvascular hyperpermeability (28). Fasting induces an increase in H3K9bhb in small intestinal crypt cells and promotes upregulation of metabolism-related genes that may better respond to metabolic stress (48). In addition, SGLT2 inhibitors also inhibit the mechanistic target of rapamycin complex 1 (mTORC1) signaling overactivation by elevating ketone bodies *in vivo*, thereby attenuating non-proteinuric and proteinuric DKD (69). BHB inhibits adipocyte lipolysis *via* nicotinic acid receptor PUMA-G resulting in alleviation of dyslipidemia (76). Thus, histone Kbhb a novel epigenetic histone regulatory mark that links metabolism to gene expression provides a new way to investigate chromatin regulation and the different functions of BHB in important human pathophysiological states including diabetes, obesity, and hyperlipidemia.

5 Conclusions and prospects

Recent advances in novel HPTMs highlight a natural connection between metabolism and gene transcription. Histone Kbhb is markedly regulated by the level of BHB concentration linking starvation-responsive metabolism and epigenetic regulation. Accumulating studies identified that BHB not only acts as an alternative energy source when glucose supply is inadequate but also regulates a variety of molecular signaling functions (38). BHB and histone Kbhb may influence a wide range of human diseases such as metabolic cardiovascular disease, kidney diseases, tumors, neuropsychiatric disorders, and metabolic diseases. Histone Kbhb, as well as non-histone Kbhb widely exist and they are

highly regulated by energy metabolism. Furthermore, Kbhb was recently discovered to improve the stability of the COVID-19 antibody (99). The complexity of the field of chromatin biology has been enriched with the discovery of an increasing number of novel HPTMs.

The writers and erasers play an important regulatory role in the process of histone Kbhb, acyltransferase p300 was found as a writer which catalyzes histone Kbhb, while HDAC1 to HDAC3, and SIRT1 to SIRT3 are the enzymes that remove Kbhb. These researches uncovered the key regulatory elements of the histone Kbhb pathway, laying the foundation for studying its role in different cellular processes. Until now, what are the main factors that determine the specificity and selectivity of p300, HDACs, or SIRT's enzymatic activity in the process of histone acylations are unclear. The dynamics and ratio of the intracellular concentration of BHB-CoA and acetyl-CoA may be one of the reasons, however, this hypothesis and mechanism need further research to prove. Furthermore, hyperketonemia is known to trigger ketoacidosis, but BHB or KD are used to improve metabolism or epilepsy, how to select the appropriate concentration of BHB without inducing ketoacidosis is still unknown. Therefore, prudent and well-designed clinical and basic experiments are required for a deeper and integrated understanding of histone Kbhb molecular basis in epigenetics and beyond, particularly given the relationship between and other histone acylations, and possible side effects of therapies that provoke ketosis.

Author contributions

WH, YX conceptualized the study, collected the funds, and reviewed the manuscript. T-TZ and XC searched the literature and prepared the manuscript. YQ-H, Y-MX and FY-X were responsible for the final approval of the version to be submitted. All authors read and approved the final manuscript.

Funding

A This work was supported by grants from the National Natural Science Foundation of China (No. 82170834, No.81970676 and No.81800741) and the Office of Science Technology and Talent Work of Luzhou (No. 2020LZXNYDP02 and No. 2021LZXNYD-G01, No. 2019LZXNYDJ40).

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated

organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

- Zhang Y, Sun Z, Jia J, Du T, Zhang N, Tang Y, et al. Overview of histone modification. *Adv Exp Med Biol* (2021) 1283:1–16. doi: 10.1007/978-981-15-8104-5_1
- Greer EL, Shi Y. Histone methylation: a dynamic mark in health, disease and inheritance. *Nat Rev Genet* (2012) 13(5):343–57. doi: 10.1038/nrg3173
- Phillips DM. The presence of acetyl groups of histones. *Biochem J* (1963) 87(2):258–63. doi: 10.1042/bj0870258
- Hans F, Dimitrov S. Histone H3 phosphorylation and cell division. *Oncogene* (2001) 20(24):3021–7. doi: 10.1038/sj.onc.1204326
- Koprinarova M, Schnakenburger M, Diederich M. Role of histone acetylation in cell cycle regulation. *Curr topics med Chem* (2016) 16(7):732–44. doi: 10.2174/1568026615666150825140822
- Grunstein M. Histone acetylation in chromatin structure and transcription. *Nature* (1997) 389(6649):349–52. doi: 10.1038/38664
- Chen CC, Carson JJ, Feser J, Tamburini B, Zabaronick S, Linger J, et al. Acetylated lysine 56 on histone H3 drives chromatin assembly after repair and signals for the completion of repair. *Cell* (2008) 134(2):231–43. doi: 10.1016/j.cell.2008.06.035
- Barski A, Cuddapah S, Cui K, Roh TY, Schones DE, Wang Z, et al. High-resolution profiling of histone methylations in the human genome. *Cell* (2007) 129(4):823–37. doi: 10.1016/j.cell.2007.05.009
- Wen W, Zhu F, Zhang J, Keum YS, Zykova T, Yao K, et al. MST1 promotes apoptosis through phosphorylation of histone H2AX. *J Biol Chem* (2010) 285(50):39108–16. doi: 10.1074/jbc.M110.151753
- Soldi M, Cuomo A, Bonaldi T. Improved bottom-up strategy to efficiently separate hypermodified histone peptides through ultra-HPLC separation on a bench top orbitrap instrument. *Proteomics* (2014) 14(19):2212–25. doi: 10.1002/pmic.201400075
- Chen Y, Sprung R, Tang Y, Ball H, Sangras B, Kim SC, et al. Lysine propionylation and butyrylation are novel post-translational modifications in histones. *Mol Cell Proteomics MCP*. (2007) 6(5):812–9. doi: 10.1074/mcp.M700021-MCP200
- Tan M, Luo H, Lee S, Jin F, Yang JS, Montellier E, et al. Identification of 67 histone marks and histone lysine crotonylation as a new type of histone modification. *Cell* (2011) 146(6):1016–28. doi: 10.1016/j.cell.2011.08.008
- Peng C, Lu Z, Xie Z, Cheng Z, Chen Y, Tan M, et al. The first identification of lysine malonylation substrates and its regulatory enzyme. *Mol Cell Proteomics MCP* (2011) 10(12):M111.012658. doi: 10.1074/mcp.M111.012658
- Zhang Z, Tan M, Xie Z, Dai L, Chen Y, Zhao Y. Identification of lysine succinylation as a new post-translational modification. *Nat Chem Biol* (2011) 7(1):58–63. doi: 10.1038/nchembio.495
- Tan M, Peng C, Anderson KA, Chhoy P, Xie Z, Dai L, et al. Lysine glutarylation is a protein posttranslational modification regulated by SIRT5. *Cell Metab* (2014) 19(4):605–17. doi: 10.1016/j.cmet.2014.03.014
- Dai L, Peng C, Montellier E, Lu Z, Chen Y, Ishii H, et al. Lysine 2-hydroxyisobutyrylation is a widely distributed active histone mark. *Nat Chem Biol* (2014) 10(5):365–70. doi: 10.1038/nchembio.1497
- Xie Z, Zhang D, Chung D, Tang Z, Huang H, Dai L, et al. Metabolic regulation of gene expression by histone lysine β -hydroxybutyrylation. *Mol Cell* (2016) 62(2):194–206. doi: 10.1016/j.molcel.2016.03.036
- Liu Y, Li M, Fan M, Song Y, Yu H, Zhi X, et al. Chromodomain γ -like protein-mediated histone crotonylation regulates stress-induced depressive behaviors. *Biol Psychiatry* (2019) 85(8):635–49. doi: 10.1016/j.biopsych.2018.11.025
- Jiang Y, Li Y, Liu C, Zhang L, Lv D, Weng Y, et al. Isonicotinylation is a histone mark induced by the anti-tuberculosis first-line drug isoniazid. *Nat Commun* (2021) 12(1):5548. doi: 10.1038/s41467-021-25867-y
- Huang H, Zhang D, Wang Y, Perez-Neut M, Han Z, Zheng YG, et al. Lysine benzoylation is a histone mark regulated by SIRT2. *Nat Commun* (2018) 9(1):3374. doi: 10.1038/s41467-018-05567-w
- Smithy J, Sidoli S, Yuan ZF, Coradin M, Bhanu NV, Marchione DM, et al. Characterization of histone acylations links chromatin modifications with metabolism. *Nat Commun* (2017) 8(1):1141. doi: 10.1038/s41467-017-01384-9
- Audia JE, Campbell RM. Histone modifications and cancer. *Cold Spring Harb Perspect Biol* (2016) 8(4):a019521. doi: 10.1101/cshperspect.a019521
- Bannister AJ, Kouzarides T. Regulation of chromatin by histone modifications. *Cell Res* (2011) 21(3):381–95. doi: 10.1038/cr.2011.22
- Abi Khalil C. The emerging role of epigenetics in cardiovascular disease. *Ther Adv Chronic Disease*. (2014) 5(4):178–87. doi: 10.1177/2040622314529325
- Aubert G, Martin OJ, Horton JL, Lai L, Vega RB, Leone TC, et al. The failing heart relies on ketone bodies as a fuel. *Circulation* (2016) 133(8):698–705. doi: 10.1161/CIRCULATIONAHA.115.017355
- Mizuno Y, Harada E, Nakagawa H, Morikawa Y, Shono M, Kugimiya F, et al. The diabetic heart utilizes ketone bodies as an energy source. *Metab: Clin exp* (2017) 77:65–72. doi: 10.1016/j.metabol.2017.08.005
- Zhang J, Jia PP, Liu QL, Cong MH, Gao Y, Shi HP, et al. Low ketolytic enzyme levels in tumors predict ketogenic diet responses in cancer cell lines *in vitro* and *in vivo*. *J Lipid Res* (2018) 59(4):625–34. doi: 10.1194/jlr.M082040
- Li B, Yu Y, Liu K, Zhang Y, Geng Q, Zhang F, et al. β -hydroxybutyrate inhibits histone deacetylase 3 to promote claudin-5 generation and attenuate cardiac microvascular hyperpermeability in diabetes. *Diabetologia* (2021) 64(1):226–39. doi: 10.1007/s00125-020-05305-2
- Mierziak J, Wojtasik W, Kulma A, Dziadas M, Kostyn K, Dymińska L, et al. 3-hydroxybutyrate is active compound in flax that upregulates genes involved in DNA methylation. *Int J Mol Sci* (2020) 21(8):2887. doi: 10.3390/ijms21082887
- Koronowski KB, Greco CM, Huang H, Kim JK, Fribourgh JL, Crosby P, et al. Ketogenesis impact on liver metabolism revealed by proteomics of lysine β -hydroxybutyrylation. *Cell Rep* (2021) 36(5):109487. doi: 10.1016/j.celrep.2021.109487
- Zhang H, Tang K, Ma J, Zhou L, Liu J, Zeng L, et al. Ketogenesis-generated β -hydroxybutyrate is an epigenetic regulator of CD8(+) T-cell memory development. *Nat Cell Biol* (2020) 22(1):18–25. doi: 10.1038/s41556-019-0440-0
- Puchalska P, Crawford PA. Metabolic and signaling roles of ketone bodies in health and disease. *Annu Rev Nutr* (2021) 41:49–77. doi: 10.1146/annurev-nutr-111120-111518
- Dąbek A, Wojtala M, Pirola L, Balcerczyk A. Modulation of cellular biochemistry, epigenetics and metabolomics by ketone bodies. implications of the ketogenic diet in the physiology of the organism and pathological states. *Nutrients* (2020) 12(3):788. doi: 10.3390/nu12030788
- Kong G, Huang Z, Ji W, Wang X, Liu J, Wu X, et al. The ketone metabolite β -hydroxybutyrate attenuates oxidative stress in spinal cord injury by suppression of class I histone deacetylases. *J Neurotrauma* (2017) 34(18):2645–55. doi: 10.1089/neu.2017.5192
- Huang H, Zhang D, Weng Y, Delaney K, Tang Z, Yan C, et al. The regulatory enzymes and protein substrates for the lysine β -hydroxybutyrylation pathway. *Sci Adv* (2021) 7(9):eabe2771. doi: 10.1126/sciadv.abe2771
- Deng Y, Xie M, Li Q, Xu X, Ou W, Zhang Y, et al. Targeting mitochondria-inflammation circuit by β -hydroxybutyrate mitigates HfPpEF. *Circ Res* (2021) 128(2):232–45. doi: 10.1161/CIRCRESAHA.120.317933
- Liu K, Li F, Sun Q, Lin N, Han H, You K, et al. p53 β -hydroxybutyrylation attenuates p53 activity. *Cell Death Dis* (2019) 10(3):243. doi: 10.1038/s41419-019-1463-y
- Newman JC, Verdin E. β -hydroxybutyrate: A signaling metabolite. *Annu Rev Nutr* (2017) 37:51–76. doi: 10.1146/annurev-nutr-071816-064916
- Scofield RF, Brady PS, Schumann WC, Kumaran K, Ohgaku S, Margolis JM, et al. On the lack of formation of l-(+)-3-hydroxybutyrate by liver. *Arch Biochem biophys* (1982) 214(1):268–72. doi: 10.1016/0003-9861(82)90030-3
- Sabari BR, Zhang D, Allis CD, Zhao Y. Metabolic regulation of gene expression through histone acylations. *Nat Rev Mol Cell Biol* (2017) 18(2):90–101. doi: 10.1038/nrm.2016.140

41. Moreno-Yruela C, Bæk M, Monda F, Olsen CA. Chiral posttranslational modification to lysine ϵ -amino groups. *Accounts Chem Res* (2022) 55(10):1456–66. doi: 10.1021/acs.accounts.2c00115
42. Lincoln BC, Des Rosiers C, Brunengraber H. Metabolism of s -3-hydroxybutyrate in the perfused rat liver. *Arch Biochem Biophys* (1987) 259(1):149–56. doi: 10.1016/0003-9861(87)90480-2
43. Hao F, Tian M, Zhang X, Jin X, Jiang Y, Sun X, et al. Butyrate enhances CPT1A activity to promote fatty acid oxidation and iTreg differentiation. *Proc Natl Acad Sci U.S.A.* (2021) 118(22):e2014681118. doi: 10.1073/pnas.2014681118
44. Martinez-Moreno JM, Fontecha-Barriuso M, Martin-Sanchez D, Guerrero-Mauvecin J, Goma-Garcés E, Fernandez-Fernandez B, et al. Epigenetic modifiers as potential therapeutic targets in diabetic kidney disease. *Int J Mol Sci* (2020) 21(11):4113. doi: 10.3390/ijms21114113
45. Sabari BR, Tang Z, Huang H, Yong-Gonzalez V, Molina H, Kong HE, et al. Intracellular crotonyl-CoA stimulates transcription through p300-catalyzed histone crotonylation. *Mol Cell* (2015) 58(2):203–15. doi: 10.1016/j.molcel.2015.02.029
46. Rabhi N, Hannou SA, Froguel P, Annicotte JS. Cofactors as metabolic sensors driving cell adaptation in physiology and disease. *Front Endocrinol* (2017) 8:304. doi: 10.3389/fendo.2017.00304
47. McDonnell E, Crown SB, Fox DB, Kitir B, Ilkayeva OR, Olsen CA, et al. Lipids reprogram metabolism to become a major carbon source for histone acetylation. *Cell Rep* (2016) 17(6):1463–72. doi: 10.1016/j.celrep.2016.10.012
48. Terranova CJ, Stemler KM, Barrodia P, Jeter-Jones SL, Ge Z, de la Cruz Bonilla M, et al. Reprogramming of H3K9bhb at regulatory elements is a key feature of fasting in the small intestine. *Cell Rep* (2021) 37(8):110044. doi: 10.1016/j.celrep.2021.110044
49. Zhang H, Chang Z, Qin LN, Liang B, Han JX, Qiao KL, et al. MTA2 triggered r-loop trans-regulates BDH1-mediated β -hydroxybutyrylation and potentiates propagation of hepatocellular carcinoma stem cells. *Signal Transduct Target Ther* (2021) 6(1):135. doi: 10.1038/s41392-021-00464-z
50. Shimazu T, Hirschev MD, Newman J, He W, Shirakawa K, Le Moan N, et al. Suppression of oxidative stress by β -hydroxybutyrate, an endogenous histone deacetylase inhibitor. *Sci (New York NY)* (2013) 339(6116):211–4. doi: 10.1126/science.1227166
51. Newman JC, Verdin E. Ketone bodies as signaling metabolites. *Trends Endocrinol Metab: TEM*. (2014) 25(1):42–52. doi: 10.1016/j.tem.2013.09.002
52. Boon R. Metabolic fuel for epigenetic: Nuclear production meets local consumption. *Front Genet* (2021) 12:768996. doi: 10.3389/fgene.2021.768996
53. Zhao S, Zhang X, Li H. Beyond histone acetylation-writing and erasing histone acylations. *Curr Opin Struct Biol* (2018) 53:169–77. doi: 10.1016/j.sbi.2018.10.001
54. Roth SY, Denu JM, Allis CD. Histone acetyltransferases. *Annu Rev Biochem* (2001) 70:81–120. doi: 10.1146/annurev.biochem.70.1.81
55. Dancy BM, Cole PA. Protein lysine acetylation by p300/CBP. *Chem Rev* (2015) 115(6):2419–52. doi: 10.1021/cr500452k
56. Huang H, Wang DL, Zhao Y. Quantitative crotonylome analysis expands the roles of p300 in the regulation of lysine crotonylation pathway. *Proteomics* (2018) 18(15):e1700230. doi: 10.1002/pmic.201700230
57. Kaczmarek Z, Ortega E, Goudarzi A, Huang H, Kim S, Márquez JA, et al. Structure of p300 in complex with acyl-CoA variants. *Nat Chem Biol* (2017) 13(1):21–9. doi: 10.1038/nchembio.2217
58. Ntorla A, Burgoyne JR. The regulation and function of histone crotonylation. *Front Cell Dev Biol* (2021) 9:624914. doi: 10.3389/fcell.2021.624914
59. Zhang X, Cao R, Niu J, Yang S, Ma H, Zhao S, et al. Molecular basis for hierarchical histone de- β -hydroxybutyrylation by SIRT3. *Cell Discov* (2019) 5:35. doi: 10.1038/s41421-019-0103-0
60. Abmayr SM, Workman JL. Histone lysine de- β -hydroxybutyrylation by SIRT3. *Cell Res* (2019) 29(9):694–5. doi: 10.1038/s41422-019-0211-2
61. Moreno-Yruela C, Zhang D, Wei W, Bæk M, Liu W, Gao J, et al. Class I histone deacetylases (HDAC1-3) are histone lysine deacetylases. *Sci Adv* (2022) 8(3):eabi6696. doi: 10.1126/sciadv.abi6696
62. Chen L, Miao Z, Xu X. β -hydroxybutyrate alleviates depressive behaviors in mice possibly by increasing the histone3-lysine9- β -hydroxybutyrylation. *Biochem Biophys Res Commun* (2017) 490(2):117–22. doi: 10.1016/j.bbrc.2017.05.184
63. Wu X, Miao D, Liu Z, Liu K, Zhang B, Li J, et al. β -hydroxybutyrate antagonizes aortic endothelial injury by promoting generation of VEGF in diabetic rats. *Tissue Cell* (2020) 64:101345. doi: 10.1016/j.tice.2020.101345
64. Luo W, Yu Y, Wang H, Liu K, Wang Y, Huang M, et al. Up-regulation of MMP-2 by histone H3K9 β -hydroxybutyrylation to antagonize glomerulosclerosis in diabetic rat. *Acta Diabetol* (2020) 57(12):1501–9. doi: 10.1007/s00592-020-01552-2
65. Nishitani S, Fukuhara A, Shin J, Okuno Y, Otsuki M, Shimomura I. Metabolomic and microarray analyses of adipose tissue of dapagliflozin-treated mice, and effects of 3-hydroxybutyrate on induction of adiponectin in adipocytes. *Sci Rep* (2018) 8(1):8805. doi: 10.1038/s41598-018-27181-y
66. Tolsma TO, Hansen JC. Post-translational modifications and chromatin dynamics. *Essays Biochem* (2019) 63(1):89–96. doi: 10.1042/ebc20180067
67. Zhang R, Erler J, Langowski J. Histone acetylation regulates chromatin accessibility: Role of H4K16 in inter-nucleosome interaction. *Biophys J* (2017) 112(3):450–9. doi: 10.1016/j.bpj.2016.11.015
68. Buenostro JD, Giresi PG, Zaba LC, Chang HY, Greenleaf WJ. Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position. *Nat Methods* (2013) 10(12):1213–8. doi: 10.1038/nmeth.2688
69. Tomita I, Kume S, Sugahara S, Osawa N, Yamahara K, Yasuda-Yamahara M, et al. SGLT2 inhibition mediates protection from diabetic kidney disease by promoting ketone body-induced mTORC1 inhibition. *Cell Metab* (2020) 32(3):404–19.e6. doi: 10.1016/j.cmet.2020.06.020
70. Chen XF, Chen X, Tang X. Short-chain fatty acid, acylation and cardiovascular diseases. *Clin Sci (Lond)* (2020) 134(6):657–76. doi: 10.1042/CS20200128
71. Bae HR, Kim DH, Park MH, Lee B, Kim MJ, Lee EK, et al. β -hydroxybutyrate suppresses inflammasome formation by ameliorating endoplasmic reticulum stress via AMPK activation. *Oncotarget* (2016) 7(41):66444–54. doi: 10.18632/oncotarget.12119
72. Chakraborty S, Galla S, Cheng X, Yeo JY, Mell B, Singh V, et al. Salt-responsive metabolite, β -hydroxybutyrate, attenuates hypertension. *Cell Rep* (2018) 25(3):677–89.e4. doi: 10.1016/j.celrep.2018.09.058
73. Place DE, Kanneganti TD. Fueling ketone metabolism quenches salt-induced hypertension. *Trends Endocrinol Metab* (2019) 30(3):145–7. doi: 10.1016/j.tem.2019.01.004
74. Graff EC, Fang H, Wanders D, Judd RL. Anti-inflammatory effects of the hydroxycarboxylic acid receptor 2. *Metabolism* (2016) 65(2):102–13. doi: 10.1016/j.metabol.2015.10.001
75. Suzuki M, Suzuki M, Kitamura Y, Mori S, Sato K, Dohi S, et al. Beta-hydroxybutyrate, a cerebral function improving agent, protects rat brain against ischemic damage caused by permanent and transient focal cerebral ischemia. *Jpn J Pharmacol* (2002) 89(1):36–43. doi: 10.1254/jjp.89.36
76. Taggart AK, Kero J, Gan X, Cai TQ, Cheng K, Ippolito M, et al. (D)-beta-hydroxybutyrate inhibits adipocyte lipolysis via the nicotinic acid receptor PUMA-G. *J Biol Chem* (2005) 280(29):26649–52. doi: 10.1074/jbc.C500213200
77. Tieu K, Perier C, Caspersen C, Teismann P, Wu DC, Yan SD, et al. D-beta-hydroxybutyrate rescues mitochondrial respiration and mitigates features of Parkinson disease. *J Clin Invest*. (2003) 112(6):892–901. doi: 10.1172/JCI200318797
78. Lutas A, Yellen G. The ketogenic diet: metabolic influences on brain excitability and epilepsy. *Trends Neurosci* (2013) 36(1):32–40. doi: 10.1016/j.tins.2012.11.005
79. Yin JX, Maalouf M, Han P, Zhao M, Gao M, Dharshaun T, et al. Ketones block amyloid entry and improve cognition in an alzheimer's model. *Neurobiol Aging* (2016) 39:25–37. doi: 10.1016/j.neurobiolaging.2015.11.018
80. Kashiwaya Y, Takeshima T, Mori N, Nakashima K, Clarke K, Veech RL. D-beta-hydroxybutyrate protects neurons in models of alzheimer's and parkinson's disease. *Proc Natl Acad Sci U S A* (2000) 97(10):5440–4. doi: 10.1073/pnas.97.10.5440
81. Hu E, Du H, Zhu X, Wang L, Shang S, Wu X, et al. Beta-hydroxybutyrate promotes the expression of BDNF in hippocampal neurons under adequate glucose supply. *Neuroscience* (2018) 386:315–25. doi: 10.1016/j.neuroscience.2018.06.036
82. Gersner R, Gal R, Levit O, Moshe H, Zangen A. Inherited behaviors, BDNF expression and response to treatment in a novel multifactorial rat model for depression. *Int J Neuropsychopharmacol* (2014) 17(6):945–55. doi: 10.1017/S1461145714000030
83. Hu E, Du H, Shang S, Zhang Y, Lu X. Beta-hydroxybutyrate enhances BDNF expression by increasing H3K4me3 and decreasing H2AK119ub in hippocampal neurons. *Front Neurosci* (2020) 14:591177. doi: 10.3389/fnins.2020.591177
84. Obokata M, Negishi K, Sunaga H, Ishida H, Ito K, Ogawa T, et al. Association between circulating ketone bodies and worse outcomes in hemodialysis patients. *J Am Heart Assoc* (2017) 6(10):e006885. doi: 10.1161/JAHA.117.006885
85. Poplawski MM, Mastaitis JW, Isoda F, Grosjean F, Zheng F, Mobbs CV. Reversal of diabetic nephropathy by a ketogenic diet. *PLoS One* (2011) 6(4):e18604. doi: 10.1371/journal.pone.0018604
86. Torres JA, Kruger SL, Broderick C, Amaralbhagya T, Agrawal S, Dodam JR, et al. Ketosis ameliorates renal cyst growth in polycystic kidney disease. *Cell Metab* (2019) 30(6):1007–23.e5. doi: 10.1016/j.cmet.2019.09.012
87. Renehan AG, Tyson M, Egger M, Heller RF, Zwahlen M. Body-mass index and incidence of cancer: a systematic review and meta-analysis of prospective

observational studies. *Lancet* (2008) 371(9612):569–78. doi: 10.1016/S0140-6736(08)60269-X

88. Freedland SJ, Mavropoulos J, Wang A, Darshan M, Demark-Wahnefried W, Aronson WJ, et al. Carbohydrate restriction, prostate cancer growth, and the insulin-like growth factor axis. *Prostate* (2008) 68(1):11–9. doi: 10.1002/pros.20683

89. Mavropoulos JC, Buschemeyer WC3rd, Tewari AK, Rokhfeld D, Pollak M, Zhao Y, et al. The effects of varying dietary carbohydrate and fat content on survival in a murine LNCaP prostate cancer xenograft model. *Cancer Prev Res (Phila)* (2009) 2(6):557–65. doi: 10.1158/1940-6207.CAPR-08-0188

90. Park SJ, Kong HK, Kim YS, Lee YS, Park JH. Inhibition of s-adenosylhomocysteine hydrolase decreases cell mobility and cell proliferation through cell cycle arrest. *Am J Cancer Res* (2015) 5(7):2127–38.

91. Belužić L, Grbeša I, Belužić R, Park JH, Kong HK, Kopjar N, et al. Knock-down of AHCY and depletion of adenosine induces DNA damage and cell cycle arrest. *Sci Rep* (2018) 8(1):14012. doi: 10.1038/s41598-018-32356-8

92. Dorman D, Wertz I, Shimizu H, Arnott D, Frantz GD, Dowd P, et al. The ubiquitin ligase COP1 is a critical negative regulator of p53. *Nature* (2004) 429(6987):86–92. doi: 10.1038/nature02514

93. Donehower LA, Harvey M, Slagle BL, McArthur MJ, Montgomery CA Jr., Butel JS, et al. Mice deficient for p53 are developmentally normal but susceptible to spontaneous tumours. *Nature* (1992) 356(6366):215–21. doi: 10.1038/356215a0

94. Wang SJ, Li D, Ou Y, Jiang L, Chen Y, Zhao Y, et al. Acetylation is crucial for p53-mediated ferroptosis and tumor suppression. *Cell Rep* (2016) 17(2):366–73. doi: 10.1016/j.celrep.2016.09.022

95. Ferrannini E, Mark M, Mayoux E. CV Protection in the EMPA-REG OUTCOME trial: A "Thrifty substrate" hypothesis. *Diabetes Care* (2016) 39(7):1108–14. doi: 10.2337/dc16-0330

96. Ohashi K, Ouchi N, Matsuzawa Y. Anti-inflammatory and anti-atherogenic properties of adiponectin. *Biochimie* (2012) 94(10):2137–42. doi: 10.1016/j.biochi.2012.06.008

97. de Boer IH. A new chapter for diabetic kidney disease. *N Engl J Med* (2017) 377(9):885–7. doi: 10.1056/NEJMe1708949

98. Strain WD, Paldanius PM. Diabetes, cardiovascular disease and the microcirculation. *Cardiovasc Diabetol* (2018) 17(1):57. doi: 10.1186/s12933-018-0703-2

99. Li Z, Zhang Y, Han M, Deng H, Wu F, Liu G, et al. Lysine β -hydroxybutyrylation improves stability of COVID-19 antibody. *Biomacromolecules* (2021). doi: 10.1021/acs.biomac.1c01435