

Review

# Ketogenic Diet and Ketone Bodies as Clinical Support for the Treatment of SARS-CoV-2—Review of the Evidence

Izabela Bolesławska <sup>1,\*</sup>, Magdalena Kowalówka <sup>1</sup>, Natasza Bolesławska-Król <sup>2</sup> and Juliusz Przysławski <sup>1</sup>

<sup>1</sup> Department of Bromatology, Poznan University of Medical Sciences, 60-806 Poznan, Poland; mkowalowka@ump.edu.pl (M.K.); jprzysla@ump.edu.pl (J.P.)

<sup>2</sup> Student Society of Radiotherapy, Collegium Medicum, University of Zielona Gora, Zyta 28, 65-046 Zielona Góra, Poland; natmeff@gmail.com

\* Correspondence: ibolesla@ump.edu.pl

**Abstract:** One of the proposed nutritional therapies to support drug therapy in COVID-19 is the use of a ketogenic diet (KD) or ketone bodies. In this review, we summarized the evidence from tissue, animal, and human models and looked at the mechanisms of action of KD/ketone bodies against COVID-19. KD/ketone bodies were shown to be effective at the stage of virus entry into the host cell. The use of  $\beta$ -hydroxybutyrate (BHB), by preventing the metabolic reprogramming associated with COVID-19 infection and improving mitochondrial function, reduced glycolysis in CD4+ lymphocytes and improved respiratory chain function, and could provide an alternative carbon source for oxidative phosphorylation (OXPHOS). Through multiple mechanisms, the use of KD/ketone bodies supported the host immune response. In animal models, KD resulted in protection against weight loss and hypoxemia, faster recovery, reduced lung injury, and resulted in better survival of young mice. In humans, KD increased survival, reduced the need for hospitalization for COVID-19, and showed a protective role against metabolic abnormalities after COVID-19. It appears that the use of KD and ketone bodies may be considered as a clinical nutritional intervention to assist in the treatment of COVID-19, despite the fact that numerous studies indicate that SARS-CoV-2 infection alone may induce ketoacidosis. However, the use of such an intervention requires strong scientific validation.

**Keywords:** COVID-19; ketogenic diet; ketone bodies; SARS-CoV-2



**Citation:** Bolesławska, I.; Kowalówka, M.; Bolesławska-Król, N.; Przysławski, J. Ketogenic Diet and Ketone Bodies as Clinical Support for the Treatment of SARS-CoV-2—Review of the Evidence. *Viruses* **2023**, *15*, 1262. <https://doi.org/10.3390/v15061262>

Academic Editor: Yinzhong Shen

Received: 20 April 2023

Revised: 23 May 2023

Accepted: 25 May 2023

Published: 27 May 2023



**Copyright:** © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Viral infections are a constant threat to public health. Morbidity and mortality associated with seasonal influenza A epidemics but also regularly emerging new viruses with pandemic potential such as Severe Acute Respiratory Syndrome (SARS), N5N1 influenza, H1N1 influenza, Middle East Respiratory Syndrome (MERS) and, most recently, Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) [1] is a key global healthcare problem, requiring the identification of new therapies capable of reducing the severity of these infections.

The SARS-CoV-2 pandemic, which has caused a wave of illness and deaths worldwide as of 2019 [2,3], has not yet been fully contained [4,5]. Although concentrated efforts by researchers and clinicians have led to the identification of new therapeutic options [4,6] and vaccines [5], full immunization with vaccines does not appear to be sufficiently effective against new variants of SARS-CoV-2 [7,8], and treatment is still mainly based on symptomatic treatment [9,10]. This has prompted the search for innovative solutions that could prevent virus entry, strengthen the body's protective barrier, support COVID-19 treatment and recovery from the disease, and protect against new SARS-CoV-2 strains.

Numerous studies conducted in this area have pointed to the prophylactic role of adequate nutrition in reducing the risk of SARS-CoV-2 infection [11–13]. More recently, the

role of clinical nutritional therapy as support for drug therapy in improving the course of COVID-19 has also been highlighted [14].

One of the dietary options being considered is the ketogenic diet (KD), which aims to induce physiological ketosis [1] by significantly restricting the offset of carbohydrates by high-fat content. Ketone bodies resulting from KD, such as  $\beta$ -hydroxybutyrate (BHB), acetoacetate (AcAc), and acetone, affect the body by producing, for example, anticonvulsant effects [15,16] used in the treatment of drug-resistant epilepsy since the early 20th century [17–19]. There are also an increasing number of studies confirming the neuroprotective effects of KD/ketone bodies [20–22], their efficacy in the prevention of Parkinson's and Alzheimer's disease [23,24], the treatment of obesity [25–28], type 2 diabetes [26,28], cardiovascular disease [29–31], or cancer [28,32,33]. Several studies have also confirmed the immunostimulatory and anti-inflammatory effects [14,34–39] of KD and ketone bodies. Induction of a state of ketosis by KD or ketone bodies and their pleiotropic effects may also prove to be an effective intervention in the prevention and/or treatment of COVID-19 coronavirus disease [40–45].

KD and beta-hydroxybutyrate (BHB), accounting for approximately 70% of the circulating pool of ketone bodies [46], exhibit a variety of molecular effects including metabolic regulation and effects on modulation of the epigenome, resulting in changes in DNA methylation, chromatin histone modifications and miRNA levels [46,47], and enhancement of cellular resistance to oxidative stress through multiple mechanisms [46,48] including regulation of the miRNA expression profile [49]. However, the molecular mechanisms underlying the effects of these diets are only partially understood and can result in both physiologically beneficial outcomes and potentially harmful effects [46,47]. Of particular concern is the potential for diabetic ketoacidosis leading to coma and even death (with ketone body concentrations exceeding 20 mM) [46,50]. There is a bidirectional relationship between SARS-CoV-2 and diabetes—SARS-CoV-2 can directly damage pancreatic beta cells, exacerbating hyperglycemia, causing or exacerbating diabetes in a significant number of patients, while the presence of diabetes can exaggerate a COVID-19 infection [51–53]. In such patients, the use of KD could exacerbate diabetes and worsen COVID-19.


Given the specificity of SARS-CoV-2, hard evidence is needed to support the actual efficacy of KD and/or ketone bodies against the virus. A review of the evidence and a look at the proposed mechanisms of action of KD/ketone bodies against SARS-CoV-2 virus was the aim of this manuscript. Notably, since the SARS-CoV-2 virus shares similarities with other viruses, including those from distant viral families [54], it causes co-infections with other viral diseases leading to worsening clinical outcomes of COVID-19 [55]. There are also concerns that the SARS-CoV-2 virus, despite pandemic containment by vaccination, could become a persistent threat to global health like the Middle East Respiratory Syndrome (MERS) coronavirus before it [56] or the Zika virus [57]. Whether in the form of an entirely new variant or evolving from the Omicron sublineage with new antigenic changes, the new variants will exhibit variability in clinically relevant features, including sensitivity to drug therapies [58]. A dietary approach using KD or ketone bodies that can change the metabolic environment to one that is unfavorable to the entry and existence of the virus in the human body may prove effective in both prevention and treatment support.

## 2. Course of SARS-CoV-2 Infection

SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2) is a single-stranded enveloped virus with positive ssRNA polarity [59–62]. Compared to previously known coronaviruses, SARS-CoV-2 has high virulence and a higher transmission rate [62].

After contact with the cell, SARS-CoV-2 virus can enter the cell either through endosomes or through plasma membrane fusion. In both cases, spike proteins (S1/S2) present on the surface of SARS-CoV-2 mediate attachment to the cell membrane by binding to angiotensin-converting enzyme type 2 (ACE2) as an entry receptor [63,64]. This is made possible by the characteristic structure of the structural spike protein (S) that forms the outer envelope of the virus, consisting of three subunits: S1, S2, and S2'. The S1 sub-

unit via the Receptor Binding Domain (RBD) of the virus binds to the ACE2 receptor on the cell surface [65–67] and then host cell surface proteases such as TMPRSS2 (transmembrane serine protease 2), cathepsin, furin, or yet other proteins (e.g., CD147, GRP78, ADAM17) act on critical cutting sites at the S1/S2 interface of the spike protein and at the S2' site [61,65,68–71]. This results in membrane fusion and viral infection [59]. The interaction between the SARS-CoV-2 coronavirus spike protein and the host cell ACE2 receptor is shown in Figure 1.



**Figure 1.** Interaction between the SARS-CoV-2 coronavirus spike protein and the host cell receptor ACE2 [72].

Once the virus has entered the host cell and released its genetic material, the viral proteins necessary for further replication and translation are synthesized. The formed new viral particles, capable of entering further host cells, are translocated close to the cell membrane and released exocytosis [63,73].

The course of the disease itself and its clinical picture depends primarily on the nature and type of immune response to infection. This is made up of many factors including individual genetic predisposition, specific immune response, and the inflammation present [74].

SARS-CoV-2 virus infection stimulates several immune response pathways in the host, including activation of both innate and acquired immune responses [75]. The defender against SARS-CoV-2 is the innate immune system eliminating SARS-CoV-2 in the best-case scenario without activation of the adaptive immune system [74,76]. However, an effective immune response against SARS-CoV-2 often requires activation of both arms of the immune system—the innate immune system including granulocytes, monocytes, macrophages, NK cells, and other cells of the innate immune system (including various dendritic cells, innate lymphoid cells, or mast cells) and the acquired (adaptive) immune system with T and B lymphocytes [77]. During SARS-CoV-2 infection, an inappropriately activated and dysregulated host immune response may cause immunopathology [78,79]. The course of infection may also be influenced by the external environment and the virus itself due to its genetic diversity, genetic evolution, and variable infectivity [77].

In the vast majority of patients, COVID-19 is asymptomatic or mild (80% of cases), but in some cases, there is a rapid and unpredictable course of the disease [77]. In about 5% of infected patients, the immune response runs uncontrollably, leading to inflammatory cascades, cytokine storms, and activation of coagulation cascades that further cause damage to other organs [60,80] and, in extreme cases, sepsis and death [81]. However, the detailed mechanisms of pathogenesis are not yet understood. A schematic of the COVID-19 run is shown in Figure 2.




Figure 2. COVID-19 flow diagram [63].

### 3. Characteristics and Performance of KD

The ketogenic diet is a dietary regimen designed to increase the synthesis and utilization of ketone bodies [82,83]. Its characteristics are high fat and low carbohydrate. This macronutrient ratio allows for the production of nutritional ketosis—glucose-dependent tissues (heart and skeletal muscle) can partially adapt to fat metabolism and utilize ketone bodies such as acetone, BHB, or AcAc [84,85] as the leading energy substrate [84,86,87]. Ketosis is usually achieved by restricting carbohydrate intake below 50 g/day (5–10%) [82,83], providing 1–1.5 g/kg/day of protein, and 60 to 90% of energy coming from fat [23,88–93] (Table 1).

The concentration of ketone bodies in the circulation under normal physiological conditions usually fluctuates around 100  $\mu$ M–1 mM [94] but can rise to 5–8 mM after a low-carbohydrate diet [86,95,96] and >20 mM in pathological conditions such as diabetic or alcoholic ketoacidosis [97].

**Table 1.** Composition of selected ketogenic diets to achieve nutritional ketosis.

Number	Type of KD	Carbohydrates /Day	Fat/Day	Protein/Day	Ketosis
1.	Classical Ketogenic Diet (CKD)—children [88–90]	10–15 g/day	90% EV	1 g/kg bw/day	✓
2.	Medium-Chain Triglyceride (MCT) [91]	50 g/day 20% EV	78 g/day 70% EV	25 g/day 10% EV	✓
3.	Low-glycemic index treatment (LGIT) [98]	40 g/day 27% EV	60 g/day 45% EV	40 g/day 28% EV	✓
4.	Medium-Chain Triglyceride-Based Diet [92]				✓
5.	Modified Atkins Diet (MAD) [93]	10 g/day 5% EV	70 g/day 70% EV	60 g/day 25% EV	✓
6.	Mediterranean-ketogenic diet (MMKD) [23]	<20 g/day <10% EV	60–65% EV	30–35% EV	✓
7.	Eko Atkins diet [99]	26% EV	43% EV	31% EV	✓
8.	Ketonformula <sup>®</sup> , 50 g + usual diet [100]				✓


% EV—% of energy; bw—body weight; ✓—yes.

Reduced by KD use, insulin levels together with high adrenaline levels lead to the release of free fatty acids and glycerol from adipocytes under the action of lipase. Free fatty acids undergo  $\beta$ -oxidation in the liver mitochondria with the production of acetyl-CoA, which under standard conditions is incorporated into the Krebs cycle by condensation of acetyl-CoA with oxaloacetate [94,97]. Low dietary carbohydrate supply increases the rate of fatty acid oxidation in the liver and consumes the hepatic pool of oxaloacetate for gluconeogenesis. Accumulated acetyl-CoA exceeds the ability to synthesize citrate thus becoming a precursor of ketone bodies [82,101,102].

The breakdown of fatty acids in the cytosol of prokaryotic cells and in the mitochondrial matrix of eukaryotic cells is preceded by the conversion of fatty acids in an ATP-mediated reaction catalyzed by acyl-CoA synthetase (so-called fatty acid thiokinase) into an active metabolite [103]. Once the acyl chains formed from fatty acids have passed through the mitochondrial membranes, they are introduced into the mitochondria via carnitine palmitoyltransferase (CPT-1) [104] and then broken down into acetyl-CoA via  $\beta$ -oxidation. Two acetyl-CoA molecules combine to form acetoacetyl-CoA (AcAc-CoA) in a reaction catalyzed by the enzyme thiolase, and then acetoacetyl-CoA reacts with the next acetyl-CoA molecule and with water to form 3-hydroxy-3-methylglutaryl-CoA (HMG-CoA) via the enzyme HMG-CoA synthase. HMG-CoA in the presence of a lyase is cleaved into acetyl-CoA and acetoacetate. In the mitochondrial matrix, acetoacetate is converted to acetone by non-enzymatic decarboxylation, or is reduced to  $\beta$ -hydroxybutyrate ( $\beta$ -OHB) via 3-hydroxybutyrate dehydrogenase [29,85,102].

The ketone bodies produced are released from the liver into the circulatory system. Acetone, as a volatile substance, does not convert back to acetyl-CoA and is removed from the body via the lungs and kidneys. After reaching extrahepatic tissues,  $\beta$ -hydroxybutyrate

(BDH1) is converted to acetoacetate via the enzyme beta-hydroxybutyrate dehydrogenase, and acetoacetate is converted back to acetyl-CoA via the enzyme beta-ketoacyl-CoA transferase, goes through the citric acid cycle, and produces 22 ATP per molecule after oxidative phosphorylation (Figure 3) [85,102].



**Figure 3.** Scheme of metabolism of ketone bodies. BDH1—D- $\beta$ -OHB dehydrogenase, SCOT—succinyl-CoA:3-oxoacid-CoA transferase, TCA—the tricarboxylic acid cycle. (A) The ketone bodies acetoacetate (AcAc),  $\beta$ -hydroxybutyrate ( $\beta$ -OHB), and acetone are majorly produced in the mitochondrial matrix of liver cells during the oxidation of fatty acids. Ketone synthesis begins in the mitochondria of the liver after transporting the fatty acyl CoA molecule into the inner mitochondrial membrane by the carnitine shuttle. The transmembrane protein to move fatty acyl CoA molecules across the mitochondrial membrane is carnitine palmitoyl transferase I (CPT-1). The fatty acyl CoA molecules undergo  $\beta$ -oxidation to become acetyl CoA molecules. Acetyl CoA molecules are either converted to malonyl CoA by acetyl CoA carboxylase or acetoacetyl CoA by 3-ketothiolase. Acetoacetyl CoA is further converted to 3-hydroxy-3-methylglutaryl CoA (HMG CoA) by HMG CoA synthase. HMG CoA synthase is the rate-limiting step for the synthesis of ketone bodies. HMG CoA is finally converted to acetoacetate by HMG CoA lyase. (B) Upon arrival to the mitochondria of distant organs, ketone bodies become utilized for energy. The first step involved is an enzyme that converts acetoacetate to acetoacetyl CoA. This enzyme is called succinyl CoA-oxoacid transferase (SCOT), and it is the rate-limiting step for the utilization of ketones for energy. High concentrations of acetoacetate feedback negatively on SCOT to decrease ketone conversion. Finally, acetoacetyl CoA is converted to acetyl CoA by methylacetoacetyl CoA thiolase. Acetyl CoA may be turned into citrate and churned through the citric acid cycle to produce FADH<sub>2</sub> and NADH, or it can be converted to oxaloacetate and used in gluconeogenesis. Abbreviations: BDH1—D- $\beta$ -OHB dehydrogenase, SCOT—succinyl-CoA:3-oxoacid-CoA transferase, TCA—the tricarboxylic acid cycle.

KD has been used to treat drug-resistant epilepsy for more than a century [17–19]. There is also a growing body of research supporting the efficacy of KD in neurodegenerative diseases [20–22], including the prevention of Parkinson’s and Alzheimer’s disease [23,24]. A body of evidence supports the beneficial effects of KD in the treatment of obesity [25–28], type 2 diabetes [26,28], cardiovascular disease [29–31], and cancer [28,32,33]. Several studies have also confirmed the immunostimulatory and anti-inflammatory effects of KD and ketone bodies [14,34–39] and their effects on the microbiome [105–107]. The experimental use of KD in mechanically ventilated patients in intensive care units has also yielded good results [108–110].

Recently, there has been a particular increase in research on KD related to the suggested role of KD and ketone bodies in preventing SARS-CoV-2 infection at both the viral entry and dissemination stages in the body. Studies in animals, human biological material, and humans have confirmed these suggestions on several key points. Thus, it appears that the use of KD and ketone bodies may be considered as an adjunctive intervention for the treatment of COVID-19, despite the fact that numerous studies indicate that COVID-19 infection alone can induce ketoacidosis, particularly in patients with diabetes [111–114], but also can cause a high increase in ketone bodies in patients without diabetes [111,115].

On the other hand, Karagannis et al. [116], in a study conducted in the peripheral blood of patients with moderate to severe COVID-19, found low BHB concentrations irrespective of calorie intake and blood glucose levels and thus concluded that ketogenesis is impaired in COVID-19 and may be mediated by a reduction in beta-hydroxybutyrate. However, in the light of the previously described reports, these conclusions need to be confirmed.

#### 4. Efficacy of KD in COVID-19

##### 4.1. Efficacy of KD/Ketone Bodies at the Stage of Virus Entry into the Cell

KD reduces ACE2 (angiotensin-converting enzyme 2) methylation and increases ACE2 mRNA expression in visceral adipose tissue and leukocytes, and decreases ACE2 and TMPRSS2 (transmembrane serine protease 2) levels in the lung.

The ACE2 used by SARS-CoV-2 to enter the cell is mainly found in the lung [117] but also in other locations including the small intestine, oesophagus, liver, colon [118], and adipose tissue [119,120].

The clinical course of infection depends largely on which host cells are infected first. Higher expression of the ACE2 gene in visceral and subcutaneous adipose tissue than in lung tissue may exacerbate infection, prolong hospitalization time, and increase the risk of death in obese patients with COVID-19 [121–124]. In a study by Izquierdo et al. [125], visceral adipose tissue and leucocyte samples from obese patients showed higher levels of ACE2 methylation and lower ACE2 gene expression than in normal-weight patients. The use of low-calorie KD reversed this unfavorable obesity-associated ACE2 methylation which resulted in an increase in ACE2 mRNA expression and reduced dangerous disease progression (Table 2).

Additionally, in a study in COVID-19-infected rats, the application of KD reduced the levels of ACE2 and TMPRSS2 (required to stimulate the S protein of the virus to allow it to enter the cell) in the lungs resulting in reduced susceptibility to COVID-19 infection [126] (Table 3). Thus, KD may act similarly to the cellular serine protease inhibitor TMPRSS2 approved for clinical use [127] which blocks SARS-CoV-2 entry and may be one option for treating the disease.

KD causes a shift of the renin–angiotensin–aldosterone system (RAS) from the AngI/ACE1/AngII/AT1 arm towards the counter-regulatory, anti-inflammatory arm of the RAS.

The renin–angiotensin–aldosterone system (RAS) plays a significant role in the regulation of the body’s water and electrolyte balance, as well as renal, neuronal, and endocrine functions related to cardiovascular control [128]. Due to the extensive and multi-organ action of the RAS, its overactivation contributes to the development of many diseases. The RAS can proceed through two pathways [129].



**Table 2.** Studies on the efficacy of KD/ketone bodies in SARS-CoV-2 on biological material from patients with COVID-19.

	Study Material/Quantity of Samples	Evaluation	Intervention/Measurement	Results	Bibliography
1	Serum from patients with moderate COVID-19 or ARDS due to COVID-19/n = 155	Evaluation of the relationship between infection-induced metabolic changes and host immune responses in severe lung infections	Cultured human TCD4+ / E ketone ester BHB (d - $\beta$ -hydroxybutyrate-(R)-1,3-butanediol monoester) added at days 0, 1, and 2 in an amount of 5 mM BHB (6 days of culture)	<ul style="list-style-type: none"> <li>- in the peripheral blood of patients with moderate to severe COVID-19, low concentrations of BHB irrespective of calorie intake and blood glucose levels</li> <li>- BHB resulted in improved mitochondrial function, and fatty acid and amino acid oxidation; reduced glycolysis in CD4 + T lymphocytes</li> <li>- cultures of human TH1 CD4 + lymphocytes with BHB showed an increase in the number of CD4+ cells and IFN<math>\gamma</math> production, and a decrease in PD-1 (programmed death receptor 1) expression</li> </ul>	Karagiannis et al. [116]
2	Immune cells from patients with COVID-19; disease severity >IV <sup>0</sup> and respiratory failure/n = 9/20	Evaluation of the efficacy of metabolic reprogramming of CD8 + T cells by ketone bodies in overcoming immune paralysis in patients with COVID-19	Peripheral blood mononuclear cells (PBMCs) from COVID-19 patients. Incubation with BHB, D/L-beta-hydroxybutyrate with a final concentration of 10 mM	<ul style="list-style-type: none"> <li>- significant increase in granzyme B-expressing CD8 + lymphocytes and increased granzyme B expression per cell,</li> <li>- profound enhancement of CD8 + immune capacity</li> <li>- significantly increased secretion of the T lymphocyte cytokines CD8 + IFN<math>\gamma</math>, TNF<math>\alpha</math>, perforin, and granzyme B</li> <li>- markedly increased cell lysis capacity of CD8 + T lymphocytes</li> <li>- CD8 + T lymphocytes showed significantly higher basal and maximal respiratory chain activity and better spare respiratory capacity</li> <li>- a tendency towards increased mitochondrial mass in CD8 + T cells</li> <li>- significantly increased mitochondrial ROS but cellular ROS levels remained unchanged</li> </ul>	Hirschberger et al. [143]

Table 2. Cont.

	Study Material/Quantity of Samples	Evaluation	Intervention/Measurement	Results	Bibliography
3	DNA methylation levels of ACE2 from data sets generated by hybridization of subcutaneous (n = 32), visceral (n = 32) adipose tissue, and leukocyte samples (n = 34).	Evaluation of ACE2 methylation levels in different depots of adipose tissue and leukocytes in obesity	DNA methylation levels of ACE2. Data were compared based on degree of obesity and after 4–6 months of weight loss following weight loss therapy based on VLCKD vs. HCD vs. bariatric surgery	- decreased levels of ACE2 methylation and increased ACE2 mRNA expression in visceral adipose tissue and leukocytes in obese patients after VLCKD	Izquierdo et al. [125]
4	Simulated Kbh antibody produced using in vitro chemical modification from a commercially obtained human serum antibody	Investigating the immunomodulatory mechanisms of $\beta$ -hydroxybutyrate	Antibody solution with addition of free $\beta$ -hydroxybutyrate	Kbh did not affect the binding capacity of B38 to the S-protein of COVID-19 virus, indicating that the chemically modified antibody retained its original antigen-binding activity	Li et al. [144]

VLCKD—very low-calorie ketogenic diet, HCD—balanced hypocaloric diet, BHB—beta-hydroxybutyrate, ARDS—acute respiratory distress syndrome, ACE2—angiotensin-converting enzyme type 2, IFN $\gamma$ —interferon gamma, TNF $\alpha$ —tumor necrosis factor  $\alpha$ , ROS—reactive oxygen species, Kbh -.

**Table 3.** Studies on the efficacy of KD in SARS-CoV-2 in animal models.

Study Design	Participants (n)/Age (year)	Evaluation	Intervention/Measurement	Duration	Results	Bibliography
1 Randomized clinical trial	Male C57BL mice infected with natural mouse beta coronavirus (mCoV) 6—mouse hepatitis virus strain A59 (MHV-A59) (reproducing the clinical features of COVID-19). Old mice (20–24 months) and young mice (2–6 months)	Effect of KD on the mouse defense response against MHV-A59 and identification of the underlying mechanism	Intranasal inoculation of mCoV-A59 into adult and old male mice. Feeding a standard vivarium feed (Harlan 2018s) or a ketogenic diet (Envigo, TD.190049)	5 days before infection	<ul style="list-style-type: none"> <li>- protection against weight loss and hypoxemia caused by infection</li> <li>- improved survival of young but not old mice</li> <li>- significantly reduced mRNA expression of the pro-inflammatory cytokines IL-1<math>\beta</math>, TNF<math>\alpha</math>, and IL-6 in the lung, VAT, and hypothalamus</li> <li>- in old mice, a significant increase in cup cells, expansion of T <math>\gamma\delta</math>, and a significant decrease in subsets of proliferative cells</li> <li>- significant decrease in monocyte population subsets, change in monocyte compartment, and loss of cluster with high levels of Chil3, Lmna, Il1r2, Lcn2, Cd33, and Cd24a; loss of monocyte subpopulation in cells with low interferon response</li> <li>- increase in T <math>\gamma\delta</math> cells, downregulation of TLR, Plk1, and aurora B signaling pathways in T <math>\gamma\delta</math> cells</li> <li>- in old mice, increased genes associated with reduced inflammation, increased lipoprotein remodeling</li> <li>- increased respiratory electron transport and complex I biogenesis in lung T <math>\gamma\delta</math> cells of old mice</li> <li>- reduced activation status of T <math>\gamma\delta</math> lymphocytes</li> <li>- decreased NLRP3 and caspase-1 mRNA in lung, visceral adipose tissue (VAT), and hypothalamus, and inflammasome activation in VAT</li> <li>- significantly reduced myeloid cell infiltration in the heart</li> <li>- BHB reduced oligomerization of ASC, which is an adaptor protein required for assembly of the inflammasome complex</li> </ul>	Ryu et al. [42]
2 Randomized clinical trial	Male albino rats of the Sprague strain/body weight 200–250 g	Measurements of ACE2, TMPRSS2, and RAS components and inflammatory genes in animal lungs and hearts	Chow diet (P: 27.0%, F: 13.0%, and C: 60.0%), high-fat sucrose-enriched diet (P: 20.0%, F: 60.0%, and C:20.0%) or KD (P:20%, F: 80%, and C: 0%).	16 weeks	<ul style="list-style-type: none"> <li>- reduced lung ACE2 and TMPRSS2 levels compared to a sucrose-enriched diet</li> <li>- KD in the lung resulted in decreased ACE1 and AT1 protein content, and a shift of the (RAS) system from the AngI/ACE1/AngII/AT1 arm towards the counter-regulatory, anti-inflammatory arm of the RAS</li> <li>- KD induced anti-inflammatory effects—reduction of tlr4 and il6r gene expression, and a tendency to reduce tnfr1 gene expression</li> </ul>	Eira et al. [126]

Table 3. Cont.


Study Design	Participants (n)/Age (year)	Evaluation	Intervention/Measurement	Duration	Results	Bibliography
3 Randomized clinical trial	K18-hACE2 mice aged 6–20 weeks infected intranasally with 60 PFU of SARS-CoV-2 (preclinical model of SARS-CoV-2 infection)	Assessment of T-cell metabolism and function	Supplementation with ketone ester 20 mg ml <sup>-1</sup> (D- - hydroxybutyrate-(R)-1,3 butanediol monoester) in drinking water	8 days after infection	- decreased glucose dependence of CD4+ lymphocytes and increased the potential to oxidize amino acids and fatty acids thereby increasing their ability to produce IFN $\gamma$ and antiviral protection - ketone ester promoted faster recovery from weight loss and reduced lung injury, resulting in improved overall survival	Karagiannis et al. [116]

KD—ketogenic diet, P—protein, F—fat, C—carbohydrates, VAT—visceral adipose tissue, ER—endoplasmic reticulum, NLRP3—NLRP3 inflammasome, BHB—beta-hydroxybutyrate, TMPRSS2—transmembrane serine protease 2, RAS—renin-angiotensin-aldosterone system, ACE1—angiotensin-converting enzyme I protein, AT1—angiotensin receptor type 1 protein.

#### 4.2. Effectiveness of KD/Ketone Bodies in Regulating Innate and Acquired Immunity

The atypical course of COVID-19 in about 20% of patients may be due to a unique dysregulation of the immune response [79]. In contrast, ketone bodies may regulate immune cells, although their effect on the immune response is not fully understood [34,36].

Previous studies suggest that BHBs, by preventing metabolic reprogramming associated with COVID-19 infection and improving mitochondrial function, resulted in reduced glycolysis in CD4+ lymphocytes, improved respiratory chain function, and could provide an alternative carbon source to OXPHOS. In studies with animal models and human biological material, it was found that the use of KD/ketone bodies could support the host immune response as shown in Figure 5.



**Figure 5.** The role of the ketogenic diet and ketone bodies in immunity. The KD expands metabolically protective  $\gamma\delta$  T cells that restrain inflammation.  $\beta$ -hydroxybutyrate inhibits NLRP3 inflammasome activation.  $\gamma\delta$  T cells in visceral adipose tissue are activated by KD and might have important roles in adipose tissue remodeling and homeostasis. HIF-1 $\alpha$  also induced expression of the proinflammatory cytokines TNF- $\alpha$  and IL-6 and decreased their mRNA expression. BHB improved respiratory chain function and prevented CD4+ and CD8+ T-lymphocyte dysfunction.

##### 4.2.1. BHB Reduces Cellular Dependence on Glucose

In healthy organisms, when immune cell activation is not required, glucose is metabolized by the glycolytic pathway of primary importance in sustaining life in almost all organisms, yielding pyruvate and NADH (nicotinamide adenine dinucleotide) as products [145]. These intermediates are then utilized in the mitochondria, in the tricarboxylic acid (TCA) cycle of cellular respiration, and oxidative phosphorylation (OXPHOS), forming intermediate compounds that are used in the electron transport chain to produce 32 molecules of adenosine triphosphate (ATP) for every molecule of glucose [146,147]. During the immune response when immune cells are activated, for example, in COVID-19 infection, effector T cells to meet the increased bioenergetic demands undergo metabolic reprogramming involving a switch from oxidative phosphorylation to glycolysis [148]. Such reprogramming results in compromised ATP production by OXPHOS [145].

A study by Karagannis et al. [116] found that T lymphocytes in bronchoalveolar lavage fluid (BALF) and blood from subjects with SARS-CoV-2-induced acute respiratory distress syndrome (ARDS) showed a significantly altered metabolic profile towards glycolysis, but still exhibited a residual capacity to oxidize fatty acids and amino acids [116]. However, in COVID-19, the available amount of amino acids and BHB is reduced [116,149], and

marked changes in lipid and amino acid metabolism are also observed [150]. Meanwhile, BHB treatment resulted in improved mitochondrial functionality for fatty acid and amino acid oxidation while leading to reduced glycolysis in CD4+ lymphocytes in patients with severe COVID-19 [116]. Similarly, the use of ketone ester in a preclinical mouse model of SARS-CoV-2 infection reduced the dependence of CD4+ lymphocytes on glucose and increased the potential to oxidize amino acids and fatty acids thereby increasing their ability to produce interferon- $\gamma$  (IFN $\gamma$ ), which plays an important role in the body's immune response during infection control [151] and antiviral protection [116].

#### 4.2.2. BHB Promotes Mitochondrial Function, Improves Respiratory Chain Function, and May Provide an Alternative Carbon Source to OXPHOS

SARS-CoV-2 virus RNA localizes to mitochondria and then manipulates mitochondrial function causing mitochondrial dysfunction [152], disrupting platelet mitochondrial respiratory chain function and mitochondrial oxidative phosphorylation (OXPHOS) [153,154], which in turn will disrupt T-cell function [148,155] and may be the cause of defective adaptive immune responses in COVID-19.

Ketone bodies through immunometabolic reprogramming in T cells can be used for energy production via mitochondrial oxidative phosphorylation (OXPHOS) [97,156].

Hirschberger et al. [143] demonstrated the efficacy of using BHB to increase the energy capacity of CD8+ cells during COVID-19. CD8+ lymphocytes under the influence of BHB showed significantly higher basal and maximal respiratory chain activity and better respiratory reserve capacity, indicating enhanced energy production in the mitochondria. BHB caused an increase in mitochondrial mass in CD8+ lymphocytes. Thus, they demonstrated that T lymphocytes were able to utilize BHB via oxidation in the Krebs cycle, which drives OXPHOS with excellent efficiency [157,158] and showed that ketone bodies direct human CD8+ T lymphocytes towards aerobic mitochondrial metabolism during COVID-19, thus enabling improved energy supply [143].

These observations corroborate an earlier study by Karagannis et al. [116], who demonstrated on cultured mouse and human TH1 CD4+ lymphocytes the ability of BHB to enhance CD4+ mitochondrial functionality and the role of BHB in supporting OXPHOS in mouse CD4+ lymphocytes. In the same study, Karagannis et al. [116] showed that providing mice with a ketogenic diet resulted in metabolic reprogramming of pulmonary CD4+ cells towards OXPHOS and a reduction in their glycolytic capacity, and that BHB provides an alternative carbon source to drive OXPHOS and the production of bioenergetic amino acids (glutamate and aspartate) and glutathione in CD4+ T cells, which is important for maintaining redox balance.

#### 4.2.3. BHB Prevents CD4+ T-Cell Dysfunction

Adequate activation for proliferation, clonal expansion, and effector function of cytotoxic CD4+ type 1 helper T cells (H1 T cells) is crucial in clearing viral infections due to their ability to produce cytokines including interferon- $\gamma$  (IFN $\gamma$ ) [159,160] and their modulation of the activity of other antibodies [116,161].

Severe COVID-19 is associated with significantly reduced CD4+ lymphocyte frequency and dysfunction [162–165]. This is caused by direct binding of the SARS-CoV-2 peak glycoprotein to the CD4+ molecule, which facilitates the entry of SARS-CoV-2 into helper T cells, and further causes cellular impairment or death [166].

In a study by Karagiannis et al. [116], when human TH1 CD4+ lymphocytes obtained from patients with severe COVID-19 with BHB were cultured, an increase in CD4+ cells and IFN $\gamma$  production, and a decrease in PD-1 (programmed death receptor 1) expression were observed. Additionally, previous studies conducted on human immune cells indicated that BHB led to a significant transcriptional upregulation of CD4+ cell cytokines, interleukin (IL)2, IL4, IL8, and IL22 which improved the immune capacity of human T cells [156]. However, according to a study by Ryu et al. [42] in an animal model, ketogenesis in infected old mice did not affect the frequency of CD4+ and effector memory CD8+.

#### 4.2.4. KD and Activation of Ketogenesis Reduced the Number of Pathogenic Monocytes in the Lung, Blocked Infiltration of a Pathogenic Subset of Monocytes in the Lung, and Resulted in the Loss of a Subpopulation of Monocytes with Low Interferon Expression

Monocytes, representing 15–30% of circulating lymphocytes [167], play an important role in both innate and adaptive immunity, inflammation, and tissue remodeling [168]. During infection, they escape from the blood vessels into the surrounding tissues and, after being transformed into macrophages along with them, constitute sentinel cells, engulf viruses, and coordinate the immune system [169]. However, an uncontrolled increase in monocyte levels leading to inflammation has been observed in patients with severe COVID-19 [170].

Ryu et al. [42], in a study with mice (Table 2), observed a significant decrease in subsets of the monocyte population in old, infected mice after KD injection. Induction of ketogenesis in mice by KD application also blocked infiltration of a pathogenic subset of monocytes in the lungs, which show high expression of S100A8/9, an immune-activating alarmin [171–173]. Uncontrolled production or sustained expression of alarmins can be dangerous to the host by inducing a cytokine storm or excessive inflammation [174,175], as observed in severe cases of COVID-19 [78,176,177]. In addition, they noted a loss of a subpopulation of monocytes with low interferon expression after KD treatment [42]. This suggests the induction of an immune response following ketogenesis in infected mice, quenching the overexpression of alarmins and thereby preventing the formation of excessive inflammation and/or cytokine storm in COVID-19.

#### 4.2.5. KD Caused a Slight Decrease in Interleukin-6 (IL-6) Levels in Humans and Reduced mRNA Expression of the Pro-Inflammatory Cytokines IL-1 $\beta$ , TNF $\alpha$ , and IL-6, the Inflammatory Genes TLR4 and ILR6, and Suppressed TNFR1 Gene Expression in Mice

Numerous studies have shown that severe and critically ill COVID-19 patients experience a ‘cytokine storm’ associated not only with excessive cytokine production, but with dysregulation of immune cell function and systemic inflammation [178]. In such patients, a slight decrease in interleukin-6 (IL-6) levels was observed following a eucaloric ketogenic diet (EKD) [14] (Table 4).

**Table 4.** Studies on the efficacy of KD in COVID-19 in humans.

Study Design	Participants (n)/Age (year)	Evaluation	Intervention	Duration	Results	Bibliography
1 Retrospective study in hospitalized patients with COVID-19	EKD n = 34 vs. ESD n = 68	30-day mortality, admission to the intensive care unit; effect of ECD on biological and inflammatory parameters, especially IL-6	ESD 1800–2100 kcal, C: 42–50%, P: 16–20%, F: 26–30%; EKD 1800–2100 kcal, C <30 g, 5–6%, P: 17–18%, and polyunsaturated/unsaturated/saturated fat ratio 3:2:1	follow the diet for a mini-mum of 7 days	<ul style="list-style-type: none"> <li>- statistically significantly higher survival rates</li> <li>- lower need for intensive care unit hospitalization</li> <li>- reduction in IL-6 between days 0 and 7</li> </ul>	Succar et al. (2021) [14]
2 Case report—COVID-19-positive woman, hospitalized for bilateral pneumonia, with sarcopenic obesity, hypertension, hyperinsulinemia, hypercholesterolemia, and hypertriglyceridemia	n = 1, age 55	Anthropometric parameters and body composition, muscle mass and strength, time required to rise from a sitting position without the use of hands, blood pressure, insulin and fasting blood glucose levels, creatinine, triglycerides, LDL cholesterol, HDL cholesterol, vitamin D, and HOMA index	VLCKD (800 kcal/day, C: 28 g (14.6%), F: 35 g (38.7%), P: 85 g (46.7%), 2–2.5 L water/day + interval training 2 times a week 30–35 min/day via the Zoom platform	6 weeks after COVID-19	<ul style="list-style-type: none"> <li>- reduction of body weight, BMI, amount of body fat, maintenance of lean mass,</li> <li>- improved muscle strength and physical performance</li> <li>- reduction in blood pressure, insulin, glycemia and HOMA, TG, LDL, and creatinine levels</li> <li>- increase in HDL</li> </ul>	Camajani et al. [179]
3 A real-world, retrospective exploratory analysis in outpatients with type 2 diabetes, obesity, and positive COVID-19	n = 339, 54.9 ± 8.7 age	hemoglobin A1c, blood glucose, ketone bodies, body weight and body mass index, and assessment of the need for hospitalization	KD—C: <50 g, ~5–10%, P: 1.5 g/kg reference body weight, ~15–20%, F: to satiety ~70–75%	before and during treatment	greater weight loss was significantly associated with a lower likelihood of hospitalization	Volk et al. [180]

KD—ketogenic diet, ESD—eucaloric standard diet, EKD—eucaloric ketogenic diet, VLCKD—very low-energy ketogenic diet, P—protein, F—fat, C—carbohydrates, BMI—body mass index, TG—triglycerides, HOMA index—insulin resistance index, hemoglobin A1c—glycated hemoglobin.

In a study by Ryu et al. [42], older mice fed a KD diet after subtotal infection with mCoV-A59 showed significantly reduced mRNA expression of the pro-inflammatory cytokines IL-1 $\beta$ , TNF $\alpha$ , and IL-6 in the lung, visceral adipose tissue (VAT), and hypothalamus. The use of KD also resulted in reduced expression of the inflammatory genes TLR4 and ILR6, and suppressed expression of the TNFR1 gene in the lungs of mice compared to a sucrose-enriched diet [126] (Table 2). Thus, KD may, by affecting signaling pathways, improve outcomes in COVID-19, including by blocking the expression of genes encoding pro-inflammatory cytokines.

Interestingly, some patients with COVID-19 in severe condition expressed low levels of cytokines in the blood resulting from immune cell depletion [79,181]. In functional studies by Hirschberg et al. [143] on cells obtained from COVID-19 patients (Table 3), incubating them with BHB resulted in increased TNF- $\alpha$  expression. This improved the CD8 + immune capacity, which may consequently counteract the severe complications of COVID-19. However, given the results of studies demonstrating the adverse effects of excess TNF- $\alpha$  in COVID-19, these reports should be approached with caution. While this effect of BHB would be beneficial in cases of immune cell depletion, it could already have adverse consequences in other patients.

#### 4.2.6. KD and Activation of Ketogenesis Resulted in Increased Homeostaticity of T $\gamma\delta$ Lymphocytes

During viral infection, T $\gamma\delta$  lymphocytes are also activated, which, although a small proportion of the total peripheral blood T cells, have the ability to respond rapidly to stimuli, making them one of the most important innate and acquired lines of defense against viral infections [182,183]. Increased activation of T $\gamma\delta$  lymphocytes was observed in patients with COVID-19, they were also more differentiated than control samples [184] and quantitatively reduced levels were observed compared to healthy patients [185,186]. Such an abnormal T $\gamma\delta$  profile in COVID-19 favoured a worse disease scenario [182].

Meanwhile, Ryu et al. [42] showed an increase in T $\gamma\delta$  cells and an increase in T  $\gamma\delta$  lymphocyte expansion in the lungs of aged mice as well as a reduced T $\gamma\delta$  lymphocyte activation state after application of KD to mCoV-A59-infected male mice reproducing the clinical features of COVID-19 for 5 days prior to infection. Application of KD resulted in a significant increase in genes associated with reduced inflammation, increased lipoprotein remodeling and downregulation of TLR, Plk1, and aurora B signaling pathways in T $\gamma\delta$  lymphocytes. Lung T $\gamma\delta$  lymphocytes from mCoV-A59-infected mice treated with KD showed increased respiratory electron transport and complex I biogenesis. In addition, Golgi retrograde transport to the ER and cell cycle were downregulated, suggesting a reduced activation status of  $\gamma\delta$  T cells. The results suggest that T $\gamma\delta$  T cells proliferated with KD are functionally more homeostatic and immunologically protective against mCoV-A59 infection mimicking SARS-CoV-2. This confirmed previous observations associated with the use of KD in influenza A virus infection, resulting in the expansion of T $\gamma\delta$  cells in the lungs of mice, improved immune barrier function, and lung barrier integrity [187–189], and thereby increased antiviral resistance. Importantly, T $\gamma\delta$  cell expansion required metabolic adaptation to KD, whereas feeding mice a high-fat, high-carbohydrate diet or providing ketone bodies did not protect against infection [34].

#### 4.2.7. KD and Ketogenesis Activation Inactivates the NLRP3 Inflammasome

The NLRP3 inflammasome [190,191], an essential component of the innate immune system, is hyperactivated in response to SARS-CoV-2 infection [40,192]. Hyperactivation of the NLRP3 inflammasome results in increased levels of caspase-1 responsible for inducing the secretion of pro-inflammatory cytokines (IL-1 $\beta$ , IL-18, IL-6, and TNF- $\alpha$ ) [191,193,194]. Abnormal activation of the NLRP3 inflammasome in COVID-19 is involved in the pathogenesis of cytokine storm [190,195], acute respiratory distress syndrome (ARDS) [193,196], and acute lung inflammation (ALI) [197], as well as pathological multi-organ damage [194,198].

Ryu et al. [199], in a study with mice and mouse macrophages after mCoV-A59 infection replicating the clinical features of COVID-19, observed that administration of a ketogenic diet for 5 days in an aging mouse model of mCoV-A59 infection, significantly decreased NLRP3 mRNA and caspase-1 in the lung, visceral adipose tissue (VAT), and hypothalamus, and inflammasome activation in the VAT. BHB also reduced oligomerization of ASC, an adaptor protein required for the assembly of the inflammasome complex. This confirms previous observations by Youm et al. [36], who showed in a mouse model of inflammatory disease that BHB caused mechanistic inhibition of NLRP3 inflammasome activation and also reduced interleukin (IL)-1 $\beta$  and IL-18 production in human monocytes. In vivo, BHB or a ketogenic diet attenuated caspase-1 activation and IL-1 $\beta$  secretion in mouse models of diseases mediated by NLRP3.

Thus, the NLRP3 inflammasome may be a potential target for the treatment of COVID-19 [191,194], which is strongly supported by studies showing improved saturation, reduced hospitalization time and reduced mortality in patients with COVID-19 after treatment with NLRP3 inflammasome inhibitors such as Tranilast [200] or colchicine [198]. The anti-inflammatory effect of metformin in COVID-19 was also associated with inhibition of NLRP3 inflammasome activation [196].

#### 4.2.8. BHB Prevents CD8+ T-Lymphocyte Dysfunction

CD8+ T lymphocytes contribute to protective immune responses against SARS-CoV-2 [201]. They play an essential role in controlling viral infection by lysing virus-infected cells and producing effector cytokines [202]. However, in severe COVID-19, profound dysfunction of CD8+ lymphocytes has been observed involving their over-activation, reduced absolute number, and functionality [201,203–208].

In a study by Hirschberger et al. [143] on immune cells from patients with severe COVID-19 infection, application of BHB resulted in an increase in CD8+ lymphocytes and a significant enhancement of CD8+ immune capacity, and increased secretion of CD8+ T-cell cytokines (IFN $\gamma$ , TNF $\alpha$ , perforin, and granzyme B). They also showed that CD8+ lymphocytes had a significantly increased capacity for cell lysis after BHB treatment. Thus, administration of BHB significantly enhanced the CD8+ T-lymphocyte functions impaired by SARS-CoV-2 infection in severely ill patients. In addition, under the influence of BHB, there was a tendency to increase mitochondrial mass in CD8+ T cells and redirect CD8+ T cells towards aerobic mitochondrial metabolism allowing for better energy supply. This reprogramming of CD8+ cells by BHB may be a promising strategy to overcome immune paralysis in patients with COVID-19. These findings confirmed previous observations on primary human cells and from an interventional immunological study with healthy volunteers, where both ketone bodies and a ketogenic diet prevented CD8+ T-cell dysfunction through immunometabolic reprogramming [156].

#### 4.2.9. BHB and KD Prevent Cell Death via the Pseudo Receptor Pathway (by Perforin and Granzymes)

In the peripheral blood mononuclear cells of patients with severe COVID-19, in addition to CD8+ lymphocyte dysfunction, significantly higher granzyme B and perforin expression was observed in CD8+ lymphocytes compared with mild disease [162]. Incubation of peripheral blood mononuclear cells obtained from patients with high COVID-19 severity and respiratory failure with BHB resulted in a significant increase in granzyme B-expressing CD8+ lymphocytes, increased granzyme B expression per cell, significantly increased secretion of perforin and granzyme B, and high cell lysis capacity [143]. This demonstrates the enhanced ability of CD8+ lymphocytes after BHB treatment to eliminate the virus from the host [209–212].

Perforins polymerize to form a channel that allows the free flow of ions and small polypeptides into the cell. The result is cell instability, impaired osmoticity, loss of energy and destruction of genetic material and, ultimately, pseudo receptor-mediated cell destruction [213].

In the absence of perforin, high viral titers activate CD8<sup>+</sup> lymphocytes, maintain high CD8 T-cell numbers for long periods of time, and produce high levels of IFN- $\gamma$  but are unable to clear the infection [214]. Cunningham et al. [212] suggested a link between perforin expression and COVID-19 resistance.

The role of granzymes, on the other hand, is to activate perforin polymerization, penetrating through perforin channels into the target cell, where they degrade integral structure proteins, cytosolic proteins, and cell organelles. They also bind to and degrade chromatin proteins. This leads to the exposure of DNA to endonucleases and its destruction [213].

#### 4.2.10. BHB as a Substrate for Histone H3K9me3

Xie et al. [215] showed that  $\beta$ -Hydroxybutyrate acts as a substrate for histone H3K9me3 (a post-translational modification of the protein), which then activates gene expression, independent of acetylation. In an unreviewed study on simulated H3K9me3 antibodies produced using in vitro chemical modification from a commercially obtained human serum antibody treated with  $\beta$ -Hydroxybutyrate, it was shown that H3K9me3 did not affect the binding capacity of B38 to the S protein of COVID-19 virus, indicating that the chemically modified antibody retained its original antigen-binding activity [144].

#### 4.3. BHB Increased Mitochondrial mROS, but without Uncontrolled ROS Expansion

In many pathological conditions, increased mitochondrial oxidative phosphorylation translates into an increase in mitochondrial reactive oxygen species (mROS) [216,217]. At the same time, mROS are very important for the homeostasis of the body [218] including an adequate T-cell immune response [219]. Nine times higher ROS production was observed in patients with COVID-19 compared to healthy subjects, and was particularly high in patients requiring mechanical ventilation [220].

Hirschberger et al. [143], in a study on cells obtained from COVID-19 patients incubated with BHB, found that mitochondrial ROS were significantly elevated after incubation with ketone bodies, indicating an increased immune capacity of CD8<sup>+</sup> cells. At the same time, cellular ROS levels remained unchanged, indicating that there was no uncontrolled ROS expansion that could impair cell viability.

#### 4.4. KD in Animal Models Resulted in Protection against Weight Loss and Hypoxemia, Faster Recovery, Reduced Lung Damage, and Resulted in Better Survival of Young Mice

In animal studies, Ryu et al. [42] also showed that KD-fed mice infected with mCoV-A59 were protected against weight loss and hypoxemia caused by the infection. Compared to the control group, young mice infected with a lethal dose of mCoV-A59 (PFU 1e6) and fed KD before infection showed better survival in contrast to old mice, which were not protected from death by KD. KD also significantly reduced myeloid cell infiltration in the heart.

Karagannis et al. [116], in a study with mice in a preclinical model of SARS-CoV-2 infection, found faster recovery from weight loss due to infection and also reduced lung damage, resulting in increased overall survival when mice were fed KD or ketone ester.

#### 4.5. KD in Humans Increased Survival, Reduced the Need for Hospitalization for COVID-19, and Showed a Protective Role against Metabolic Abnormalities after COVID-19

Previous studies have shown that the use of KD resulted in a milder course and lower mortality in patients with COVID-19 and also reduced obesity.

Sukkar et al. [14] in a pilot study showed higher survival and significantly less need for ICU hospitalization in COVID-19 patients following a eucaloric ketogenic diet (EKD 1800–2100, carbohydrates <30 g, 5–6% of total energy) from admission compared with patients following a standard diet. The results of this study indicated for the first time the role of clinical KD-based nutritional therapy as a pathophysiological support for drug therapy in improving the prognosis of COVID-19. The EKD used proved to be safe and no adverse effects were observed.

In a retrospective study, Volk et al. [180] demonstrated a significant reduction in body weight in patients with type 2 diabetes and obesity who used KD (5–10% energy in the form of carbohydrate, 15–20% protein, and 70–75% fat) before COVID-19 diagnosis and during treatment. Both obesity and type 2 diabetes predispose to a more severe course of COVID-19. Obesity through common pathophysiological features with COVID-19, such as chronic inflammation, immune dysregulation, oxidative stress, increased cytokine production, and endothelial dysfunction worsens the course of the disease. An additional increased number of adipocytes express ACE2 and increased adipose tissue constitutes a larger viral reservoir [44,221]. The introduction of KD through weight reduction resulted in reduced inflammation and thus a lighter course of COVID-19 and less need for hospitalization. KD, by limiting carbohydrate supply and reducing glucose and insulin levels, may also have reversed the dysfunction of the immune response that increases the risk of spreading invasive pathogens in patients with diabetes [222].

Reductions in body weight, body mass index (BMI), amount of body fat with preservation of lean mass, and improvements in muscle strength and physical performance were observed by Camajani et al. [179] in a woman with sarcopenic obesity (SO) and dyslipidemia, following hospitalization for bilateral interstitial pneumonia due to COVID-19, using VLCKD (800 kcal/day, carbohydrate 28 g (14.6%), fat 35 g (38.7%), and protein 85 g (46.7%)) combined with interval training for 6 weeks after treatment. Given that SO exacerbates the decline in skeletal muscle mass and strength, in turn, the reduced muscle mass is significantly associated with the occurrence of complications after COVID-19 [223], and the use of VLCKD reversed the expected adverse outcomes.

In the same patient with SO, a decrease in blood pressure, insulin, fasting glycemia, HOMA index, triglycerides, and LDL cholesterol was observed with a concomitant increase in HDL cholesterol [179]. These findings suggest a protective role of KD against metabolic abnormalities after COVID-19.

Previous studies have indicated beneficial metabolic effects of KD use, with COVID-19 diabetes (type 1, type 2) [224,225], hypertension, and cardiovascular disease [1,225] being associated with an increased risk of severe disease and higher mortality [224]. Thus, improving parameters related to the course of these diseases may reduce the severity of the course and mortality of COVID-19 [226–228].

#### 4.6. Future Research Directions

Many years of research using ketogenic diets to treat epilepsy, obesity, and type 2 diabetes have demonstrated the safety of KD. However, they do have some limitations as clinical interventions, including the difficulty in implementing large-scale dietary changes and strict patient adherence to these recommendations. Despite the encouraging results, it is important to note the limitations of the studies analyzed such as the small sample size and lack of randomized controlled trials, which may affect the generalizability of the results. Furthermore, animal studies should be extrapolated to humans with great caution.

To confirm the efficacy of KD/ketone bodies in the prevention and treatment of COVID-19, randomized clinical trials involving COVID-19 patients are needed, with assessment of the duration and severity of the course of infection and patient mortality. These trials should analyze the outcomes of patients at different stages of COVID-19 disease progression following the introduction of KD or ketone bodies therapy, which would identify the time at which such an intervention would be most effective and eliminate groups of patients at risk of developing ketoacidosis.

Soliman et al. [43], to support the treatment of SARS-CoV-2, suggested altering host lipid metabolism through the use of KD combined with fasting consisting of a ketogenic breakfast combined with medium-chain fatty acid (MCT) supplementation from coconut oil, a lunch with MCT inclusion, an 8–12 h fast and a dinner rich in fruit and vegetables. Cooper et al. [229] in a study involving patients with COVID-19 and a history of hyperglycemia and/or hyperinsulinemia proposed an algorithm of restricted carbohydrate management, administered either enterally or parenterally, restricting carbohydrates and intravenous

solutions containing dextrose in the diet to a minimum. Restriction of the introduced diet would be dependent on continuously monitored blood glucose, insulin, and ketone bodies. The algorithm should include supplementation with vitamin D, magnesium, and zinc.

Although the results obtained to date do not allow a clear recommendation for the use of KD and/or ketone bodies for the treatment of COVID-19, a model of nutritional management using KD can be proposed, which would involve a low-carbohydrate diet that increases blood ketones throughout the course of the disease, while monitoring patient outcomes and adjusting carbohydrate restriction according to the course of the disease. This would maintain normal blood glucose levels and stimulate immune cell function in the early stages, reduce glucose metabolism and associated pro-inflammatory signaling during symptom severity, and promote anti-inflammatory processes when respiratory difficulties arise. Regardless of the timing of the introduction of KD or ketone bodies therapy, it could have the effect of reducing the incidence of progression to ARDS, protecting organs from oxidative and inflammatory damage, and shortening the duration of the disease. The use of KD could also have a positive effect in patients who have survived severe ARDS causing psychiatric problems and chronic fatigue [3] due to the mitigation of cell death and tissue damage.

## 5. Conclusions

There is a wealth of scientific evidence on the effectiveness of nutritional interventions in disease prevention and treatment. It is also known that a good nutritional status of the body has a great impact on the course of COVID-19. However, to date, the best nutritional approach for use during the SARS-CoV-2 pandemic has not been established. This review examines the evidence on the areas of efficacy of the use of ketogenic diets or ketone bodies for prevention of or during COVID-19 infection in animal models, in humans, and in tissue models. The efficacy of KD or ketone bodies at the stage of SARS-CoV-2 virus entry into the cell has been shown to result from (a) KD decreasing ACE2 methylation and increasing ACE2 mRNA expression in visceral adipose tissue and leukocytes, downregulation of ACE2 and TMPRSS2 key elements used by SARS-CoV-2 to infect cells in the lung, and (b) a shift of the renin–angiotensin–aldosterone system (RAS) from the AngI/ACE1/AngII/AT1 arm towards the anti-inflammatory RAS arm. The efficacy of KD/ketone bodies in the regulation of innate and acquired immunity was also confirmed by (a) BHB reducing cellular dependence on glucose, (b) BHB promoting mitochondrial function and improving respiratory chain function, which could become an alternative carbon source to OXPHOS, (c) BHB preventing CD4+ and CD8+ T-lymphocyte dysfunction, (d) KD and activation of ketogenesis reducing the number of pathogenic monocytes in the lungs, (e) KD inducing a slight decrease in IL-6 levels in humans and decreased mRNA expression of the pro-inflammatory cytokines IL-1 $\beta$ , TNF $\alpha$ , and IL-6, the inflammatory genes TLR4 and ILR6, and inhibiting TNFR1 gene expression in mice, (f) increased homeostaticity of T  $\gamma\delta$  lymphocytes, (g) inactivation of the NLRP3 inflammasome, and (h) prevention of cell death via the pseudo receptor pathway (via perforin and granzymes). At the same time, BHB increased mitochondrial mROS, but without uncontrolled ROS expansion. In animal models, KD resulted in protection against weight loss and hypoxemia, faster recovery, reduced lung damage, and better survival of young mice. In humans, KD also increased survival, reduced the need for COVID-19 hospitalization, and showed a protective role against metabolic disorders after COVID-19.

In conclusion, it can be stated that the multidirectional beneficial effects of KD and ketone bodies pave the way for research into metabolic treatment in COVID-19 [4]; however, the selection of KD effective in the prevention and/or treatment of COVID-19 is fraught with problems and requires further studies that would not only confirm its efficacy but also identify the best time to apply it during infection, define the KD model, and identify contraindications for the use of such an intervention against COVID-19.

**Author Contributions:** Conceptualization, I.B.; formal analysis, J.P.; resources, I.B. and N.B.-K.; writing—original draft preparation, I.B.; writing—review and editing, M.K., N.B.-K. and J.P.; visualization, I.B.; project administration, I.B.; funding acquisition, J.P. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Data sharing is not applicable to this article.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Stubbs, B.J.; Koutnik, A.P.; Goldberg, E.L.; Upadhyay, V.; Turnbaugh, P.J.; Verdin, E.; Newman, J.C. Investigating Ketone Bodies as Immunometabolic Countermeasures against Respiratory Viral Infections. *Med* **2020**, *1*, 43–65. [\[CrossRef\]](#)
2. Fagbule, O.F. 2019 Novel Coronavirus. *Ann. Ib. Postgrad. Med.* **2019**, *17*, 108–110.
3. Gupta, P.; Goyal, K.; Kanta, P.; Ghosh, A.; Singh, M.P. Novel 2019-Coronavirus on New Year's Eve. *Indian J. Med. Microbiol.* **2019**, *37*, 459–477. [\[CrossRef\]](#)
4. Wen, W.; Chen, C.; Tang, J.; Wang, C.; Zhou, M.; Cheng, Y.; Zhou, X.; Wu, Q.; Zhang, X.; Feng, Z.; et al. Efficacy and Safety of Three New Oral Antiviral Treatment (Molnupiravir, Fluvoxamine and Paxlovid) for COVID-19: A Meta-Analysis. *Ann. Med.* **2022**, *54*, 516–523. [\[CrossRef\]](#)
5. Calvo Fernández, E.; Zhu, L.Y. Racing to Immunity: Journey to a COVID-19 Vaccine and Lessons for the Future. *Br. J. Clin. Pharmacol.* **2021**, *87*, 3408–3424. [\[CrossRef\]](#)
6. *Coronavirus Disease 2019 (COVID-19) Treatment Guidelines*; National Institutes of Health: Bethesda, MD, USA, 2021.
7. Fiolet, T.; Kherabi, Y.; MacDonald, C.-J.; Ghosn, J.; Peiffer-Smadja, N. Comparing COVID-19 Vaccines for Their Characteristics, Efficacy and Effectiveness against SARS-CoV-2 and Variants of Concern: A Narrative Review. *Clin. Microbiol. Infect.* **2022**, *28*, 202–221. [\[CrossRef\]](#)
8. Hadj Hassine, I. COVID-19 Vaccines and Variants of Concern: A Review. *Rev. Med. Virol.* **2022**, *32*, e2313. [\[CrossRef\]](#)
9. Rehman, S.U.; Rehman, S.U.; Yoo, H.H. COVID-19 Challenges and Its Therapeutics. *Biomed. Pharmacother.* **2021**, *142*, 112015. [\[CrossRef\]](#)
10. Parums, D.V. Editorial: Current Status of Oral Antiviral Drug Treatments for SARS-CoV-2 Infection in Non-Hospitalized Patients. *Med. Sci. Monit.* **2022**, *28*, e935952. [\[CrossRef\]](#)
11. Calder, P.C. Nutrition, Immunity and COVID-19. *BMJ Nutr. Prev. Health* **2020**, *3*, 74–92. [\[CrossRef\]](#)
12. Iddir, M.; Brito, A.; Dingeo, G.; Fernandez Del Campo, S.S.; Samouda, H.; La Frano, M.R.; Bohn, T. Strengthening the Immune System and Reducing Inflammation and Oxidative Stress through Diet and Nutrition: Considerations during the COVID-19 Crisis. *Nutrients* **2020**, *12*, 1562. [\[CrossRef\]](#)
13. Zabetakis, I.; Lordan, R.; Norton, C.; Tsoupras, A. COVID-19: The Inflammation Link and the Role of Nutrition in Potential Mitigation. *Nutrients* **2020**, *12*, 1466. [\[CrossRef\]](#)
14. Sukkar, S.G.; Cogorno, L.; Pisciotta, L.; Pasta, A.; Vena, A.; Gradasci, R.; Dentone, C.; Guidido, E.; Martino, E.; Beltrami, S.; et al. Clinical Efficacy of Eucaloric Ketogenic Nutrition in the COVID-19 Cytokine Storm: A Retrospective Analysis of Mortality and Intensive Care Unit Admission. *Nutrition* **2021**, *89*, 111236. [\[CrossRef\]](#)
15. Poff, A.M.; Rho, J.M.; D'Agostino, D.P. Ketone Administration for Seizure Disorders: History and Rationale for Ketone Esters and Metabolic Alternatives. *Front. Neurosci.* **2019**, *13*, 1041. [\[CrossRef\]](#)
16. Zarnowska, I.M. Therapeutic Use of the Ketogenic Diet in Refractory Epilepsy: What We Know and What Still Needs to Be Learned. *Nutrients* **2020**, *12*, 2616. [\[CrossRef\]](#)
17. Ułamek-Kozioł, M.; Czuczwar, S.J.; Januszewski, S.; Pluta, R. Ketogenic Diet and Epilepsy. *Nutrients* **2019**, *11*, 2510. [\[CrossRef\]](#)
18. Masino, S.A.; Rho, J.M. Metabolism and Epilepsy: Ketogenic Diets as a Homeostatic Link. *Brain Res.* **2019**, *1703*, 26–30. [\[CrossRef\]](#)
19. Fan, Y.; Wang, H.; Liu, X.; Zhang, J.; Liu, G. Crosstalk between the Ketogenic Diet and Epilepsy: From the Perspective of Gut Microbiota. *Mediat. Inflamm.* **2019**, *2019*, 8373060. [\[CrossRef\]](#)
20. Włodarek, D. Role of Ketogenic Diets in Neurodegenerative Diseases (Alzheimer's Disease and Parkinson's Disease). *Nutrients* **2019**, *11*, 169. [\[CrossRef\]](#)
21. Jensen, N.J.; Wodschow, H.Z.; Nilsson, M.; Rungby, J. Effects of Ketone Bodies on Brain Metabolism and Function in Neurodegenerative Diseases. *Int. J. Mol. Sci.* **2020**, *21*, 8767. [\[CrossRef\]](#)
22. Ludwig, D.S. The Ketogenic Diet: Evidence for Optimism but High-Quality Research Needed. *J. Nutr.* **2020**, *150*, 1354–1359. [\[CrossRef\]](#)
23. Neth, B.J.; Mintz, A.; Whitlow, C.; Jung, Y.; Solingapuram Sai, K.; Register, T.C.; Kellar, D.; Lockhart, S.N.; Hoscheidt, S.; Maldjian, J.; et al. Modified Ketogenic Diet Is Associated with Improved Cerebrospinal Fluid Biomarker Profile, Cerebral

- Perfusion, and Cerebral Ketone Body Uptake in Older Adults at Risk for Alzheimer's Disease: A Pilot Study. *Neurobiol. Aging* **2020**, *86*, 54–63. [[CrossRef](#)]
24. Phillips, M.C.L.; Murtagh, D.K.J.; Gilbertson, L.J.; Asztely, F.J.S.; Lynch, C.D.P. Low-Fat versus Ketogenic Diet in Parkinson's Disease: A Pilot Randomized Controlled Trial. *Mov. Disord.* **2018**, *33*, 1306–1314. [[CrossRef](#)]
25. Bruci, A.; Tuccinardi, D.; Tozzi, R.; Balena, A.; Santucci, S.; Frontani, R.; Mariani, S.; Basciani, S.; Spera, G.; Gnessi, L.; et al. Very Low-Calorie Ketogenic Diet: A Safe and Effective Tool for Weight Loss in Patients with Obesity and Mild Kidney Failure. *Nutrients* **2020**, *12*, 333. [[CrossRef](#)]
26. Goday, A.; Bellido, D.; Sajoux, I.; Crujeiras, A.B.; Burguera, B.; García-Luna, P.P.; Oleaga, A.; Moreno, B.; Casanueva, F.F. Short-Term Safety, Tolerability and Efficacy of a Very Low-Calorie-Ketogenic Diet Interventional Weight Loss Program versus Hypocaloric Diet in Patients with Type 2 Diabetes Mellitus. *Nutr. Diabetes* **2016**, *6*, e230. [[CrossRef](#)]
27. Hall, K.D.; Bemis, T.; Brychta, R.; Chen, K.Y.; Courville, A.; Crayner, E.J.; Goodwin, S.; Guo, J.; Howard, L.; Knuth, N.D.; et al. Calorie for Calorie, Dietary Fat Restriction Results in More Body Fat Loss than Carbohydrate Restriction in People with Obesity. *Cell Metab.* **2015**, *22*, 427–436. [[CrossRef](#)]
28. Cohen, C.W.; Fontaine, K.R.; Arend, R.C.; Alvarez, R.D.; Leath, C.A.; Huh, W.K.; Bevis, K.S.; Kim, K.H.; Straughn, J.M.; Gower, B.A. A Ketogenic Diet Reduces Central Obesity and Serum Insulin in Women with Ovarian or Endometrial Cancer. *J. Nutr.* **2018**, *148*, 1253–1260. [[CrossRef](#)]
29. Harvey, K.L.; Holcomb, L.E.; Kolwicz, S.C., Jr. Ketogenic Diets and Exercise Performance. *Nutrients* **2019**, *11*, 2296. [[CrossRef](#)]
30. Kosinski, C.; Jornayvaz, F.R. Effects of Ketogenic Diets on Cardiovascular Risk Factors: Evidence from Animal and Human Studies. *Nutrients* **2017**, *9*, 517. [[CrossRef](#)]
31. Merra, G.; Gratteri, S.; De Lorenzo, A.; Barrucco, S.; Perrone, M.A.; Avolio, E.; Bernardini, S.; Marchetti, M.; Di Renzo, L. Effects of Very-Low-Calorie Diet on Body Composition, Metabolic State, and Genes Expression: A Randomized Double-Blind Placebo-Controlled Trial. *Eur. Rev. Med. Pharmacol. Sci.* **2017**, *21*, 329–345.
32. Hagihara, K.; Kajimoto, K.; Osaga, S.; Nagai, N.; Shimosegawa, E.; Nakata, H.; Saito, H.; Nakano, M.; Takeuchi, M.; Kanki, H.; et al. Promising Effect of a New Ketogenic Diet Regimen in Patients with Advanced Cancer. *Nutrients* **2020**, *12*, 1473. [[CrossRef](#)]
33. Khodabakhshi, A.; Akbari, M.E.; Mirzaei, H.R.; Mehrad-Majd, H.; Kalamian, M.; Davoodi, S.H. Feasibility, Safety, and Beneficial Effects of MCT-Based Ketogenic Diet for Breast Cancer Treatment: A Randomized Controlled Trial Study. *Nutr. Cancer* **2020**, *72*, 627–634. [[CrossRef](#)]
34. Goldberg, E.L.; Molony, R.D.; Kudo, E.; Sidorov, S.; Kong, Y.; Dixit, V.D.; Iwasaki, A. Ketogenic Diet Activates Protective  $\Gamma\delta$  T Cell Responses against Influenza Virus Infection. *Sci. Immunol.* **2019**, *4*, eaav2026. [[CrossRef](#)]
35. Qiao, G.; Chen, P.; Sun, Q.; Zhang, M.; Zhang, J.; Li, Z.; Li, Q. Poly- $\beta$ -Hydroxybutyrate (PHB) in Bioflocs Alters Intestinal Microbial Community Structure, Immune-Related Gene Expression and Early Cyprinid Herpesvirus 2 Replication in Gibel Carp (*Carassius Auratus* Gibelio). *Fish. Shellfish. Immunol.* **2020**, *97*, 72–82. [[CrossRef](#)]
36. Youm, Y.-H.; Nguyen, K.Y.; Grant, R.W.; Goldberg, E.L.; Bodogai, M.; Kim, D.; D'Agostino, D.; Planavsky, N.; Lupfer, C.; Kanneganti, T.D.; et al. The Ketone Metabolite  $\beta$ -Hydroxybutyrate Blocks NLRP3 Inflammasome-Mediated Inflammatory Disease. *Nat. Med.* **2015**, *21*, 263–269. [[CrossRef](#)]
37. Goldberg, E.L.; Asher, J.L.; Molony, R.D.; Shaw, A.C.; Zeiss, C.J.; Wang, C.; Morozova-Roche, L.A.; Herzog, R.L.; Iwasaki, A.; Dixit, V.D.  $\beta$ -Hydroxybutyrate Deactivates Neutrophil NLRP3 Inflammasome to Relieve Gout Flares. *Cell Rep.* **2017**, *18*, 2077–2087. [[CrossRef](#)]
38. Trotta, M.C.; Maisto, R.; Guida, F.; Boccella, S.; Luongo, L.; Balta, C.; D'Amico, G.; Herman, H.; Hermenean, A.; Bucolo, C.; et al. The Activation of Retinal HCA2 Receptors by Systemic Beta-Hydroxybutyrate Inhibits Diabetic Retinal Damage through Reduction of Endoplasmic Reticulum Stress and the NLRP3 Inflammasome. *PLoS ONE* **2019**, *14*, e0211005. [[CrossRef](#)]
39. Qian, J.; Zhu, W.; Lu, M.; Ni, B.; Yang, J. D- $\beta$ -hydroxybutyrate Promotes Functional Recovery and Relieves Pain Hypersensitivity in Mice with Spinal Cord Injury. *Br. J. Pharmacol.* **2017**, *174*, 1961–1971. [[CrossRef](#)]
40. Paoli, A.; Gorini, S.; Caprio, M. The Dark Side of the Spoon—Glucose, Ketones and COVID-19: A Possible Role for Ketogenic Diet? *J. Transl. Med.* **2020**, *18*, 441. [[CrossRef](#)]
41. Bradshaw, P.C.; Seeds, W.A.; Miller, A.C.; Mahajan, V.R.; Curtis, W.M. COVID-19: Proposing a Ketone-Based Metabolic Therapy as a Treatment to Blunt the Cytokine Storm. *Oxid. Med. Cell. Longev.* **2020**, *2020*, 6401341. [[CrossRef](#)]
42. Ryu, S.; Shchukina, I.; Youm, Y.-H.; Qing, H.; Hilliard, B.K.; Dlugos, T.; Zhang, X.; Yasumoto, Y.; Booth, C.J.; Fernández-Hernando, C.; et al. Ketogenesis Restrains Aging-Induced Exacerbation of COVID in a Mouse Model. *bioRxiv* **2020**. [[CrossRef](#)]
43. Soliman, S.; Faris, M.E.; Ratemi, Z.; Halwani, R. Switching Host Metabolism as an Approach to Dampen SARS-CoV-2 Infection. *Ann. Nutr. Metab.* **2020**, *76*, 297–303. [[CrossRef](#)]
44. Gangitano, E.; Tozzi, R.; Gandini, O.; Watanabe, M.; Basciani, S.; Mariani, S.; Lenzi, A.; Gnessi, L.; Lubrano, C. Ketogenic Diet as a Preventive and Supportive Care for COVID-19 Patients. *Nutrients* **2021**, *13*, 1004. [[CrossRef](#)]
45. Shaheen, A. Can Ketone Bodies Inactivate Coronavirus Spike Protein? The Potential of Biocidal Agents against SARS-CoV-2. *Bioessays* **2021**, *43*, e2000312. [[CrossRef](#)]

46. Dąbek, A.; Wojtala, M.; Pirola, L.; Balcerzyk, A. Modulation of Cellular Biochemistry, Epigenetics and Metabolomics by Ketone Bodies. Implications of the Ketogenic Diet in the Physiology of the Organism and Pathological States. *Nutrients* **2020**, *12*, 788. [CrossRef]
47. Ungaro, P.; Nettore, I.C.; Franchini, F.; Palatucci, G.; Muscogiuri, G.; Colao, A.; Macchia, P.E. Epigenome Modulation Induced by Ketogenic Diets. *Nutrients* **2022**, *14*, 3245. [CrossRef]
48. Alhamzah, S.A.; Gatar, O.M.; Alruwaili, N.W. Effects of Ketogenic Diet on Oxidative Stress and Cancer: A Literature Review. *Adv. Cancer Biol.-Metastasis* **2023**, *7*, 100093. [CrossRef]
49. Cannataro, R.; Caroleo, M.C.; Fazio, A.; La Torre, C.; Plastina, P.; Gallelli, L.; Lauria, G.; Cione, E. Ketogenic Diet and MicroRNAs Linked to Antioxidant Biochemical Homeostasis. *Antioxidants* **2019**, *8*, 269. [CrossRef]
50. Green, A.; Bishop, R.E. Ketoacidosis—Where Do the Protons Come From? *Trends Biochem. Sci.* **2019**, *44*, 484–489. [CrossRef]
51. Nagdev, G.; Chavan, G.; Gadkari, C.; Sahu, G. Diabetic Ketoacidosis and COVID-19: A Retrospective Observational Study. *Cureus* **2022**, *14*, e30895. [CrossRef]
52. de Sá-Ferreira, C.O.; da Costa, C.H.M.; Guimarães, J.C.W.; Sampaio, N.S.; Silva, L.d.M.L.; de Mascarenhas, L.P.; Rodrigues, N.G.; dos Santos, T.L.; Campos, S.; Young, E.C. Diabetic Ketoacidosis and COVID-19: What Have We Learned so Far? *Am. J. Physiol. Endocrinol. Metab.* **2022**, *322*, E44–E53. [CrossRef]
53. Santharaman, A.; Raj, K.; Sankaramangalam, K.; Dewan, S.; Sapkota, S.; Chandna, S.; Shah, M.; Singh, D.; Bassi, M.; Luo, H.; et al. COVID-19 Related Predisposition to Diabetic Ketoacidosis. *Cureus* **2023**, *15*, e36674. [CrossRef]
54. Ghavasieh, A.; Bontorin, S.; Artime, O.; Verstraete, N.; De Domenico, M. Multiscale Statistical Physics of the Pan-Viral Interactome Unravels the Systemic Nature of SARS-CoV-2 Infections. *Commun. Phys.* **2021**, *4*, 83. [CrossRef]
55. Kim, K.W.; Deveson, I.W.; Pang, C.N.I.; Yeang, M.; Naing, Z.; Adikari, T.; Hammond, J.M.; Stevanovski, I.; Beukers, A.G.; Verich, A.; et al. Respiratory Viral Co-Infections among SARS-CoV-2 Cases Confirmed by Virome Capture Sequencing. *Sci. Rep.* **2021**, *11*, 3934. [CrossRef]
56. CSR MERS Outbreaks. Available online: <http://www.emro.who.int/health-topics/mers-cov/mers-outbreaks.html> (accessed on 15 May 2023).
57. Countries and Territories with Current or Previous Zika Virus Transmission. Available online: <https://www.who.int/publications/m/item/countries-and-territories-with-current-or-previous-zika-virus-transmission> (accessed on 15 May 2023).
58. Carabelli, A.M.; Peacock, T.P.; Thorne, L.G.; Harvey, W.T.; Hughes, J.; de Silva, T.I.; Peacock, S.J.; Barclay, W.S.; de Silva, T.I.; Towers, G.J.; et al. SARS-CoV-2 Variant Biology: Immune Escape, Transmission and Fitness. *Nat. Rev. Microbiol.* **2023**, *21*, 162–177. [CrossRef]
59. Mohamadian, M.; Chiti, H.; Shoghli, A.; Biglari, S.; Parsamanesh, N.; Esmaeilzadeh, A. COVID-19: Virology, biology and novel laboratory diagnosis. *J. Gene Med.* **2021**, *23*, e3303. [CrossRef]
60. Samudrala, P.K.; Kumar, P.; Choudhary, K.; Thakur, N.; Wadekar, G.S.; Dayaramani, R.; Agrawal, M.; Alexander, A. Virology, Pathogenesis, Diagnosis and in-Line Treatment of COVID-19. *Eur. J. Pharmacol.* **2020**, *883*, 173375. [CrossRef]
61. Velusamy, P.; Kiruba, K.; Su, C.-H.; Arun, V.; Anbu, P.; Gopinath, S.C.B.; Vaseeharan, B. SARS-CoV-2 Spike Protein: Site-Specific Breakpoints for the Development of COVID-19 Vaccines. *J. King Saud. Univ. Sci.* **2021**, *33*, 101648. [CrossRef]
62. Rasmussen, S.A.; Smulian, J.C.; Lednický, J.A.; Wen, T.S.; Jamieson, D.J. Coronavirus Disease 2019 (COVID-19) and Pregnancy: What Obstetricians Need to Know. *Am. J. Obstet. Gynecol.* **2020**, *222*, 415–426. [CrossRef]
63. dos Santos, W.G. Natural History of COVID-19 and Current Knowledge on Treatment Therapeutic Options. *Biomed. Pharmacother.* **2020**, *129*, 110493. [CrossRef]
64. Todorova, V.; Ivanov, K.; Ivanova, S. Comparison between the Biological Active Compounds in Plants with Adaptogenic Properties (*Rhaponticum Carthamoides*, *Lepidium Meyenii*, *Eleutherococcus Senticosus* and *Panax Ginseng*). *Plants* **2021**, *11*, 64. [CrossRef]
65. Papa, G.; Mallery, D.L.; Albecka, A.; Welch, L.G.; Cattin-Ortolá, J.; Luptak, J.; Paul, D.; McMahon, H.T.; Goodfellow, I.G.; Carter, A.; et al. Furin cleavage of SARS-CoV-2 Spike promotes but is not essential for infection and cell-cell fusion. *PLoS Pathog.* **2021**, *17*, e1009246. [CrossRef]
66. Kadam, S.B.; Sukhramani, G.S.; Bishnoi, P.; Pable, A.A.; Barvkar, V.T. SARS-CoV-2, the Pandemic Coronavirus: Molecular and Structural Insights. *J. Basic. Microbiol.* **2021**, *61*, 180–202. [CrossRef]
67. Scialo, F.; Daniele, A.; Amato, F.; Pastore, L.; Matera, M.G.; Cazzola, M.; Castaldo, G.; Bianco, A. ACE2: The Major Cell Entry Receptor for SARS-CoV-2. *Lung* **2020**, *198*, 867–877. [CrossRef]
68. Jackson, C.B.; Farzan, M.; Chen, B.; Choe, H. Mechanisms of SARS-CoV-2 Entry into Cells. *Nat. Rev. Mol. Cell. Biol.* **2022**, *23*, 3–20. [CrossRef]
69. Trougakos, I.P.; Stamatelopoulos, K.; Terpos, E.; Tsitsilonis, O.E.; Aivalioti, E.; Paraskevis, D.; Kastritis, E.; Pavlakis, G.N.; Dimopoulos, M.A. Insights to SARS-CoV-2 Life Cycle, Pathophysiology, and Rationalized Treatments That Target COVID-19 Clinical Complications. *J. Biomed. Sci.* **2021**, *28*, 9. [CrossRef]
70. Osman, E.E.A.; Rehemtulla, A.; Neamati, N. Why All the Fury over Furin? *J. Med. Chem.* **2022**, *65*, 2747–2784. [CrossRef]
71. Lippi, G.; Lavie, C.J.; Henry, B.M.; Sanchis-Gomar, F. Do Genetic Polymorphisms in Angiotensin Converting Enzyme 2 (ACE2) Gene Play a Role in Coronavirus Disease 2019 (COVID-19)? *Clin. Chem. Lab. Med.* **2020**, *58*, 1415–1422. [CrossRef]

72. Rabaan, A.A.; Al-Ahmed, S.H.; Albayat, H.; Alwarthan, S.; Alhajri, M.; Najim, M.A.; AlShehail, B.M.; Al-Adsani, W.; Alghadeer, A.; Abduljabbar, W.A.; et al. Variants of SARS-CoV-2: Influences on the Vaccines' Effectiveness and Possible Strategies to Overcome Their Consequences. *Medicina* **2023**, *59*, 507. [\[CrossRef\]](#)
73. Kuczyńska, K.; Zawilska, J.B.; Badura, J.; Strehl, B. SARS-CoV-2 virus: Origin, structure and replication cycle. *Farm. Pol.* **2021**, *77*, 143–149. [\[CrossRef\]](#)
74. Li, Q.; Wang, Y.; Sun, Q.; Knopf, J.; Herrmann, M.; Lin, L.; Jiang, J.; Shao, C.; Li, P.; He, X.; et al. Immune Response in COVID-19: What Is Next? *Cell Death Differ.* **2022**, *29*, 1107–1122. [\[CrossRef\]](#)
75. Mohammed, R.N.; Tamjidifar, R.; Rahman, H.S.; Adili, A.; Ghoreishizadeh, S.; Saeedi, H.; Thangavelu, L.; Shomali, N.; Aslam-inabad, R.; Marofi, F.; et al. A Comprehensive Review about Immune Responses and Exhaustion during Coronavirus Disease (COVID-19). *Cell Commun. Signal.* **2022**, *20*, 79. [\[CrossRef\]](#)
76. Vafaeinezhad, A.; Atashzar, M.R.; Baharlou, R. The Immune Responses against Coronavirus Infections: Friend or Foe? *Int. Arch. Allergy Immunol.* **2021**, *182*, 863–876. [\[CrossRef\]](#)
77. Schultze, J.L.; Aschenbrenner, A.C. COVID-19 and the Human Innate Immune System. *Cell* **2021**, *184*, 1671–1692. [\[CrossRef\]](#)
78. Xiong, Y.; Liu, Y.; Cao, L.; Wang, D.; Guo, M.; Jiang, A.; Guo, D.; Hu, W.; Yang, J.; Tang, Z.; et al. Transcriptomic Characteristics of Bronchoalveolar Lavage Fluid and Peripheral Blood Mononuclear Cells in COVID-19 Patients. *Emerg. Microbes Infect.* **2020**, *9*, 761–770. [\[CrossRef\]](#)
79. McKechnie, J.L.; Blish, C.A. The Innate Immune System: Fighting on the Front Lines or Fanning the Flames of COVID-19? *Cell Host Microbe* **2020**, *27*, 863–869. [\[CrossRef\]](#)
80. Ochani, R.K.; Asad, A.; Yasmin, F.; Shaikh, S.; Khalid, H.; Batra, S.; Sohail, M.R.; Mahmood, S.F.; Ochani, R.; Arshad, M.H.; et al. COVID-19 Pandemic: From Origins to Outcomes. A Comprehensive Review of Viral Pathogenesis, Clinical Manifestations, Diagnostic Evaluation, and Management. *Infez. Med.* **2021**, *29*, 20–36.
81. Ponti, G.; Maccaferri, M.; Ruini, C.; Tomasi, A.; Ozben, T. Biomarkers Associated with COVID-19 Disease Progression. *Crit. Rev. Clin. Lab. Sci.* **2020**, *57*, 389–399. [\[CrossRef\]](#)
82. Longo, R.; Peri, C.; Cricri, D.; Coppi, L.; Caruso, D.; Mitro, N.; De Fabiani, E.; Crestani, M. Ketogenic Diet: A New Light Shining on Old but Gold Biochemistry. *Nutrients* **2019**, *11*, 2497. [\[CrossRef\]](#)
83. Gershuni, V.M.; Yan, S.L.; Medici, V. Nutritional Ketosis for Weight Management and Reversal of Metabolic Syndrome. *Curr. Nutr. Rep.* **2018**, *7*, 97–106. [\[CrossRef\]](#)
84. Kolb, H.; Kempf, K.; Röhlings, M.; Lenzen-Schulte, M.; Schloot, N.C.; Martin, S. Ketone Bodies: From Enemy to Friend and Guardian Angel. *BMC Med.* **2021**, *19*, 313. [\[CrossRef\]](#)
85. Dhillon, K.K.; Gupta, S. Biochemistry, Ketogenesis. In *StatPearls*; StatPearls Publishing: Treasure Island, FL, USA, 2022.
86. Garcia, E.; Shalaurova, I.; Matyus, S.P.; Oskardmay, D.N.; Otvos, J.D.; Dullaart, R.P.F.; Connelly, M.A. Ketone Bodies Are Mildly Elevated in Subjects with Type 2 Diabetes Mellitus and Are Inversely Associated with Insulin Resistance as Measured by the Lipoprotein Insulin Resistance Index. *J. Clin. Med.* **2020**, *9*, 321. [\[CrossRef\]](#)
87. Hui, S.; Cowan, A.J.; Zeng, X.; Yang, L.; TeSlaa, T.; Li, X.; Bartman, C.; Zhang, Z.; Jang, C.; Wang, L.; et al. Quantitative Fluxomics of Circulating Metabolites. *Cell Metab.* **2020**, *32*, 676–688.e4. [\[CrossRef\]](#)
88. Wheless, J.W. History of the Ketogenic Diet. *Epilepsia* **2008**, *49*, 3–5. [\[CrossRef\]](#)
89. Peterman, M.G. The Ketogenic diet in epilepsy. *J. Am. Med. Assoc.* **1925**, *84*, 1979–1983. [\[CrossRef\]](#)
90. Sampaio, L.P. de B. Ketogenic Diet for Epilepsy Treatment. *Arq. Neuropsiquiatr.* **2016**, *74*, 842–848. [\[CrossRef\]](#)
91. Huttenlocher, P.R.; Wilbourn, A.J.; Signore, J.M. Medium-Chain Triglycerides as a Therapy for Intractable Childhood Epilepsy. *Neurology* **1971**, *21*, 1097–1103. [\[CrossRef\]](#)
92. Lee, J.E.; Titcomb, T.J.; Bisht, B.; Rubenstein, L.M.; Louison, R.; Wahls, T.L. A Modified MCT-Based Ketogenic Diet Increases Plasma  $\beta$ -Hydroxybutyrate but Has Less Effect on Fatigue and Quality of Life in People with Multiple Sclerosis Compared to a Modified Paleolithic Diet: A Waitlist-Controlled, Randomized Pilot Study. *J. Am. Coll. Nutr.* **2021**, *40*, 13–25. [\[CrossRef\]](#)
93. Kossoff, E.H.; Dorward, J.L. The Modified Atkins Diet. *Epilepsia* **2008**, *49*, 37–41. [\[CrossRef\]](#)
94. Cox, P.J.; Kirk, T.; Ashmore, T.; Willerton, K.; Evans, R.; Smith, A.; Murray, A.J.; Stubbs, B.; West, J.; McLure, S.W.; et al. Nutritional Ketosis Alters Fuel Preference and Thereby Endurance Performance in Athletes. *Cell Metab.* **2016**, *24*, 256–268. [\[CrossRef\]](#)
95. Kanikarla-Marie, P.; Jain, S.K. Hyperketonemia and Ketosis Increase the Risk of Complications in Type 1 Diabetes. *Free Radic. Biol. Med.* **2016**, *95*, 268–277. [\[CrossRef\]](#)
96. Zhu, H.; Bi, D.; Zhang, Y.; Kong, C.; Du, J.; Wu, X.; Wei, Q.; Qin, H. Ketogenic Diet for Human Diseases: The Underlying Mechanisms and Potential for Clinical Implementations. *Signal Transduct. Target. Ther.* **2022**, *7*, 11. [\[CrossRef\]](#) [\[PubMed\]](#)
97. Puchalska, P.; Crawford, P.A. Multi-Dimensional Roles of Ketone Bodies in Fuel Metabolism, Signaling, and Therapeutics. *Cell Metab.* **2017**, *25*, 262–284. [\[CrossRef\]](#)
98. Pfeifer, H.H.; Lyczkowski, D.A.; Thiele, E.A. Low Glycemic Index Treatment: Implementation and New Insights into Efficacy. *Epilepsia* **2008**, *49*, 42–45. [\[CrossRef\]](#)
99. Jenkins, D.J.A.; Wong, J.M.W.; Kendall, C.W.C.; Esfahani, A.; Ng, V.W.Y.; Leong, T.C.K.; Faulkner, D.A.; Vidgen, E.; Paul, G.; Mukherjee, R.; et al. Effect of a 6-Month Vegan Low-Carbohydrate ('Eco-Atkins') Diet on Cardiovascular Risk Factors and Body Weight in Hyperlipidaemic Adults: A Randomised Controlled Trial. *BMJ Open* **2014**, *4*, e003505. [\[CrossRef\]](#)

100. Ota, M.; Matsuo, J.; Ishida, I.; Takano, H.; Yokoi, Y.; Hori, H.; Yoshida, S.; Ashida, K.; Nakamura, K.; Takahashi, T.; et al. Effects of a Medium-Chain Triglyceride-Based Ketogenic Formula on Cognitive Function in Patients with Mild-to-Moderate Alzheimer's Disease. *Neurosci. Lett.* **2019**, *690*, 232–236. [CrossRef]
101. Grabacka, M.; Pierzchalska, M.; Dean, M.; Reiss, K. Regulation of Ketone Body Metabolism and the Role of PPAR $\alpha$ . *Int. J. Mol. Sci.* **2016**, *17*, 2093. [CrossRef]
102. Pondel, N.; Liškiewicz, D.; Liškiewicz, A. The ketogenic diet—Mechanism of action and prospects for therapeutic use: Data from clinical trials. *Postępy Biochem.* **2020**, *66*, 270–286. [CrossRef]
103. Talley, J.T.; Mohiuddin, S.S. *Biochemistry, Fatty Acid Oxidation*; StatPearls Publishing: Treasure Island, FL, USA, 2023.
104. Nałęcz, K.A.; Nałęcz, M.J. Carnitine—Mitochondria and beyond. *Postępy Biochem.* **2016**, *62*, 85–93.
105. Cabrera-Mulero, A.; Tinahones, A.; Bandera, B.; Moreno-Indias, I.; Macías-González, M.; Tinahones, F.J. Keto Microbiota: A Powerful Contributor to Host Disease Recovery. *Rev. Endocr. Metab. Disord.* **2019**, *20*, 415–425. [CrossRef]
106. Murtaza, N.; Burke, L.M.; Vlahovich, N.; Charlesson, B.; O' Neill, H.; Ross, M.L.; Campbell, K.L.; Krause, L.; Morrison, M. The Effects of Dietary Pattern during Intensified Training on Stool Microbiota of Elite Race Walkers. *Nutrients* **2019**, *11*, 261. [CrossRef]
107. Le, H.H.; Johnson, E.L. Going Keto? Say BHB-Ye Bye to Your Gut Bifidobacteria. *Cell Host Microbe* **2020**, *28*, 3–5. [CrossRef]
108. El Koofy, N.M.; Rady, H.I.; Abdallah, S.M.; Bazaraa, H.M.; Rabie, W.A.; El-Ayadi, A.A. The Effect of High Fat Dietary Modification and Nutritional Status on the Outcome of Critically Ill Ventilated Children: Single-Center Study. *Korean J. Pediatr.* **2019**, *62*, 344–352. [CrossRef]
109. al-Saady, N.M.; Blackmore, C.M.; Bennett, E.D. High Fat, Low Carbohydrate, Enteral Feeding Lowers PaCO<sub>2</sub> and Reduces the Period of Ventilation in Artificially Ventilated Patients. *Intensive Care Med.* **1989**, *15*, 290–295. [CrossRef]
110. Abd El Sabour Faramawy, M.; Abd Allah, A.; El Batrawy, S.; Amer, H. Impact of High Fat Low Carbohydrate Enteral Feeding on Weaning from Mechanical Ventilation. *Egypt. J. Chest Dis. Tuberc.* **2014**, *63*, 931–938. [CrossRef]
111. Li, J.; Wang, X.; Chen, J.; Zuo, X.; Zhang, H.; Deng, A. COVID-19 Infection May Cause Ketosis and Ketoacidosis. *Diabetes Obes. Metab.* **2020**, *22*, 1935–1941. [CrossRef]
112. Gera, S.; Longendyke, R.L.; Minich, N.M.; Malay, S.; Wood, J.R. The COVID-19 Pandemic and Associated Worsening of Diabetic Ketoacidosis Presentation in Youth. *Diabet. Med.* **2021**, *38*, e14610. [CrossRef]
113. Reddy, P.K.; Kuchay, M.S.; Mehta, Y.; Mishra, S.K. Diabetic Ketoacidosis Precipitated by COVID-19: A Report of Two Cases and Review of Literature. *Diabetes Metab. Syndr.* **2020**, *14*, 1459–1462. [CrossRef]
114. Diabetic Ketoacidosis in a Patient with COVID-19. Available online: <https://www.consultant360.com/article/consultant360/diabetic-ketoacidosis-patient-covid-19> (accessed on 11 September 2022).
115. Bruzzone, C.; Bizkarguenaga, M.; Gil-Redondo, R.; Diercks, T.; Arana, E.; García de Vicuña, A.; Seco, M.; Bosch, A.; Palazón, A.; San Juan, I.; et al. SARS-CoV-2 Infection Dysregulates the Metabolomic and Lipidomic Profiles of Serum. *iScience* **2020**, *23*, 101645. [CrossRef]
116. Karagiannis, F.; Peukert, K.; Surace, L.; Michla, M.; Nikolka, F.; Fox, M.; Weiss, P.; Feuerborn, C.; Maier, P.; Schulz, S.; et al. Impaired Ketogenesis Ties Metabolism to T Cell Dysfunction in COVID-19. *Nature* **2022**, *609*, 801–807. [CrossRef]
117. Al-Benna, S. Association of High Level Gene Expression of ACE2 in Adipose Tissue with Mortality of COVID-19 Infection in Obese Patients. *Obes. Med.* **2020**, *19*, 100283. [CrossRef] [PubMed]
118. D'Amico, F.; Baumgart, D.C.; Danese, S.; Peyrin-Biroulet, L. Diarrhea During COVID-19 Infection: Pathogenesis, Epidemiology, Prevention, and Management. *Clin. Gastroenterol. Hepatol.* **2020**, *18*, 1663–1672. [CrossRef] [PubMed]
119. Bogdański, A.; Słaba, A.; Kręgielska-Narozna, M. Obesity as a risk factor increasing the chance of a more severe course of COVID-19. *Forum Zaburzeń Metab.* **2020**, *11*, 105–111.
120. Iannelli, A.; Favre, G.; Frey, S.; Esnault, V.; Gugenheim, J.; Bouam, S.; Schiavo, L.; Tran, A.; Alifano, M. Obesity and COVID-19: ACE 2, the Missing Tile. *Obes. Surg.* **2020**, *30*, 4615–4617. [CrossRef] [PubMed]
121. Paravidino, V.B.; Leite, T.H.; Mediano, M.F.F.; Sichieri, R.; Azevedo e Silva, G.; Cravo, V.; Balduino, A.; Salgueiro, E.; Besen, B.A.M.P.; Moreira, R. de C.; et al. Association between Obesity and COVID-19 Mortality and Length of Stay in Intensive Care Unit Patients in Brazil: A Retrospective Cohort Study. *Sci. Rep.* **2022**, *12*, 13737. [CrossRef]
122. Kuehn, B.M. More Severe Obesity Leads to More Severe COVID-19 in Study. *JAMA* **2021**, *325*, 1603. [CrossRef]
123. Lighter, J.; Phillips, M.; Hochman, S.; Sterling, S.; Johnson, D.; Francois, F.; Stachel, A. Obesity in Patients Younger Than 60 Years Is a Risk Factor for COVID-19 Hospital Admission. *Clin. Infect. Dis.* **2020**, *71*, 896–897. [CrossRef]
124. Deng, M.; Qi, Y.; Deng, L.; Wang, H.; Xu, Y.; Li, Z.; Meng, Z.; Tang, J.; Dai, Z. Obesity as a Potential Predictor of Disease Severity in Young COVID-19 Patients: A Retrospective Study. *Obesity* **2020**, *28*, 1815–1825. [CrossRef]
125. Izquierdo, A.G.; Carreira, M.C.; Boughanem, H.; Moreno-Navarrete, J.M.; Nicoletti, C.F.; Oliver, P.; de Luis, D.; Nonino, C.B.; Portillo, M.P.; Martínez-Olmos, M.A.; et al. Adipose Tissue and Blood Leukocytes ACE2 DNA Methylation in Obesity and after Weight Loss. *Eur. J. Clin. Investig.* **2022**, *52*, e13685. [CrossRef]
126. Da Eira, D.; Jani, S.; Ceddia, R.B. Obesogenic and Ketogenic Diets Distinctly Regulate the SARS-CoV-2 Entry Proteins ACE2 and TMPRSS2 and the Renin-Angiotensin System in Rat Lung and Heart Tissues. *Nutrients* **2021**, *13*, 3357. [CrossRef]
127. Hoffmann, M.; Kleine-Weber, H.; Schroeder, S.; Krüger, N.; Herrler, T.; Erichsen, S.; Schiergens, T.S.; Herrler, G.; Wu, N.-H.; Nitsche, A.; et al. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* **2020**, *181*, 271–280.e8. [CrossRef] [PubMed]

128. Guo, D.F.; Sun, Y.L.; Hamet, P.; Inagami, T. The Angiotensin II Type 1 Receptor and Receptor-Associated Proteins. *Cell Res.* **2001**, *11*, 165–180. [[CrossRef](#)]
129. Méndez-García, L.A.; Escobedo, G.; Minguer-Urbe, A.G.; Viurcos-Sanabria, R.; Aguayo-Guerrero, J.A.; Carrillo-Ruiz, J.D.; Solleiro-Villavicencio, H. Role of the Renin-Angiotensin System in the Development of COVID-19-Associated Neurological Manifestations. *Front. Cell. Neurosci.* **2022**, *16*, 977039. [[CrossRef](#)]
130. Ames, M.K.; Atkins, C.E.; Pitt, B. The Renin-Angiotensin-Aldosterone System and Its Suppression. *J. Vet. Intern. Med.* **2019**, *33*, 363–382. [[CrossRef](#)]
131. Chaszczewska-Markowska, M.; Sagan, M.; Bogunia-Kubik, K. The renin-angiotensin-aldosterone system (RAAS)—Physiology and molecular mechanisms of functioning. *Postepy Hig. Med. Dosw.* **2016**, *70*, 917–927. [[CrossRef](#)] [[PubMed](#)]
132. Frankowska, S.; Zając, K.; Zargaryan, K.; Zawilska, J. Angiotensin converting enzyme 2 (ACE2)—The major receptor for SARS-CoV-2 virus. *Farm. Pol.* **2021**, *77*, 150–154. [[CrossRef](#)]
133. Grace, J.A.; Klein, S.; Herath, C.B.; Granzow, M.; Schierwagen, R.; Masing, N.; Walther, T.; Sauerbruch, T.; Burrell, L.M.; Angus, P.W.; et al. Activation of the MAS Receptor by Angiotensin-(1-7) in the Renin-Angiotensin System Mediates Mesenteric Vasodilatation in Cirrhosis. *Gastroenterology* **2013**, *145*, 874–884.e5. [[CrossRef](#)] [[PubMed](#)]
134. Fressatto de Godoy, M.A.; Pernomian, L.; de Oliveira, A.M.; Rattan, S. Biosynthetic Pathways and the Role of the Mas Receptor in the Effects of Angiotensin-(1-7) in Smooth Muscles. *Int. J. Hypertens.* **2012**, *2012*, 121740. [[CrossRef](#)]
135. Roshanravan, N.; Ghaffari, S.; Hedayati, M. Angiotensin Converting Enzyme-2 as Therapeutic Target in COVID-19. *Diabetes Metab. Syndr. Clin. Res. Rev.* **2020**, *14*, 637–639. [[CrossRef](#)]
136. Santos, R.A.S.; Sampaio, W.O.; Alzamora, A.C.; Motta-Santos, D.; Alenina, N.; Bader, M.; Campagnole-Santos, M.J. The ACE2/Angiotensin-(1-7)/MAS Axis of the Renin-Angiotensin System: Focus on Angiotensin-(1-7). *Physiol. Rev.* **2018**, *98*, 505–553. [[CrossRef](#)] [[PubMed](#)]
137. Behl, T.; Kaur, I.; Bungau, S.; Kumar, A.; Uddin, M.S.; Kumar, C.; Pal, G.; Sahil; Shrivastava, K.; Zengin, G.; et al. The Dual Impact of ACE2 in COVID-19 and Ironic Actions in Geriatrics and Pediatrics with Possible Therapeutic Solutions. *Life Sci.* **2020**, *257*, 118075. [[CrossRef](#)]
138. Ingraham, N.E.; Barakat, A.G.; Reilkoff, R.; Bezdicek, T.; Schacker, T.; Chipman, J.G.; Tignanelli, C.J.; Puskarich, M.A. Understanding the Renin-Angiotensin-Aldosterone-SARS-CoV-Axis: A Comprehensive Review. *Eur. Respir. J.* **2020**, *56*, 2000912. [[CrossRef](#)] [[PubMed](#)]
139. Deshotels, M.R.; Xia, H.; Sriramula, S.; Lazartigues, E.; Filipeanu, C.M. Angiotensin II Mediates Angiotensin Converting Enzyme Type 2 Internalization and Degradation through an Angiotensin II Type I Receptor-Dependent Mechanism. *Hypertension* **2014**, *64*, 1368–1375. [[CrossRef](#)] [[PubMed](#)]
140. Akoumianakis, I.; Filippatos, T. The Renin-Angiotensin-Aldosterone System as a Link between Obesity and Coronavirus Disease 2019 Severity. *Obes. Rev.* **2020**, *21*, e13077. [[CrossRef](#)] [[PubMed](#)]
141. Bean, D.M.; Kraljevic, Z.; Searle, T.; Bendayan, R.; Pickles, A.; Folarin, A.; Roguski, L.; Noor, K.; Shek, A.; O’Gallagher, K.; et al. Treatment with ACE-Inhibitors Is Associated with Less Severe Disease with SARS-COVID-19 Infection in a Multi-Site UK Acute Hospital Trust. *MedRxiv* **2020**. [[CrossRef](#)]
142. Meng, J.; Xiao, G.; Zhang, J.; He, X.; Ou, M.; Bi, J.; Yang, R.; Di, W.; Wang, Z.; Li, Z.; et al. Renin-Angiotensin System Inhibitors Improve the Clinical Outcomes of COVID-19 Patients with Hypertension. *Emerg. Microbes Infect.* **2020**, *9*, 757–760. [[CrossRef](#)]
143. Hirschberger, S.; Gellert, L.; Effinger, D.; Muenchhoff, M.; Herrmann, M.; Briegel, J.-M.; Zwißler, B.; Kreth, S. Ketone Bodies Improve Human CD8+ Cytotoxic T-Cell Immune Response During COVID-19 Infection. *Front. Med.* **2022**, *9*, 923502. [[CrossRef](#)]
144. Li, Z.; Zhang, Y.; Han, M.; Deng, H.; Wu, F.; Liu, G.; Chen, G.-Q. Lysine  $\beta$ -Hydroxybutyrylation Improves Stability of COVID-19 Antibody. *Biomacromolecules* **2022**, *23*, 454–463. [[CrossRef](#)]
145. Santos, A.F.; Póvoa, P.; Paixão, P.; Mendonça, A.; Taborda-Barata, L. Changes in Glycolytic Pathway in SARS-CoV-2 Infection and Their Importance in Understanding the Severity of COVID-19. *Front. Chem.* **2021**, *9*, 685196. [[CrossRef](#)]
146. Chen, W.; Gong, P.; Guo, J.; Li, H.; Li, R.; Xing, W.; Yang, Z.; Guan, Y. Glycolysis regulates pollen tube polarity via Rho GTPase signaling. *PLoS Genet.* **2018**, *14*, e1007373. [[CrossRef](#)]
147. Ostroukhova, M.; Goplen, N.; Karim, M.Z.; Michalec, L.; Guo, L.; Liang, Q.; Alam, R. The Role of Low-Level Lactate Production in Airway Inflammation in Asthma. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2012**, *302*, L300–L307. [[CrossRef](#)]
148. Leavy, O. Defective Mitochondria Disrupt CD8+ T Cells. *Nat. Rev. Immunol.* **2016**, *16*, 535. [[CrossRef](#)] [[PubMed](#)]
149. Atila, A.; Alay, H.; Yaman, M.E.; Akman, T.C.; Cadirci, E.; Bayrak, B.; Celik, S.; Atila, N.E.; Yaganoglu, A.M.; Kadioglu, Y.; et al. The Serum Amino Acid Profile in COVID-19. *Amino Acids* **2021**, *53*, 1569–1588. [[CrossRef](#)] [[PubMed](#)]
150. Masoodi, M.; Peschka, M.; Schmiedel, S.; Haddad, M.; Frye, M.; Maas, C.; Lohse, A.; Huber, S.; Kirchhof, P.; Nofer, J.-R.; et al. Disturbed Lipid and Amino Acid Metabolisms in COVID-19 Patients. *J. Mol. Med.* **2022**, *100*, 555–568. [[CrossRef](#)] [[PubMed](#)]
151. Januarie, K.C.; Uhuo, O.V.; Iwuoha, E.; Feleni, U. Recent Advances in the Detection of Interferon-Gamma as a TB Biomarker. *Anal. Bioanal. Chem.* **2022**, *414*, 907–921. [[CrossRef](#)]
152. Altay, O.; Arif, M.; Li, X.; Yang, H.; Aydın, M.; Alkurt, G.; Kim, W.; Akyol, D.; Zhang, C.; Dinler-Doganay, G.; et al. Combined Metabolic Activators Accelerates Recovery in Mild-to-Moderate COVID-19. *Adv. Sci.* **2021**, *8*, e2101222. [[CrossRef](#)]
153. Sumbalova, Z.; Kucharska, J.; Palacka, P.; Rausova, Z.; Langsjoen, P.H.; Langsjoen, A.M.; Gvozdzakova, A. Platelet Mitochondrial Function and Endogenous Coenzyme Q10 Levels Are Reduced in Patients after COVID-19. *Bratisl. Lek. Listy* **2022**, *123*, 9–15. [[CrossRef](#)]

154. Guarnieri, J.W.; Dybas, J.M.; Fazelinia, H.; Kim, M.S.; Frere, J.; Zhang, Y.; Albrecht, Y.S.; Murdock, D.G.; Angelin, A.; Singh, L.N.; et al. Targeted down regulation of core mitochondrial genes during SARS-CoV-2 infection. *bioRxiv* **2022**. [[CrossRef](#)]
155. Pearce, E.L.; Poffenberger, M.C.; Chang, C.-H.; Jones, R.G. Fueling Immunity: Insights into Metabolism and Lymphocyte Function. *Science* **2013**, *342*, 1242454. [[CrossRef](#)]
156. Hirschberger, S.; Strauß, G.; Effinger, D.; Marstaller, X.; Ferstl, A.; Müller, M.B.; Wu, T.; Hübner, M.; Rahmel, T.; Mascolo, H.; et al. Very-low-carbohydrate Diet Enhances Human T-cell Immunity through Immunometabolic Reprogramming. *EMBO Mol. Med.* **2021**, *13*, e14323. [[CrossRef](#)]
157. Zhang, H.; Tang, K.; Ma, J.; Zhou, L.; Liu, J.; Zeng, L.; Zhu, L.; Xu, P.; Chen, J.; Wei, K.; et al. Ketogenesis-Generated  $\beta$ -Hydroxybutyrate Is an Epigenetic Regulator of CD8+ T-Cell Memory Development. *Nat. Cell Biol.* **2020**, *22*, 18–25. [[CrossRef](#)] [[PubMed](#)]
158. Cotter, D.G.; Schugar, R.C.; Crawford, P.A. Ketone Body Metabolism and Cardiovascular Disease. *Am. J. Physiol. Heart Circ. Physiol.* **2013**, *304*, H1060–H1076. [[CrossRef](#)] [[PubMed](#)]
159. Lu, Y.-J.; Barreira-Silva, P.; Boyce, S.; Powers, J.; Cavallo, K.; Behar, S.M. CD4 T Cell Help Prevents CD8 T Cell Exhaustion and Promotes Control of Mycobacterium Tuberculosis Infection. *Cell. Rep.* **2021**, *36*, 109696. [[CrossRef](#)]
160. Wang, Y.; Zheng, J.; Islam, M.S.; Yang, Y.; Hu, Y.; Chen, X. The Role of CD4+FoxP3+ Regulatory T Cells in the Immunopathogenesis of COVID-19: Implications for Treatment. *Int. J. Biol. Sci.* **2021**, *17*, 1507–1520. [[CrossRef](#)] [[PubMed](#)]
161. Wik, J.A.; Skålhegg, B.S. T Cell Metabolism in Infection. *Front. Immunol.* **2022**, *13*, 840610. [[CrossRef](#)]
162. Kang, C.K.; Han, G.-C.; Kim, M.; Kim, G.; Shin, H.M.; Song, K.-H.; Choe, P.G.; Park, W.B.; Kim, E.S.; Kim, H.B.; et al. Aberrant Hyperactivation of Cytotoxic T-Cell as a Potential Determinant of COVID-19 Severity. *Int. J. Infect. Dis.* **2020**, *97*, 313–321. [[CrossRef](#)]
163. Paletta, A.; Di Diego García, F.; Varese, A.; Erra Diaz, F.; García, J.; Cisneros, J.C.; Ludueña, G.; Mazzitelli, I.; Pisarevsky, A.; Cabrerizo, G.; et al. Platelets Modulate CD4+ T-Cell Function in COVID-19 through a PD-L1 Dependent Mechanism. *Br. J. Haematol.* **2022**, *197*, 283–292. [[CrossRef](#)]
164. Russo, C.; Raiden, S.; Algeri, S.; De Carli, N.; Davenport, C.; Sarli, M.; Bruera, M.J.; Seery, V.; Sananez, I.; Simaz, N.; et al. Extracellular ATP and Imbalance of CD4+ T Cell Compartment in Pediatric COVID-19. *Front. Cell. Infect. Microbiol.* **2022**, *12*, 893044. [[CrossRef](#)]
165. Salomé, B.; Horowitz, A. Impaired CD4 T-cell Response to SARS-CoV-2: Rationale for PD-1 Blockade in Patients with Cancer and COVID-19? *Cancer Discov.* **2021**, *11*, 1877–1878. [[CrossRef](#)]
166. Davanzo, G.G.; Codo, A.C.; Brunetti, N.S.; Boldrini, V.; Knittel, T.L.; Monterio, L.B.; de Moraes, D.; Ferrari, A.J.R.; de Souza, G.F.; Muraro, S.P.; et al. SARS-CoV-2 Uses CD4 to Infect T Helper Lymphocytes. *MedRxiv* **2020**. [[CrossRef](#)]
167. Kennedy, A.R.; Pissios, P.; Otu, H.; Roberson, R.; Xue, B.; Asakura, K.; Furukawa, N.; Marino, F.E.; Liu, F.-F.; Kahn, B.B.; et al. A High-Fat, Ketogenic Diet Induces a Unique Metabolic State in Mice. *Am. J. Physiol. Endocrinol. Metab.* **2007**, *292*, E1724–E1739. [[CrossRef](#)] [[PubMed](#)]
168. Kratochvil, R.M.; Kubes, P.; Deniset, J.F. Monocyte Conversion during Inflammation and Injury. *Arterioscler. Thromb. Vasc. Biol.* **2017**, *37*, 35–42. [[CrossRef](#)] [[PubMed](#)]
169. Junqueira, C.; Crespo, Á.; Ranjbar, S.; de Lacerda, L.B.; Lewandrowski, M.; Ingber, J.; Parry, B.; Ravid, S.; Clark, S.; Schrimpf, M.R.; et al. Fc $\gamma$ R-Mediated SARS-CoV-2 Infection of Monocytes Activates Inflammation. *Nature* **2022**, *606*, 576–584. [[CrossRef](#)]
170. Stephenson, E.; Reynolds, G.; Botting, R.A.; Calero-Nieto, F.J.; Morgan, M.D.; Tuong, Z.K.; Bach, K.; Sungnak, W.; Worlock, K.B.; Yoshida, M.; et al. Single-Cell Multi-Omics Analysis of the Immune Response in COVID-19. *Nat. Med.* **2021**, *27*, 904–916. [[CrossRef](#)] [[PubMed](#)]
171. Wang, S.; Song, R.; Wang, Z.; Jing, Z.; Wang, S.; Ma, J. S100A8/A9 in Inflammation. *Front. Immunol.* **2018**, *9*, 1298. [[CrossRef](#)]
172. Yang, D.; Han, Z.; Oppenheim, J.J. Alarmins and Immunity. *Immunol. Rev.* **2017**, *280*, 41–56. [[CrossRef](#)]
173. Szmyt, A.; Kasprzak, A. S100 family proteins in the pathogenesis of gallbladder stones. *Now. Lek.* **2011**, *80*, 213–218.
174. Chan, J.K.; Roth, J.; Oppenheim, J.J.; Tracey, K.J.; Vogl, T.; Feldmann, M.; Horwood, N.; Nanchahal, J. Alarmins: Awaiting a Clinical Response. *J. Clin. Invest.* **2012**, *122*, 2711–2719. [[CrossRef](#)]
175. Cher, J.Z.B.; Akbar, M.; Kitson, S.; Crowe, L.A.N.; Garcia-Melchor, E.; Hannah, S.C.; McLean, M.; Fazzi, U.G.; Kerr, S.C.; Murrell, G.A.C.; et al. Alarmins in Frozen Shoulder: A Molecular Association between Inflammation and Pain. *Am. J. Sport Med.* **2018**, *46*, 671–678. [[CrossRef](#)]
176. Qin, C.; Zhou, L.; Hu, Z.; Zhang, S.; Yang, S.; Tao, Y.; Xie, C.; Ma, K.; Shang, K.; Wang, W.; et al. Dysregulation of Immune Response in Patients with Coronavirus 2019 (COVID-19) in Wuhan, China. *Clin. Infect. Dis.* **2020**, *71*, 762–768. [[CrossRef](#)]
177. Yang, X.; Yu, Y.; Xu, J.; Shu, H.; Xia, J.; Liu, H.; Wu, Y.; Zhang, L.; Yu, Z.; Fang, M.; et al. Clinical Course and Outcomes of Critically Ill Patients with SARS-CoV-2 Pneumonia in Wuhan, China: A Single-Centered, Retrospective, Observational Study. *Lancet Respir. Med.* **2020**, *8*, 475–481. [[CrossRef](#)] [[PubMed](#)]
178. Sohn, K.M.; Lee, S.-G.; Kim, H.J.; Cheon, S.; Jeong, H.; Lee, J.; Kim, I.S.; Silwal, P.; Kim, Y.J.; Paik, S.; et al. COVID-19 Patients Upregulate Toll-like Receptor 4-Mediated Inflammatory Signaling That Mimics Bacterial Sepsis. *J. Korean Med. Sci.* **2020**, *35*, e343. [[CrossRef](#)] [[PubMed](#)]

179. Camajani, E.; Feraco, A.; Basciani, S.; Gnassi, L.; Barrea, L.; Armani, A.; Caprio, M. VLCKD in Combination with Physical Exercise Preserves Skeletal Muscle Mass in Sarcopenic Obesity after Severe COVID-19 Disease: A Case Report. *Healthcare* **2022**, *10*, 573. [[CrossRef](#)]
180. Volk, B.M.; Roberts, C.G.P.; VanTieghem, M.; George, M.P.; Adams, R.N.; Athinarayanan, S.J.; McKenzie, A.L. Reduced COVID-19 Severity Elicited by Weight Loss from a Medically Supervised Ketogenic Diet in a Geographically Diverse Ambulatory Population with Type 2 Diabetes and Obesity. *BMJ Nutr. Prev. Health* **2022**, *5*, e000444. [[CrossRef](#)] [[PubMed](#)]
181. Thevarajan, I.; Nguyen, T.H.O.; Koutsakos, M.; Druce, J.; Caly, L.; van de Sandt, C.E.; Jia, X.; Nicholson, S.; Catton, M.; Cowie, B.; et al. Breadth of Concomitant Immune Responses Prior to Patient Recovery: A Case Report of Non-Severe COVID-19. *Nat. Med.* **2020**, *26*, 453–455. [[CrossRef](#)]
182. Cerapio, J.P.; Perrier, M.; Pont, F.; Tosolini, M.; Laurent, C.; Bertani, S.; Fournie, J.-J. Single-Cell RNAseq Profiling of Human  $\Gamma\delta$  T Lymphocytes in Virus-Related Cancers and COVID-19 Disease. *Viruses* **2021**, *13*, 2212. [[CrossRef](#)] [[PubMed](#)]
183. Hayday, A.C.; Vantourout, P. The Innate Biologies of Adaptive Antigen Receptors. *Annu. Rev. Immunol.* **2020**, *38*, 487–510. [[CrossRef](#)]
184. Carissimo, G.; Xu, W.; Kwok, I.; Abdad, M.Y.; Chan, Y.-H.; Fong, S.-W.; Puan, K.J.; Lee, C.Y.-P.; Yeo, N.K.-W.; Amrun, S.N.; et al. Whole Blood Immunophenotyping Uncovers Immature Neutrophil-to-VD2 T-Cell Ratio as an Early Marker for Severe COVID-19. *Nat. Commun.* **2020**, *11*, 5243. [[CrossRef](#)]
185. Lei, L.; Qian, H.; Yang, X.; Zhang, X.; Zhang, D.; Dai, T.; Guo, R.; Shi, L.; Cheng, Y.; Zhang, B.; et al. The phenotypic changes of  $\gamma\delta$  T cells in COVID-19 patients. *J. Cell. Mol. Med.* **2020**, *24*, 11603–11606. [[CrossRef](#)]
186. Rijkers, G.; Vervenne, T.; van der Pol, P. More Bricks in the Wall against SARS-CoV-2 Infection: Involvement of  $\Gamma\delta$  T Cells. *Cell. Mol. Immunol.* **2020**, *17*, 771–772. [[CrossRef](#)]
187. Guo, X.-Z.J.; Dash, P.; Crawford, J.C.; Allen, E.K.; Zamora, A.E.; Boyd, D.F.; Duan, S.; Bajracharya, R.; Awad, W.A.; Apiwattanakul, N.; et al. Lung  $\Gamma\delta$  T Cells Mediate Protective Responses during Neonatal Influenza Infection That Are Associated with Type 2 Immunity. *Immunity* **2018**, *49*, 531–544.e6. [[CrossRef](#)] [[PubMed](#)]
188. Sabbaghi, A.; Miri, S.M.; Keshavarz, M.; Mahooti, M.; Zebardast, A.; Ghaemi, A. Role of  $\Gamma\delta$  T Cells in Controlling Viral Infections with a Focus on Influenza Virus: Implications for Designing Novel Therapeutic Approaches. *Virol. J.* **2020**, *17*, 174. [[CrossRef](#)] [[PubMed](#)]
189. Kumar, A.; Singh, B.; Tiwari, R.; Singh, V.K.; Singh, S.S.; Sundar, S.; Kumar, R. Emerging Role of  $\Gamma\delta$  T Cells in Protozoan Infection and Their Potential Clinical Application. *Infect. Genet. Evol.* **2022**, *98*, 105210. [[CrossRef](#)]
190. Ratajczak, M.Z.; Bujko, K.; Ciechanowicz, A.; Sielatycka, K.; Cymer, M.; Marlicz, W.; Kucia, M. SARS-CoV-2 Entry Receptor ACE2 Is Expressed on Very Small CD45- Precursors of Hematopoietic and Endothelial Cells and in Response to Virus Spike Protein Activates the Nlrp3 Inflammasome. *Stem Cell Rev. Rep.* **2021**, *17*, 266–277. [[CrossRef](#)] [[PubMed](#)]
191. Rodrigues, T.S.; de Sá, K.S.G.; Ishimoto, A.Y.; Becerra, A.; Oliveira, S.; Almeida, L.; Gonçalves, A.V.; Perucello, D.B.; Andrade, W.A.; Castro, R.; et al. Inflammasomes Are Activated in Response to SARS-CoV-2 Infection and Are Associated with COVID-19 Severity in Patients. *J. Exp. Med.* **2021**, *218*, e20201707. [[CrossRef](#)]
192. Amin, S.; Aktar, S.; Rahman, M.M.; Chowdhury, M.M.H. NLRP3 Inflammasome Activation in COVID-19: An Interlink between Risk Factors and Disease Severity. *Microbes Infect.* **2022**, *24*, 104913. [[CrossRef](#)] [[PubMed](#)]
193. Freeman, T.L.; Swartz, T.H. Targeting the NLRP3 Inflammasome in Severe COVID-19. *Front. Immunol.* **2020**, *11*, 1518. [[CrossRef](#)]
194. Zhao, N.; Di, B.; Xu, L.-L. The NLRP3 Inflammasome and COVID-19: Activation, Pathogenesis and Therapeutic Strategies. *Cytokine Growth Factor Rev.* **2021**, *61*, 2–15. [[CrossRef](#)]
195. Dutta, D.; Liu, J.; Xiong, H. NLRP3 Inflammasome Activation and SARS-CoV-2-Mediated Hyperinflammation, Cytokine Storm and Neurological Syndromes. *Int. J. Physiol. Pathophysiol. Pharmacol.* **2022**, *14*, 138–160.
196. Xian, H.; Liu, Y.; Rundberg Nilsson, A.; Gatchalian, R.; Crother, T.R.; Tourtellotte, W.G.; Zhang, Y.; Aleman-Muench, G.R.; Lewis, G.; Chen, W.; et al. Metformin Inhibition of Mitochondrial ATP and DNA Synthesis Abrogates NLRP3 Inflammasome Activation and Pulmonary Inflammation. *Immunity* **2021**, *54*, 1463–1477.e11. [[CrossRef](#)]
197. Grailer, J.J.; Canning, B.A.; Kalbitz, M.; Haggadone, M.D.; Dhond, R.M.; Andjelkovic, A.V.; Zetoune, F.S.; Ward, P.A. Critical Role for the NLRP3 Inflammasome during Acute Lung Injury. *J. Immunol.* **2014**, *192*, 5974–5983. [[CrossRef](#)]
198. Kow, C.S.; Lee, L.-H.; Ramachandram, D.S.; Hasan, S.S.; Ming, L.C.; Goh, H.P. The Effect of Colchicine on Mortality Outcome and Duration of Hospital Stay in Patients with COVID-19: A Meta-Analysis of Randomized Trials. *Immun. Inflamm. Dis.* **2022**, *10*, 255–264. [[CrossRef](#)]
199. Ryu, S.; Shchukina, I.; Youm, Y.H.; Qing, H.; Hilliard, B.; Dlugos, T.; Zhang, X.; Yasumoto, Y.; Booth, C.J.; Fernández-Hernando, C.; et al. Ketogenic diet restrains aging-induced exacerbation of coronavirus infection in mice. *Elife* **2021**, *10*, e66522. [[CrossRef](#)] [[PubMed](#)]
200. Saeedi-Boroujeni, A.; Nashibi, R.; Ghadiri, A.A.; Nakajima, M.; Salmanzadeh, S.; Mahmoudian-Sani, M.-R.; Hanafi, M.G.; Sharhani, A.; Khodadadi, A. Tranilast as an Adjunctive Therapy in Hospitalized Patients with Severe COVID-19: A Randomized Controlled Trial. *Arch. Med. Res.* **2022**, *53*, 368–377. [[CrossRef](#)] [[PubMed](#)]
201. Rha, M.-S.; Shin, E.-C. Activation or Exhaustion of CD8+ T Cells in Patients with COVID-19. *Cell. Mol. Immunol.* **2021**, *18*, 2325–2333. [[CrossRef](#)] [[PubMed](#)]

202. Azkur, A.K.; Akdis, M.; Azkur, D.; Sokolowska, M.; van de Veen, W.; Brügggen, M.; O'Mahony, L.; Gao, Y.; Nadeau, K.; Akdis, C.A. Immune Response to SARS-CoV-2 and Mechanisms of Immunopathological Changes in COVID-19. *Allergy* **2020**, *75*, 1564–1581. [[CrossRef](#)]
203. Mortezaee, K.; Majidpoor, J. CD8+ T Cells in SARS-CoV-2 Induced Disease and Cancer—Clinical Perspectives. *Front. Immunol.* **2022**, *13*, 864298. [[CrossRef](#)]
204. De Biasi, S.; Meschiari, M.; Gibellini, L.; Bellinazzi, C.; Borella, R.; Fidanza, L.; Gozzi, L.; Iannone, A.; Lo Tartaro, D.; Mattioli, M.; et al. Marked T Cell Activation, Senescence, Exhaustion and Skewing towards TH17 in Patients with COVID-19 Pneumonia. *Nat. Commun.* **2020**, *11*, 3434. [[CrossRef](#)]
205. Zheng, H.-Y.; Zhang, M.; Yang, C.-X.; Zhang, N.; Wang, X.-C.; Yang, X.-P.; Dong, X.-Q.; Zheng, Y.-T. Elevated Exhaustion Levels and Reduced Functional Diversity of T Cells in Peripheral Blood May Predict Severe Progression in COVID-19 Patients. *Cell. Mol. Immunol.* **2020**, *17*, 541–543. [[CrossRef](#)]
206. Zheng, M.; Gao, Y.; Wang, G.; Song, G.; Liu, S.; Sun, D.; Xu, Y.; Tian, Z. Functional exhaustion of antiviral lymphocytes in COVID-19 patients. *Cell. Mol. Immunol.* **2020**, *17*, 533–535. [[CrossRef](#)]
207. Diao, B.; Wang, C.; Tan, Y.; Chen, X.; Liu, Y.; Ning, L.; Chen, L.; Li, M.; Liu, Y.; Wang, G.; et al. Reduction and Functional Exhaustion of T Cells in Patients With Coronavirus Disease 2019 (COVID-19). *Front. Immunol.* **2020**, *11*, 827. [[CrossRef](#)]
208. Ganji, A.; Farahani, I.; Khansarnejad, B.; Ghazavi, A.; Mosayebi, G. Increased Expression of CD8 Marker on T-Cells in COVID-19 Patients. *Blood Cells Mol. Dis.* **2020**, *83*, 102437. [[CrossRef](#)] [[PubMed](#)]
209. Charles, A.; Janeway, J.; Travers, P.; Walport, M.; Shlomchik, M.J. T cell-mediated cytotoxicity. In *Immunobiology: The Immune System in Health and Disease*, 5th ed.; Garland Publishing: New York, NY, USA, 2001.
210. Mescher, M.F.; Curtsinger, J.M.; Agarwal, P.; Casey, K.A.; Gerner, M.; Hammerbeck, C.D.; Popescu, F.; Xiao, Z. Signals Required for Programming Effector and Memory Development by CD8+ T Cells. *Immunol. Rev.* **2006**, *211*, 81–92. [[CrossRef](#)] [[PubMed](#)]
211. Osińska, I.; Popko, K.; Demkow, U. Perforin: An Important Player in Immune Response. *Cent. Eur. J. Immunol.* **2014**, *39*, 109–115. [[CrossRef](#)]
212. Cunningham, L.; Kimber, I.; Basketter, D.; Simmonds, P.; McSweeney, S.; Tziotzios, C.; McFadden, J.P. Perforin, COVID-19 and a Possible Pathogenic Auto-inflammatory Feedback Loop. *Scand. J. Immunol.* **2021**, *94*, e13102. [[CrossRef](#)] [[PubMed](#)]
213. Paduch, R.; Klatka, M.; Klatka, J. Types of cell death. *Pom. J. Life Sci.* **2015**, *61*, 411–418.
214. Gupta, M.; Greer, P.; Mahanty, S.; Shieh, W.-J.; Zaki, S.R.; Ahmed, R.; Rollin, P.E. CD8-Mediated Protection against Ebola Virus Infection Is Perforin Dependent. *J. Immunol.* **2005**, *174*, 4198–4202. [[CrossRef](#)] [[PubMed](#)]
215. Xie, Z.; Zhang, D.; Chung, D.; Tang, Z.; Huang, H.; Dai, L.; Qi, S.; Li, J.; Colak, G.; Chen, Y.; et al. Metabolic Regulation of Gene Expression by Histone Lysine  $\beta$ -Hydroxybutyrylation. *Mol. Cell* **2016**, *62*, 194–206. [[CrossRef](#)] [[PubMed](#)]
216. Liemburg-Apers, D.C.; Willems, P.H.G.M.; Koopman, W.J.H.; Grefte, S. Interactions between Mitochondrial Reactive Oxygen Species and Cellular Glucose Metabolism. *Arch. Toxicol.* **2015**, *89*, 1209–1226. [[CrossRef](#)]
217. Shenoy, S. Coronavirus (COVID-19) Sepsis: Revisiting Mitochondrial Dysfunction in Pathogenesis, Aging, Inflammation, and Mortality. *Inflamm. Res.* **2020**, *69*, 1077–1085. [[CrossRef](#)]
218. Zhang, B.; Pan, C.; Feng, C.; Yan, C.; Yu, Y.; Chen, Z.; Guo, C.; Wang, X. Role of Mitochondrial Reactive Oxygen Species in Homeostasis Regulation. *Redox Rep.* **2022**, *27*, 45–52. [[CrossRef](#)]
219. Bassoy, E.Y.; Walch, M.; Martinvalet, D. Reactive Oxygen Species: Do They Play a Role in Adaptive Immunity? *Front. Immunol.* **2021**, *12*, 4826. [[CrossRef](#)] [[PubMed](#)]
220. Veenith, T.; Martin, H.; Le Breuille, M.; Whitehouse, T.; Gao-Smith, F.; Duggal, N.; Lord, J.M.; Mian, R.; Sarphe, D.; Moss, P. High Generation of Reactive Oxygen Species from Neutrophils in Patients with Severe COVID-19. *Sci. Rep.* **2022**, *12*, 10484. [[CrossRef](#)] [[PubMed](#)]
221. Angelidi, A.M.; Kokkinos, A.; Katechaki, E.; Ros, E.; Mantzoros, C.S. Mediterranean Diet as a Nutritional Approach for COVID-19. *Metabolism* **2021**, *114*, 154407. [[CrossRef](#)]
222. Berbudi, A.; Rahmadika, N.; Tjahjadi, A.I.; Ruslami, R. Type 2 Diabetes and Its Impact on the Immune System. *Curr. Diabetes Rev.* **2020**, *16*, 442–449. [[CrossRef](#)] [[PubMed](#)]
223. Schiaffino, S.; Albano, D.; Cozzi, A.; Messina, C.; Arioli, R.; Brà, C.; Bruno, A.; Carbonaro, L.A.; Carriero, A.; Carriero, S.; et al. CT-Derived Chest Muscle Metrics for Outcome Prediction in Patients with COVID-19. *Radiology* **2021**, *300*, E328–E336. [[CrossRef](#)] [[PubMed](#)]
224. Khunti, K.; Del Prato, S.; Mathieu, C.; Kahn, S.E.; Gabbay, R.A.; Buse, J.B. COVID-19, Hyperglycemia, and New-Onset Diabetes. *Diabetes Care* **2021**, *44*, 2645–2655. [[CrossRef](#)]
225. CDC COVID-19 Response Team. Preliminary Estimates of the Prevalence of Selected Underlying Health Conditions among Patients with Coronavirus Disease 2019—United States, 12 February–28 March 2020. *Morb. Mortal. Wkly. Rep.* **2020**, *69*, 382–386. [[CrossRef](#)]
226. Mahat, R.K.; Rathore, V.; Singh, N.; Singh, N.; Singh, S.K.; Shah, R.K.; Garg, C. Lipid Profile as an Indicator of COVID-19 Severity: A Systematic Review and Meta-Analysis. *Clin. Nutr. ESPEN* **2021**, *45*, 91–101. [[CrossRef](#)]
227. Chidambaram, V.; Kumar, A.; Majella, M.G.; Seth, B.; Sivakumar, R.K.; Voruganti, D.; Bavineni, M.; Baghal, A.; Gates, K.; Kumari, A.; et al. HDL Cholesterol Levels and Susceptibility to COVID-19. *eBioMedicine* **2022**, *82*, 104166. [[CrossRef](#)]

228. Bilge, M.; Kibar Akilli, I. Low-Density Lipoprotein Cholesterol Levels and Disease Severity in COVID-19 Pneumonia. *Med. J. Bakirkoy* **2021**, *17*, 274–279. [[CrossRef](#)]
229. Cooper, I.D.; Crofts, C.A.P.; DiNicolantonio, J.J.; Malhotra, A.; Elliott, B.; Kyriakidou, Y.; Brookler, K.H. Relationships between Hyperinsulinaemia, Magnesium, Vitamin D, Thrombosis and COVID-19: Rationale for Clinical Management. *Open Heart* **2020**, *7*, e001356. [[CrossRef](#)] [[PubMed](#)]

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.