

Hormonal and dietary factors in acne vulgaris versus controls

Thomas Jonathan Stewart, BBioMedSc MBBS FRACGP^{a,c} and Carl Bazergy, MBBS FRACGP^b

^aDarlinghurst Medical Centre, Darlinghurst, Sydney, Australia; ^bKogarah Railway Medical Centre, Kogarah, Sydney, Australia; ^cSchool of Medicine, University of New South Wales, Sydney, Australia

ABSTRACT

Background: Acne vulgaris is an inflammatory skin disorder with not as yet fully understood pathogenesis. In this controlled study, we assessed acne vulgaris patients for several possible pathogenic factors such as vitamin D deficiency, vegan diet, increased body mass index (BMI) and positive anti-transglutaminase antibody. **Methods:** We screened 10 years of records at a family medicine clinic for patients diagnosed with acne vulgaris. In eligible subjects, we collected data regarding 25-hydroxyvitamin D levels, BMI, dietary preference and serum IgA tissue transglutaminase levels. Controls were age- (+/- 12 months) and sex-matched patients seen during the study period without a diagnosis of acne vulgaris. **Results:** 453 patients were given a diagnosis of acne vulgaris during the study period. Compared with controls, we found significant associations between vitamin D deficiency (<50nmol/L), and/or positive transglutaminase antibody level (>4.0U/mL) and a diagnosis of acne vulgaris. **Conclusions:** Our study adds important information to the current body of literature in pursuit of elucidating the pathogenesis of this complex multifactorial disease.

ARTICLE HISTORY

Received 14 December 2017
Accepted 13 February 2018

KEYWORDS

Acne vulgaris; vitamin D deficiency; anti-transglutaminase antibody; obesity; diet

Introduction

Acne vulgaris is a chronic inflammatory disorder of the pilosebaceous unit typically affecting areas with a high density of hormonally-responsive sebaceous glands such as the face, neck, chest, upper back, and upper arms.

Pathogenesis is complex and remains incompletely understood. Follicular hyperkeratinization, sebum production, *Propionibacterium acnes* and inflammation appear to be involved.¹ The role of factors such as vitamin D, diet, obesity and autoinflammation is evolving.

In vitro, vitamin D has been shown to regulate proliferation and differentiation of keratinocytes and sebocytes, and may also have anti-comedogenic properties. Associations between acne vulgaris and vitamin D deficiency have been demonstrated in a few small cohorts.²⁻⁴

Studies investigating the influence of diet on acne pathogenesis have been unconvincing and focused mainly on specific foods (e.g. milk),⁵⁻⁷ which may be artificial in place of diet category. The relationship

between body mass index (BMI) and acne has received limited attention and remains uncertain.

Anti-tissue transglutaminase antibody (anti-TTG) is a serum autoantibody utilised primarily in the diagnosis of coeliac disease. Anti-TTG positivity has been associated with vitamin D deficiency in coeliac disease, and has also been seen in several other autoinflammatory conditions.⁸⁻¹⁰

The aim of this study was to determine if there are any associations between serum vitamin D levels, vegan diet, increased BMI and/or anti-TTG, and acne vulgaris in a large Australian population.

Results

453 patients (205 males and 248 females) were enrolled in the study and their mean age was 22.8 years. The control group comprised 101 males and 149 females with a mean age of 23.1 years. Serum vitamin D levels, diet, body mass index, and positive tissue transglutaminase antibodies for both groups are detailed in Table 1.

CONTACT Thomas Jonathan Stewart  thomas_stewart@live.com  506/22 Danks street, Waterloo 2017, Sydney, Australia.

© 2018 Thomas Jonathan Stewart and Carl Bazergy. Published with license by Taylor & Francis

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

Table 1. Hormonal and dietary factors in acne patients vs controls.

	Patients (n = 453)	Controls (n = 250)	Chi ² (1df)	P value
Vitamin D deficiency	192 (42.4%)	61 (24.4%)	22.62	<i>P</i> < 0.0005*
Vegan diet	8 (1.7%)	9 (3.6%)	2.3	<i>P</i> = 0.1297
Increased BMI	128 (28.2%)	67 (26.8%)	0.17	<i>P</i> = 0.6798
Positive tissue transglutaminase antibody	32 (7.0%)	5 (2.0%)	8.29	<i>P</i> = 0.0040*

Coexistence of vitamin D deficiency and positive tissue transglutaminase in patients versus controls was statistically significant (chi² (1) = 5.10, Fisher's exact = 0.042). No other significant associations existed between any of the measured factors.

Results (Table 1) showed a statistically significant difference between acne and control groups for vitamin D deficiency (chi² (1) = 22.62, *p* < 0.0005) or positive tissue transglutaminase antibody (chi² (1) = 8.29, *p* = 0.004). The difference was not significant for vegan diet (chi² (1) = 2.3, *p* = 0.1297) or increased BMI (chi² (1) = 0.17, *p* = 0.6798).

When all possible combinations of factors were assessed for association with acne vulgaris vs controls, only Vitamin D deficiency occurring with positive tissue transglutaminase antibody was statistically significant (chi² (1) = 5.89, Fisher's exact = 0.041).

Discussion

We found higher prevalence of vitamin D deficiency in patients compared with controls which has been reported previously in smaller series. Vitamin D supplementation has been shown to produce improvements in acne inflammation.² There is consistency at a molecular level where Vitamin D has been demonstrated to reduce serum inflammatory biomarker expression (e.g. IL-6, IL-8, MMP-9) and inhibit P. acnes-induced Th-17 differentiation with decreased IL-17 expression, as well as induce antimicrobial peptide production in sebocytes.¹¹⁻¹³ The association between vitamin D deficiency and anti-TTG is explained by impaired absorption across an inflamed intestinal epithelium.

Acne patients were more than twice as likely to have a non-vegan diet compared with controls. Fundamentally, vegan diets contain no milk and generally lower glycemic index foods. Multiple studies have shown an association between acne and milk intake.⁵⁻⁷ Low glycemic loads, with or without metformin, has been associated with greater reduction in acne lesion counts compared with high loads.^{14,15} Milk

consumption and high glycemic loads have been independently associated with increased levels of serum insulin growth factor-1, offering a possible mechanism.¹⁶⁻¹⁹

We found no significant association between increased BMI and acne vulgaris. Past study results have varied on this connection. Positive correlations have been found between low BMI and reduced risk for acne, and equally, rising BMI and increased risk.^{20,21} Vitamin D deficiency has been shown to occur at higher rates in obesity compared with controls, possibly because of decreased bioavailability from dietary and cutaneous sources due to deposition in body fat compartments.²² Based on our findings, any effect of vitamin D deficiency on acne pathogenesis appears to be independent of BMI.

In recent years, acne vulgaris has been increasingly described as a symptom of various autoimmune inflammatory syndromes (e.g. PAPA, PASH).²³⁻²⁶ Autoinflammatory diseases are signified by presence of an inheritance pattern and dysregulated innate immunity. Pathogenesis is not fully understood but is thought to be mediated primarily by abnormal interleukin-1 signaling.²⁷ This mechanism has received additional support from successful treatment with IL-1 antagonists.²⁸

Celiac disease has been associated with several autoimmune conditions and autoinflammatory diseases overlap with, and may even mimic autoimmune diseases.²⁹ Innate immunity has been shown to play an important role in celiac pathogenesis and may be necessary to trigger the gliadin-specific T-cell response in genetically-predisposed individuals.³⁰ Anti-TTG-positive celiac disease was associated with acne diagnosis in a single study.³¹ Future studies might better explore this association after exclusion of vitamin D deficiency.

Our study is limited by its retrospectivity. We categorized patients simply as either having, or not having a vegan diet. There is a variety of vegan diets featuring different food types. For this reason, breakdown of subtypes of vegan diets might have proved more

enlightening. This study might also be criticized for not confirming acne diagnoses with a skin specialist, however family physicians have demonstrated equivalence with dermatologists for diagnosis of several common skin diagnoses, including acne.³²

Our findings reaffirm vitamin D and dietary factors as potential pathogenic players in clinically-evident acne vulgaris. In addition, we introduce a novel association between anti-tissue transglutaminase antibody and acne, which may act independently or dependently of Vitamin D status. This, along with our other identified associations warrant further investigation prospectively in larger cohorts. We lend credence to acne vulgaris as a highly complex disease with developing pathogenesis.

Materials and methods

We retrospectively screened records from January 2008 to May 2017 at a family medicine clinic for consecutive patients 16 years or older diagnosed with acne vulgaris who had presented primarily about their skin. In this population, BMI and 25-hydroxyvitamin D levels had been recorded within one month of acne diagnosis. Vitamin D testing was performed as part of general health screening. Vitamin D deficiency was defined as <50 nmol/L.

Routine Workup of vitamin D deficiency at our practice included enquiry about patients diet and testing of serum IgA tissue transglutaminase and total IgA antibody levels. Increased BMI was defined as >30 kg/m² and a tissue transglutaminase antibody level >4.0 U/ml was considered positive.

Controls were age- (+/- 12 months) and sex-matched patients seen during the study period without a diagnosis of acne vulgaris presenting for wellness checks. Patients were excluded if they had therapeutic interventions (within 3 months) which might have affected vitamin D levels including bisphosphonates, systemic corticosteroids, vitamin D or calcium supplementation, or IgA deficiency.

Statistical analyses were performed using STATA V14.2. Pearson's chi-squared and Fisher's exact tests were used to statistically analyse the data from the patient and control groups. P values of <0.05 were considered statistically significant.

Internal board review or ethics approval was not sought as all study data had been collected as part of the patients routine care at a private institution and is

owned in its entirety by the second author. Study procedures were carried out in accordance with the Helsinki Declaration of 1975.

References

1. Contassot E, French LE. New insights into acne pathogenesis: Propionibacterium acnes activates the inflammasome. *J Invest Dermatol.* 2014;134(2):310–3. doi:10.1038/jid.2013.505. PMID:24424454.
2. Lim SK, Ha JM, Lee YH, Lee Y, Seo YJ, Kim CD, Lee JH, Im M. Comparison of vitamin D levels in patients with and without acne: A case-control study combined with a randomised controlled trial. *PLoS One.* 2016; 11(8):e01611162. doi:10.1371/journal.pone.01611162. PMID:27560161.
3. Yildizgoren MT, Togrul AK. Preliminary evidence for vitamin D deficiency in nodulocystic acne. *Dermatoendocrinol.* 2015;6:1.e983687.
4. Toosi P, Azizian Z, Yavari H, Fakhim TH, Amini SH, Enamzade R. Serum 25-hydroxy vitamin D levels in patients with acne vulgaris and its association with disease severity. *Clin Cases in Min Bone Metab.* 2015;12(3):238–42.
5. Adebamowo CA, Spiegelman D, Danby FW, Frazier AL, Willett WC, Holmes MD. High school dietary dairy intake and teenage acne. *J Am Acad Dermatol.* 2005;52(2):207. doi:10.1016/j.jaad.2004.08.007. PMID:15692464.
6. Adebamowo CA, Spiegelman D, Berkeley CS, Danby FW, Rockett HH, Colditz GA, Willett WC, Holmes MD. Milk consumption and acne in teenaged boys. *J Am Acad Dermatol.* 2008;58(5):787. doi:10.1016/j.jaad.2007.08.049. PMID:18194824.
7. Adebamowo CA, Spiegelman D, Berkeley CS, Danby FW, Rockett HH, Colditz GA, Willett WC, Holmes MD. Milk consumption and acne in adolescent girls. *Dermatol Online J.* 2006;12(4):1. PMID:17083856.
8. Farrace MG, Picarelli A, Tola MD. Presence of anti-tissue transglutaminase antibodies in inflammatory intestinal disease: An apoptosis associated event? Cell death and differentiation. 2001;8:767–70. doi:10.1038/sj.cdd.4400880. PMID:11464221.
9. Tola MD, Sabbatella L, Anania MC, Viscido A, Caprilli R, Pica R, Paoluzi P, Picarelli A. Author information. Anti-tissue transglutaminase antibodies in inflammatory bowel disease: new evidence. *Clin Chem Lab Med.* 2004;42(10):1092–7. doi:10.1515/CCLM.2004.225. PMID:15552265.
10. Gabriel S, Mihaela I, Angela B, Mariana A, Doru D. Prevalence of IgA antitissue transglutaminase antibodies in children with type 1 diabetes mellitus. *J Clin Lab Anal.* 2011;25(3):156–61. doi:10.1002/jcla.20449. PMID:21567461.
11. Lee WJ, Choi YH, Sohn MY. Expression of inflammatory biomarkers from cultured sebocytes was influenced by treatment with Vitamin D. *Indian J Dermatol.* 2013;58:327. doi:10.4103/0019-5154.113959. PMID:23919024.

12. Agak GW, Qin M, Nobe J, Kim MH, Krutzik SR, Tristan GR, Elashoff D, Garbán HJ, Kim J. Propionibacterium acnes induced IL-17 response in acne vulgaris that is regulated by vitamin A and vitamin D. *J Invest Dermatol.* 2014;134:366–73. doi:10.1038/jid.2013.334. PMID:23924903.
13. Lee WJ, Cha HW, Sohn MY, Lee SJ, Kim do W. Vitamin D increases expression of cathelicidin in cultured sebocytes. *Arch Dermatol Res.* 2012;304:627–32. doi:10.1007/s00403-012-1255-z. PMID:22695798.
14. Smith RN, Mann NJ, Braue A. The effect of a high-protein, low glycemic-load diet versus conventional high glycemic-load diet on biochemical parameters associated with acne vulgaris: a randomized, investigator-masked, controlled trial. *J Am Acad Dermatol.* 2007;57(2):247. doi:10.1016/j.jaad.2007.01.046. PMID:17448569.
15. Fabbrocini G, Izzo R, Faggiano A, Del Prete M, Donnarumma M, Marasca C, Marciello F, Savastano R, Montfrecola G, Colao A. Low glycaemic diet and metformin therapy: a new approach in male subjects with acne resistant to common treatments. *Clin Exp Dermatol.* 2016;41(1):38. doi:10.1111/ced.12673. PMID:26053680.
16. Holmes MD, Pollak MN, Willet WC, Hankinson SE. Dietary correlates of plasma insulin-like growth factor I and insulin-like growth factor binding protein 3 concentrations. *Cancer Epidemiol Biomarkers Prev.* 2002;11(9):852. PMID:12223429.
17. Runchey SS, Pollak MN, Valsta LM. Glycemic load effect on fasting and post prandial serum glucose, insulin, IGF-1 and IGFBP-3 in a randomised, controlled feeding study. *Eur J Clin Nutr.* 2012;66(10):1146–52. doi:10.1038/ejcn.2012.107. PMID:22892437.
18. Aizawa H, Niimura M. Elevated serum insulin-like growth factor-1 (IGF-1) levels in women with postadolescent acne. *J Dermatol.* 1995 Apr;22(4):249–52.
19. Vora S, O'Vhal A, Jerajani H, Nair N, Chakraborty A. Correlation of facial sebum to serum insulin-like growth factor-1 in patients with acne. *Br J Dermatol.* 2008 Sep;159(4):990–1. doi:10.1111/j.1346-8138.1995.tb03381.x.
20. Di Landro A, Gazzaniga S, Parazzini F, Ingordo V, Cusano F, Atzori L, Cutri FT, Musumeci ML, Zinetti C, Pezzarossa E, et al. Family history, body mass index, selected dietary factors, menstrual history, and risk of moderate to severe acne in adolescents and young adults. *J Am Acad Dermatol.* 2012;67(6):1129. doi:10.1016/j.jaad.2012.02.018. PMID:22386050.
21. Halvorsen JA, Vleugels RA, Bjerntness E, Lien L. A population-based study of acne and body mass index in adolescents. *Arch Dermatol.* 2012 Jan;148(1):131–2. doi:10.1001/archderm.148.1.131.
22. Worstmas J, Matsuoka LY, Chen TC, Lu Z, Holick MF. Decreased bioavailability of vitamin D in obesity. *Am J Clin Nutr.* 2000;72(3):690–3. doi:10.1093/ajcn/72.3.690. PMID:10966885.
23. Lindor NM, Arsenault TM, Solomon H, Seidman CE, McEvoy MT. A new autosomal dominant disorder of pyogenic sterile arthritis, pyoderma gangrenosum, and acne: PAPA Syndrome. *Mayo Clin Proc.* 1997;72:611–5. doi:10.1016/S0025-6196(11)63565-9. PMID:9212761.
24. Braun-Falco M, Koverysky O, Lohse P. Pyoderma gangrenosum, acne and suppurative hidradenitis (PASH) – a new autoimmune syndrome distinct form PAPA syndrome. *J Am Acad Dermatol.* 2012;66(3):409–15. doi:10.1016/j.jaad.2010.12.025. PMID:21745697.
25. Marzano AV, Trevisan V, Gattorno M, Ceccherini I, De Simone C, Crosti C. Pyogenic arthritis, pyoderma gangrenosum, acne, and hidradenitis suppurativa (PAPASH): A new autoimmune syndrome associated with a novel mutation of the PSTPIP1 gene. *JAMA Dermatol.* 2013;149(6):762–4. doi:10.1001/jamadermatol.2013.2907. PMID:23571383.
26. Saraceno R, Banino G, Chiricozzi A, Zangrilli A, Chimenti S. PsAPASH: A new syndrome associated with hidradenitis suppurativa with response to tumor necrosis factor inhibition. *J Am Acad Dermatol.* 2015;72(1):e42–4. doi:10.1016/j.jaad.2014.10.002. PMID:25497954.
27. Vinkel C, Thomsen SF. Autoinflammatory syndromes associated with hidradenitis suppurativa and/or acne. *Int J Dermatol.* 2017;56(8):811–8. doi:10.1111/ijd.13603. PMID:28345207.
28. Hoffman HM. Therapy of autoinflammatory syndromes. *J Allergy Clin Immunol.* 2015;124(6):1129–40. doi:10.1016/j.jaci.2009.11.001.
29. Kiss MH, Magalhães CS. Autoinflammatory diseases: Mimics of autoimmunity or part of its spectrum? Case presentation. *J Clin Immunol.* 2008;28:84–9. doi:10.1007/s10875-008-9179-2.
30. Mauri L, Ciacci C, Ricciardelli I, Yacca L, Raia V, Auricchio S, Picard J, Osman M, Quarantino S, Londei M. Association between innate response to gliadin and activation of pathogenic T cells in coeliac disease. *Lancet.* 2003;362(9377):30. doi:10.1016/S0140-6736(03)13803-2. PMID:12853196.
31. Lebwohl B, Sundstrom A, Jabri B, Kupfer SS, Green PH, Ludvigsson JF. Isotretinoin use and celiac disease: a population-based cross-sectional study. *Am J Clin Dermatol.* 2014;16(6):537–42. doi:10.1007/s40257-014-0090-8.
32. Castillo-Arenas E, Garrido V, Serrano-Ortega S. Skin conditions in primary care: An analysis of referral demand. *Acta Dermosifilogr.* 2014;105:271–5. doi:10.1016/j.ad.2013.10.003.